

The Bancroft Library

University of California • Berkeley

THE THEODORE P. HILL COLLECTION
of
EARLY AMERICAN MATHEMATICS BOOKS

Digitized by the Internet Archive
in 2008 with funding from
Microsoft Corporation



A
COURSE
OF
MATHEMATICS,
FOR THE
USE OF ACADEMIES,
AS WELL AS
PRIVATE TUITION.

IN TWO VOLUMES.

BY
CHARLES HUTTON, LL.D. F. R. S.
Late Professor of Mathematics in the Royal Military Academy.

THE THIRD AMERICAN EDITION:
From the Fifth, Sixth, and Seventh London Editions.

REVISED, CORRECTED, AND IMPROVED.

TO WHICH IS ADDED

An Elementary Essay on Descriptive Geometry.

BY ROBERT ADRAIN, LL. D. F. A. P. S. F. A. A. S. &c.
And Professor of Mathematics and Natural Philosophy, in Columbia
College, New-York.

VOL. I.

NEW-YORK:

PUBLISHED BY SAMUEL CAMPBELL & SON, EVERT DUYCKINCK, T. & J. SWORDS,
R. M'DERMUT, T. A. RONALDS, COLLINS & HANNAY, AND GEORGE LONG.
George Long, Printer, No. 71, Pearl-Street.

1822.

Southern District of New-York, ss.

BE IT REMEMBERED, That on the twenty-seventh day of February, in the forty-sixth year of the Independence of the United States of America, *George Long*, of the said district, hath deposited in this office the title of a book, the right whereof he claims as proprietor, in the words following, to wit:

“ A Course of Mathematics, for the Use of Academies, as well as Private Tuition. In two Volumes. By Charles Hutton, LL.D. F. R. S. Late Professor of Mathematics in the Royal Military Academy. The Third American Edition. From the Fifth, Sixth, and Seventh London Editions. Revised, corrected, and improved. To which is added, An Elementary Essay on Descriptive Geometry, by Robert Adrain, LL.D. F. A. P. S. F. A. A. S. &c. and Professor of Mathematics and Natural Philosophy, in Columbia College, New-York.”

In conformity to the act of Congress of the United States, entitled, “ An act for the encouragement of learning, by securing the copies of maps, charts, and books, to the authors and proprietors of such copies, during the time therein mentioned.” And also to an act, entitled, “ An act, supplementary to an act, entitled, An act for the encouragement of learning, by securing the copies of maps, charts, and books, to the authors and proprietors of such copies, during the times therein mentioned, and extending the benefits thereof to the arts of designing, engraving, and etching historical and other prints.”

JAMES DILL,
Clerk of the Southern District of New-York.

P R E F A C E.

A SHORT and Easy Course of the Mathematical Sciences has long been considered as a desideratum for the use of Students in the different schools of education : one that should hold a middle rank between the mere voluminous and bulky collections of this kind, and the mere abstract and brief common-place forms, of principles and memorandums.

For long experience, in all Seminaries of Learning, has shown, that such a work was very much wanted, and would prove a great and general benefit ; as, for want of it, recourse has always been obliged to be had to a number of other books by different authors ; selecting a part from one and a part from another as seemed most suitable to the purpose in hand, and rejecting the other parts—a practice which occasioned much expense and trouble, in procuring and using such a number of odd volumes, of various forms and modes of composition ; besides wanting the benefit of uniformity and reference, which are found in a regular series of composition.

To remove these inconveniences, the Author of the present work has been induced from time to time, to compose various parts of this Course of Mathematics ; which the experience of many years' use in the Academy has enabled him to adapt and improve to the most useful form and quantity for the benefit of instruction there. And, to render that benefit more eminent and lasting, the Master General of the Ordnance has been pleased to give it its present form, by ordering it to be enlarged and printed, for the use of the Royal Military Academy.

As this work has been composed expressly with the intention of adapting it to the purposes of academical education, it is not designed to hold out the expectation of an entire new mass of inventions and discoveries : but rather to collect and arrange the most useful known principles of mathematics, disposed in a convenient practical form, demonstrated in a plain and concise way, and illustrated with suitable examples, rejecting whatever seemed to be matters of mere curiosity, and retaining only such parts and branches, as have a direct tendency and application to some useful purpose in life or profession.

It is however expected that much that is new will be found in many parts of these volumes ; as well in the matter, as in the arrangement and manner of demonstration, throughout the whole work, especially in the geometry, which is rendered much more easy and simple than heretofore ; and in the conic sections

sections, which are here treated in a manner at once new, easy, and natural ; so much so indeed that all the propositions and their demonstrations in the ellipsis, are the very same, word for word, as those in the hyperbola, using only, in a very few places, the word *sum*, for the word *difference* : also in many of the mechanical and philosophical parts which follow in the second volume. In the conic sections too, it may be observed, that the first theorem of each section only is proved from the cone itself, and all the rest of the theorems are deduced from the first, or from each other, in a very plain and simple manner.

Besides renewing most of the rules, and introducing every where new examples, this edition is much enlarged in several places ; particularly by extending the tables of squares and cubes, square roots and cube roots, to 1000 numbers, which will be found of great use in many calculations ; also by the tables of logarithms, sines, and tangents, at the end of the second volume ; by the addition of Cardan's rules for resolving cubic equations ; with tables and rules for annuities ; and many other improvements in different parts of the work.

Though the several parts of this course of mathematics are ranged in the order naturally required by such elements, yet students may omit any of the particulars that may be thought the least necessary to their several purposes ; or they may study and learn various parts in a different order from their present arrangement in the book, at the discretion of the tutor. So, for instance, all the notes at the foot of the pages may be omitted, as well as many of the rules ; particularly the 1st or Common Rule for the Cube Root, p. 85, may well be omitted, being more tedious than useful. Also the chapters on Surds and Infinite Series, in the Algebra ; or these might be learned after Simple Equations. Also Compound Interest and Annuities at the end of the Algebra. Also any part of the Geometry, in vol. 1 ; any of the branches in vol. 2, at the discretion of the preceptor. And, in any of the parts, he may omit some of the examples, or he may give more than are printed in the book ; or he may very profitably vary or change them by altering the numbers occasionally.—As to the quantity of writing ; the author would recommend, that the student copy out into his fair book no more than the chief rules which he is directed to learn off by rote, with the work of one example only to each rule, set down at full length ; omitting to set down the work of all the other examples, how many soever he may be directed to work out upon his slate or waste paper.—In short, a great deal of the business, as to the quantity, and order, and manner, must depend on the judgment of the discreet and prudent tutor or director.

[Dr.

THE beneficial improvements lately made, and still making in the plan of the scientific education of the Cadets, in the Royal Military Academy at Woolwich, having rendered a further extension of the Mathematical Course adviseable, I was honoured with the orders of his Lordship the Master General of the Ordnance, to prepare a third volume, in addition to the two former volumes of the Course. to contain such additions to some of the subjects before treated of in those two volumes, with such other new branches of military science, as might appear best adapted to promote the ends of this important institution. From my advanced age, and the precarious state of my health, I was desirous of declining such a task, and pleaded my doubts of being able, in such a state, to answer satisfactorily his lordship's wishes. This difficulty however was obviated by the reply, that, to preserve a uniformity between the former and the additional parts of the Course, it was requisite that I should undertake the direction of the arrangement, and compose such parts of the work as might be found convenient, or as related to topics in which I had made experiments or improvements; and for the rest, I might take to my assistance the aid of any other person I might think proper. With this kind indulgence, being encouraged to exert my best endeavours, I immediately announced my wish to request the assistance of Dr. Gregory of the Royal Military Academy, than whom, both for his extensive scientific knowledge, and his long experience, I know of no person more fit to be associated in the due performance of such a task. Accordingly, this volume is to be considered as the joint composition of that gentleman and myself, having each of us taken and prepared, in nearly equal portions, separate chapters and branches of the work, being such as in the compass of this volume, with the advice and assistance of the Lieut. Governor, were deemed among the most useful additional subjects for the purposes of the education established in the Academy.

The several parts of the work, and their arrangement, are as follow.—In the first chapter are contained all the propositions of the course of *Conic Sections*, first printed for the use of the Academy in the year 1787, which remained, after those that were selected for the second volume of this Course: to which is added a tract on the algebraic equations of the several conic sections, serving as a brief introduction to the algebraic properties of curve lines.

The 2d chapter contains a short geometrical treatise on the elements of *Isoperimetry* and the *maxima and minima of surfaces and solids*; in which several propositions usually investigated by fluxionary processes are effected geometrically; and in which, indeed, the principal results deduced by Thos. Simpson, Horsley, Legendre, and Lhuillier, are thrown into the compass of one short tract.

The 3d and 4th chapters exhibit a concise but comprehensive view of the *trigonometrical analysis*, or that in which the chief theorems of Plane and Spherical Trigonometry are deduced algebraically by means of what is commonly denominated the *Arithmetic of Sines*. A comparison of the modes of investigation adopted in these chapters, and those pursued in that part of the second volume of this course which is devoted to Trigonometry, will enable a student to trace the relative advantages of the algebraical and geometrical methods of treating this useful branch of science. The fourth chapter includes also a disquisition on the nature and measure of *solid angles*, in which the theory of that peculiar class of geometrical magnitudes is so represented, as to render their mutual comparison (a thing hitherto supposed impossible, except in one or two very obvious cases) a matter of perfect ease and simplicity.

Chapter the fifth relates to Geodesic Operations, and that more extensive kind of *Trigonometrical Surveying* which is employed with a view to determine the geographical situation of places, the magnitude of kingdoms, and the figure of the earth. This chapter is divided into two sections; in the first of which is presented a general account of this kind of surveying; and in the second, solutions of the most important problems connected with these operations. This portion of the volume it is hoped will be found highly useful; as there is no work which contains a concise and connected account of this kind of surveying and its dependent problems; and it cannot fail to be interesting to those who know how much honour redounds to this country from the great skill, accuracy, and judgment, with which the trigonometrical survey of England has long been carried on.

In the 6th and 7th chapters are developed the principles of *Polygonometry*, and those which relate to the *Division of lands* and other surfaces, both by geometrical construction and by computation.

The 8th chapter contains a view of the nature and solution of *equations* in general, with a selection of the best rules for equations of different degrees. Chapter the 9th is devoted to

the

the nature and properties of *curves*, and the *construction of equations*. These chapters are manifestly connected, and show how the mutual relations subsisting between equations of different degrees, and curves of various orders, serve for the reciprocal illustration of the properties of both.

In the 10th chapter the subjects of *Fluents* and *Fluxional equations* are concisely treated. The various forms of Fluents comprised in the useful table of them in the 2d volume, are investigated : and several other rules are given ; such as it is believed will tend much to facilitate the progress of students in this interesting department of science, especially those which relate to the mode of finding fluents by continuation.

The 11th chapter contains solutions of the most useful problems concerning the *maximum effects of machines in motion* ; and develops those principles which should constantly be kept in view by those who would labour beneficially for the improvement of machines.

In the 12th chapter will be found the theory of the *pressure of earth and fluids* against walls and fortifications ; and the theory which leads to the best construction of *powder magazines* with equilibrated roofs.

The 13th chapter is devoted to that highly interesting subject, as well to the philosopher as to military men, the *theory and practice of gunnery*. Many of the difficulties attending this abstruse enquiry are surmounted by assuming the results of accurate experiments, as to the resistance experienced by bodies moving through the air, as the basis of the computations. Several of the most useful problems are solved by means of this expedient, with a facility scarcely to be expected, and with an accuracy far beyond our most sanguine expectations.

The 14th and last chapter contains a promiscuous but extensive collection of problems in *statics, dynamics, hydrostatics, hydraulics, projectiles, &c. &c.* ; serving at once to exercise the pupil in the various branches of mathematics comprised in the Course, to demonstrate their utility especially to those devoted to the military profession, to excite a thirst for knowledge, and in several important respects, to gratify it. This volume being professedly supplementary to the preceding two volumes of the Course, may best be used in tuition by a kind of mutual incorporation of its contents with those of the second volume. The method of effecting this will, of course, vary according to circumstances, and the precise employments for which the pupils are destined : but in general it is presumed the following may be advantageously adopted. Let the first seven chapters be taught, immediately after the
Conic

Conic Sections in the 2d volume. Then let the substance of the 2d volume succeed, as far as the Practical Exercises on Natural Philosophy, inclusive. Let the 8th and 9th chapters in this 3d vol. precede the treatise on Fluxions in the 2d ; and when the pupil has been taught the part relating to *fluents* in that treatise, let him immediately be conducted through the 10th chapter of the 2d volume. After he has gone over the remainder of the Fluxions with the applications to tangents, radii of curvature, rectifications, quadratures, &c. the 11th and 12th chapters of the 3d vol. should be taught. The problems in the 13th and 14th chapters must be blended with the practical exercises at the end of the 2d volume, in such manner as shall be found best suited to the capacity of the student, and best calculated to ensure his thorough comprehension of the several curious problems contained in those portions of the work.

In the composition of this 3d volume, as well as in that of the preceding parts of the Course, the great object kept constantly in view has been *utility*, especially to gentlemen intended for the Military Profession. To this end, all such investigations as might serve merely to display ingenuity or talent, without any regard to practical benefit, have been carefully excluded. The student has put into his hands the two powerful instruments of the ancient and the modern or sublime geometry ; he is taught the use of both, and their relative advantages are so exhibited as to guard him, it is hoped, from any undue and exclusive preference for either. Much novelty of matter is not to be expected in a work like this ; though, considering its magnitude, and the frequency with which several of the subjects have been discussed, a candid reader will not, perhaps, be entirely disappointed in this respect. Perspicuity and condensation have been uniformly aimed at through the performance ; and a small clear type, with a full page, have been chosen for the introduction of a large quantity of matter.

A candid public will accept as an apology for any slight disorder or irregularity that may appear in the composition and arrangement of this Course, the circumstance of the different volumes having been prepared at widely distant times, and with gradually expanding views. But, on the whole, I trust it will be found that, with the assistance of my friend and coadjutor in this supplementary volume, I have now produced a Course of Mathematics, in which a great variety of useful subjects are introduced, and treated with greater perspicuity and correctness, than in any three volumes of equal size in any language.

CHA. HUTTON.

PREFACE,

BY THE AMERICAN EDITOR.

THE last English edition of *Hutton's Course of Mathematics*, in three volumes octavo, may be considered as one of the best systems of Mathematics in the English language. Its great excellence consists in the judicious selection made by the authors of the work, who have constantly aimed at such things as are most necessary in the useful arts of life. To this may be added the easy and perspicuous manner in which the subject is treated—a quality of primary importance in a treatise intended for beginners, and containing the elements of science.

The third volume of the English edition having been but lately published, is scarcely known at present in this country—it is but justice to its excellent authors to state, that they have collected in it a great number of the most interesting subjects in Analytical and Mechanical Science. Analytical Trigonometry, Plane and Spherical, Trigonometrical Surveying, Maxima and Minima of Geometrical Quantities, Motion of Machines and their Maximum Effects, Practical Gunnery, &c. are among the most important subjects in Mathematics, and are discussed in the volume just mentioned in such a manner as not only to prove highly useful to pupils, but also to such as are engaged in various departments of Practical Science.

As the work, after the publication of the third volume, embraced most subjects of curiosity or utility in Mathematics, it has been thought unnecessary to enlarge its size by much additional matter. The present edition however, differs in
several

several respects from the last English one ; and it is presumed, that this difference will be found to consist of improvements. These are principally as follows :

In the first place, it was thought advisable to publish the work in two volumes instead of three ; the two volumes being still of a convenient size for the use of students.

Secondly, a new arrangement of various parts of the work has been adopted. Several parts of the third volume of the English edition treated of subjects already discussed in the preceding volumes ; in such cases, when it was practicable, the additions in the third volume have been properly incorporated with the corresponding subjects that preceded them ; and, in general, such a disposition of the various departments of the work has been made as seemed best calculated to promote the improvement of the pupil, and exhibit the respective places of the various branches in the scale of science.

In the third place, several notes have been added ; and numerous corrections have been made in various places of the work : it were tedious and unnecessary to enumerate all these at present ; it may suffice to remark the few following :

In pages 58, 59, vol. 1, a note is added on the reduction of fractions to the least common denominator ; and for common cases an easier rule is given, than has been before presented to the public.

In page 169. vol. 1, a useful note is added respecting the degree of accuracy resulting from the application of logarithms. This note will appear the more necessary, when we observe such oversights committed by authors of experience.

In several places of the last or seventh London edition, the corrections made in the first American edition have been adopted. The definition of Surds which had been improperly given in the fifth and sixth London editions, is now correctly in the seventh ; agreeably to the mode prescribed in the first American edition. This erroneous definition of Surds is still retained in the large Algebra of Bonnycastle, published

lished in London, in 1820. The true definition is given in the small work of the same author, by the editor, Mr. Ryan, in the New-York edition of 1822.

The erroneous computation of the value x in the equation $x^x = 100$, which was pointed out and corrected in the first American edition, is expunged from the seventh London edition. The solution of this problem was subject to the same error in both the treatises of Algebra by Bonnycastle. The American edition of 1822 is correct, but the larger Algebra, published in London, in 1820, still retains the error.

In the second volume, page 24, American edition, a very simple solution was given to the problem in the sixth example. This problem was solved by a cubic in the sixth London edition; and in the seventh, the solution is reduced from a cubic to a quadratic; but notwithstanding this improvement of the solution, it is still inferior to that given in the first American edition.

There are in the last London edition several errors continued from the sixth edition, which had been corrected in the first American edition. Among these we may notice the demonstration given to the third theorem in Spherics. The demonstration is founded on the assumption, that an angle of a spherical triangle is greater than the angle contained by the chords of the sides containing the spherical angle.

See on this subject the note page 555, vol. II.

In the mensuration, page 411, vol. 1, a remark is added respecting the magnitude of the earth. Dr. Hutton has commonly used a diameter of $7957\frac{3}{4}$ English miles, merely because it gives the round number 25,000 for the circumference: in a few places he has used a diameter of 7930. Having some years ago discovered the proper method of ascertaining the most probable magnitude and figure of the earth, from the admeasurement of several degrees of the meridian, I found the ratio of the axis to the equatorial diameter, to be as 320 to 321, and the diameter, when the earth is considered as a globe, to be 7918.7 English miles.

In the additions immediately preceding the Table of Logarithms in the second volume, a new method is given for ascertaining

certaining the vibrations of a variable pendulum. This problem was solved by Dr. Hutton, in his *Select Exercises*, 1787, and he has given the same solution in the present work, see page 537, vol. 2. The method used by the Doctor appears to me to be erroneous; but in order that such as would judge for themselves on this abstruse question, may have a fair opportunity of deciding between us, the Doctor's solution is given as well as my own.

It may be proper to observe, with respect to the new solution, as well as Dr. Hutton's that the resulting formula does not show the relation between the time and any number of vibrations *actually* performed; but merely gives the limit to which this relation approaches, when the horizontal velocity is indefinitely diminished. If therefore we would use the new formula as an *approximation* in very small finite vibrations, the times must not be extended without limitation.

Besides the numerous corrections in this third American edition, there is added to the second volume an elementary treatise on *Descriptive Geometry*, in which the principles and fundamental problems are given in a simple and easy manner, with a select number of useful applications, in Spherics, Conics, Sections, &c.

ROBERT ADRAIN.

Columbia College, New-York.
May 1, 1822.

CONTENTS

OF VOLUME I.

	Page
GENERAL Principles	1
ARITHMETIC.	
<i>Notation and Numeration</i>	4
<i>Roman Notation</i>	7
<i>Addition</i>	8
<i>Subtraction</i>	11
<i>Multiplication</i>	13
<i>Division</i>	18
<i>Reduction</i>	23
<i>Compound Addition</i>	32
— <i>Subtraction</i>	36
— <i>Multiplication</i>	38
— <i>Division</i>	41
<i>Golden Rule, or Rule of Three</i>	44
<i>Compound Proportion</i>	49
<i>Vulgar Fractions</i>	51
<i>Reduction of Vulgar Fractions</i>	52
<i>Addition of Vulgar Fractions</i>	61
	<i>Subtraction</i>

	Page
<i>Subtraction of Vulgar Fractions</i>	62
<i>Multiplication of Vulgar Fractions</i>	63
<i>Division of Vulgar Fractions</i>	64
<i>Rule of Three in Vulgar Fractions</i>	65
<i>Decimal Fractions</i>	66
<i>Addition of Decimals</i>	67
<i>Subtraction of Decimals</i>	68
<i>Multiplication of Decimals</i>	68
<i>Division of Decimals</i>	70
<i>Reduction of Decimals</i>	73
<i>Rule of three in Decimals</i>	76
<i>Duodecimals</i>	77
<i>Involution</i>	78
<i>Evolution</i>	80
<i>To extract the Square Root</i>	81
<i>To extract the Cube Root</i>	85
<i>To extract any Root whatever</i>	88
<i>Table of Squares, Cubes, Square-Roots and Cube-Roots</i>	90
<i>Ratios, Proportions, and Progressions</i>	110
<i>Arithmetical Progression</i>	111
<i>Geometrical Progression</i>	116
<i>Musical Proportion</i>	119
<i>Fellowship, or Partnership</i>	ib.
<i>Single Fellowship</i>	120
<i>Double Fellowship</i>	122
<i>Simple Interest</i>	124
<i>Compound Interest</i>	127
<i>Alligation Medial</i>	129
<i>Alligation Alternate</i>	131
<i>Single Position</i>	135
<i>Double Position</i>	137
<i>Permutations and Combinations</i>	140
<i>Practical Questions</i>	150

LOGARITHMS.

<i>Of Logarithms</i>	155
<i>To</i>	

CONTENTS.

xv

	Page
<i>To compute Logarithms</i>	159
<i>Description and use of the Table of Logarithms</i>	163
<i>Multiplication by Logarithms</i>	167
<i>Division by Logarithms</i>	169
<i>Involution by Logarithms</i>	169
<i>Evolution by Logarithms</i>	170

ALGEBRA.

<i>Definitions and Notation</i>	171
<i>Addition</i>	175
<i>Subtraction</i>	180
<i>Multiplication</i>	181
<i>Division</i>	184
<i>Algebraic Fractions</i>	188
<i>Involution</i>	199
<i>Evolution</i>	202
<i>Surds</i>	206
<i>Infinite Series</i>	213
<i>Arithmetical Proportion</i>	218
<i>Piling of Balls</i>	223
<i>Geometrical Proportion</i>	228
<i>Simple Equations</i>	230
<i>Quadratic Equations</i>	249
<i>Resolution of Cubic and Higher Equations</i>	257
<i>Simple Interest</i>	266
<i>Compound Interest</i>	267
<i>Annuities</i>	270

GEOMETRY.

<i>Definitions</i>	275
<i>Axioms—Theorems</i>	281
<i>Of Ratios and Proportions</i>	319
<i>Of Planes and Solids—Definitions</i>	336
<i>Problems</i>	353
<i>Applications of Algebra to Geometry</i>	369
<i>Plane Trigonometry</i>	377

Heights

	Page
<i>Heights and Distances</i>	398
<i>Mensuration of Planes</i>	402
<i>Mensuration of Solids</i>	419
<i>Land Surveying</i>	429
<i>Artificers' Work</i>	457
<i>Timber Measuring</i>	466
<i>Conic Sections</i>	469
<i>Of the Ellipse</i>	473
<i>Of the Hyperbola</i>	491
<i>Of the Parabola</i>	514
<i>Of the Conic Sections as expressed by Algebraic equations, called the</i> <i>Equations of the Curve</i>	532
<i>Elements of Isoperimetry</i>	535
<i>Problems relative to the Divison of Surfaces</i>	558
<i>Construction of Geometrical Problems</i>	571
<i>Practical Exercises in Mensuration</i>	575

A
COURSE
OF
MATHEMATICS, &c.

GENERAL PRINCIPLES.

1. **Q**UANTITY, or **M**MAGNITUDE, is any thing that will admit of increase or decrease ; or that is capable of any sort of calculation or mensuration : such as numbers, lines, space, time, motion, weight.

2. **M**ATHEMATICS is the science which treats of all kinds of quantity whatever, that can be numbered or measured.— That part which treats of numbering is called *Arithmetic* ; and that which concerns measuring, or figured extension, is called *Geometry*.—These two, which are conversant about multitude and magnitude, being the foundation of all the other parts, are called *Pure* or *Abstract Mathematics* ; because they investigate and demonstrate the properties of abstract numbers and magnitudes of all sorts. And when these two parts are applied to particular or practical subjects, they constitute the branches or parts called *Mixed Mathematics*.— Mathematics is also distinguished into *Speculative* and *Practical* : viz. *Speculative*, when it is concerned in discovering properties and relations ; and *Practical*, when applied to practice and real use concerning physical objects.

3. In Mathematics are several general terms or principles ; such as Definitions, Axioms, Propositions, Theorems, Problems, Lemmas, Corollaries, Scholiums, &c.

4. *A Definition* is the explication of any term or word in a science ; showing the sense and meaning in which the term is employed.—Every Definition ought to be clear, and expressed in words that are common and perfectly well understood.

5. *A Proposition* is something proposed to be proved, or something required to be done ; and is accordingly either a Theorem or a Problem.

6. *A Theorem* is a demonstrative proposition ; in which some property is asserted, and the truth of it required to be proved. Thus, when it is said that, The sum of the three angles of any triangle is equal to two right angles, this is a Theorem, the truth of which is demonstrated by Geometry.—A set or collection of such Theorems constitutes a *Theory*.

7. *A Problem* is a proposition or a question requiring something to be done ; either to investigate some truth or property, or to perform some operation. As, to find out the quantity or sum of all the three angles of any triangle, or to draw one line perpendicular to another.—*A Limited Problem* is that which has but one answer. *An Unlimited Problem* is that which has innumerable answers. And a *Determinate Problem* is that which has a certain number of answers.

8. *Solution* of a Problem, is the method of finding the answer. *A Numerical* or *Numeral Solution*, is the answer given in numbers. *A Geometrical Solution*, is the answer given by the principles of Geometry. And a *Mechanical Solution*, is one which is gained by trials.

9. *A Lemna* is a preparatory proposition, laid down in order to shorten the demonstration of the main proposition which follows it.

10. *A Corollary*, or *Consectary*, is a consequence drawn immediately from some proposition or other premises.

11. *A Scholium* is a remark or observation made on some foregoing proposition or premises.

12. *An Axiom*, or *Maxim*, is a self-evident proposition ; requiring no formal demonstration to prove the truth of it ; but is received and assented to as soon as mentioned. Such as, The whole of any thing is greater than a part of it ; or, The whole is equal to all its parts taken together ; or, Two quantities that are each of them equal to a third quantity, are equal to each other

13. *A Postulate, or Petition*, is something required to be done, which is so easy and evident that no person will hesitate to allow it.

14. *An hypothesis* is a supposition assumed to be true, in order to argue from, or to found upon it, the reasoning and demonstration of some proposition.

15. *Demonstration* is the collecting the several arguments and proofs, and laying them together in proper order, to show the truth of the proposition under consideration.

16. *A Direct, Positive, or Affirmative Demonstration*, is that which concludes with the direct and certain proof of the proposition in hand.—This kind of Demonstration is most satisfactory to the mind; for which reason it is called sometimes an *Ostensive Demonstration*.

17. *An Indirect, or Negative Demonstration*, is that which shows a proposition to be true, by proving that some absurdity would necessarily follow if the proposition advanced were false. This is also sometimes called *Reductio ad Absurdum*; because it shows the absurdity and falsehood of all suppositions contrary to that contained in the proposition.

18. *Method* is the art of disposing a train of arguments in a proper order, to investigate either the truth or falsity of a proposition, or to demonstrate it to others when it has been found out.—This is either Analytical or Synthetical.

19. *Analysis*, or the *Analytic Method* in Geometry, is the art or mode of finding out the truth of a proposition, by first supposing the thing to be done, and then reasoning back, step by step, till we arrive at some known truth.—This is also called the *Method of Invention, or Resolution*; and is that which is commonly used in Algebra.

20. *Synthesis*, or the *Synthetic Method*, is the searching out truth, by first laying down some simple and easy principles, and pursuing the consequences flowing from them till we arrive at the conclusion.—This is also called the *Method of Composition*; and is the reverse of the Analytic method, as this proceeds from known principles to an unknown conclusion; while the other goes in a retrograde order, from the thing sought, considered as if it were true, to some known principle or fact. And therefore, when any truth has been found out by the Analytic method, it may be demonstrated by a process in the contrary order, by Synthesis.

ARITHMETIC.

ARITHMETIC is the art or science of numbering ; being that branch of Mathematics which treats of the nature and properties of numbers.—When it treats of whole numbers, it is called *Vulgar*, or *Common Arithmetic* ; but when of broken numbers, or parts of numbers, it is called *Fractions*.

Unity, or an *Unit*, is that by which every thing is called one ; being the beginning of number ; as, one man, one ball, one gun.

Number is either simply one, or a compound of several units ; as, one man, three men, ten men.

An *Integer*, or *Whole Number*, is some certain precise quantity of units ; as, one, three, ten.—These are so called as distinguished from *Fractions*, which are broken numbers, or parts of numbers ; as, one-half, two-thirds, or three-fourths.

 NOTATION AND NUMERATION.

NOTATION, or NUMERATION, teaches to denote or express any proposed number either by words or characters ; or to read and write down any sum or number.

The numbers in Arithmetic are expressed by the following ten digits, or Arabic numeral figures, which were introduced into Europe by the Moors, about eight or nine hundred years since ; viz. 1 one, 2 two, 3 three, 4 four, 5 five, 6 six, 7 seven, 8 eight, 9 nine, 0 cipher, or nothing. These characters or figures were formerly all called by the general name of *Ciphers* ; whence it came to pass that the art of Arithmetic was then often called *Ciphering*. Also the first nine are called *Significant Figures*, as distinguished from the cipher, which is of itself quite insignificant.

Besides this value of those figures, they have also another which depends on the place they stand in when joined together ; as in the following table :

Units

&c.	Hundreds of Millions	Tens of Millions	Millions	Hundreds of Thousands	Tens of Thousands	Thousands	Hundreds	Tens	Units
0	0	0	0	0	0	0	0	0	0
			1	1	1	1	1	1	1
			2	2	2	2	2	2	2
			3	3	3	3	3	3	3
			4	4	4	4	4	4	4
			5	5	5	5	5	5	5
			6	6	6	6	6	6	6
			7	7	7	7	7	7	7
			8	8	8	8	8	8	8
			9	9	9	9	9	9	9

Here any figure in the first place, reckoning from right to left, denotes only its own simple value ; but that in the second place, denotes ten times its simple value ; and that in the third place, a hundred times its simple value ; and so on : the value of any figure, in each successive place, being always ten times its former value.

Thus, in the number 1796, the 6 in the first place denotes only six units, or simply six ; 9 in the second place signifies nine tens, or ninety ; 7 in the third place, seven hundred ; and the 1 in the fourth place, one thousand ; so that the whole number is read thus, one thousand seven hundred and ninety-six.

As to the cipher, 0, though it signify nothing of itself, yet being joined on the right-hand side to other figures, it increases their value in the same ten-fold proportion : thus, 5 signifies only five ; but 50 denotes 5 tens, or fifty ; and 500 is five hundred ; and so on.

For the more easily reading of large numbers, they are divided into periods and half-periods, each half-period consisting of three figures ; the name of the first period being units ; of the second, millions ; of the third, millions of millions, or bi-millions, contracted to billions : of the fourth, millions of millions of millions, or tri-millions, contracted to trillions, and so on. Also the first part of any period is so many units of it, and the latter part so many thousands.

The following Table contains a summary of the whole doctrine.

Periods.	Quadrill; Trillions; Billions; Millions; Units.									
Half-per.	th.	un.	th.	un.	th.	un.	th.	un.	th.	un.
Figures.	123,456;789,098;765,432;101,234;567,890.									

NUMERATION is the reading of any number in words that is proposed or set down in figures; which will be easily done by help of the following rule, deduced from the foregoing tables and observations—viz.

Divide the figures in the proposed number, as in the summary above, into periods and half periods; then begin at the left-hand side, and read the figures with the names set to them in the two foregoing tables.

EXAMPLES.

Express in words the following numbers; viz.

34	15080	13405670
96	72003	47050023
180	109026	309025600
304	483500	4723507689
6134	2500639	274856390000
9028	7523000	6578600307024

NOTATION is the setting down in figures any number proposed in words; which is done by setting down the figures instead of the words or names belonging to them in the summary above; supplying the vacant places with ciphers where any words do not occur.

EXAMPLES.

Set down in figures the following numbers:

Fifty-seven.

Two hundred eighty six.

Nine thousand two hundred and ten.

Twenty-seven thousand five hundred and ninety-four.

Six hundred and forty thousand, four hundred and eighty one.

Three millions, two hundred sixty thousand, one hundred and six.

Four

Four hundred and eight millions, two hundred and fifty-five thousand, one hundred and ninety-two.

Twenty-seven thousand and eight millions, ninety-six thousand two hundred and four.

Two hundred thousand and five hundred and fifty millions, one hundred and ten thousand, and sixteen.

Twenty-one billions, eight hundred and ten millions, sixty-four thousand, one hundred and fifty.

OF THE ROMAN NOTATION.

The Romans, like several other nations, expressed their numbers by certain letters of the alphabet. The Romans used only seven numeral letters, being the seven following capitals: viz. I for *one*; V for *five*; X for *ten*; L for *fifty*; C for an *hundred*; D for *five hundred*; M for a *thousand*. The other numbers they expressed by various repetitions and combinations of these, after the following manner:

1 = I	
2 = II	
3 = III	As often as any character is repeated, so many times is its value repeated.
4 = IIII or IV.	A less character before a greater diminishes its value.
5 = V	
6 = VI	A less character after a greater increases its value.
7 = VII	
8 = VIII	
9 = IX	
10 = X	
50 = L	
100 = C	
500 = D or IO	For every O annexed, this becomes 10 times as many.
1000 = M or CIO	For every C and O, placed one at each end, it becomes 10 times as much.
2000 = MM	
5000 = \overline{V} or IOO	A bar over any number increases it 1000 fold.
6000 = \overline{VI}	
10000 = \overline{X} or CCIOO	
50000 = \overline{L} or IOOO	
60000 = \overline{LX}	
100000 = \overline{C} or CCCIOOO	
1000000 = \overline{M} or CCCCIOOOO	
2000000 = \overline{MM}	
&c.	&c.

EXPLANATION OF CERTAIN CHARACTERS.

There are various characters or marks used in Arithmetic, and Algebra, to denote several of the operations and propositions; the chief of which are as follows :

- $+$ signifies *plus*, or addition.
 $-$ - - *minus*, or subtraction.
 \times or - multiplication.
 \div - - division.
 $:$: : : - proportion.
 $=$ - - equality.
 $\sqrt{\quad}$ - - square root.
 $\sqrt[3]{\quad}$ - - cube root, &c.
 ∞ - - diff. between two numbers when it is not known which is the greater.

Thus,

$5 + 3$, denotes that 3 is to be added to 5.

$6 - 2$, denotes that 2 is to be taken from 6.

7×3 , or $7 \cdot 3$, denotes that 7 is to be multiplied by 3.

$8 \div 4$, denotes that 8 is to be divided by 4.

$2 : 3 :: 4 : 6$, shows that 2 is to 3 as 4 is to 6.

$6 + 4 = 10$, shows that the sum of 6 and 4 is equal to 10.

$\sqrt{3}$, or $3\frac{1}{2}$, denotes the square root of the number 3.

$\sqrt[3]{5}$, or $5\frac{1}{3}$, denotes the cube root of the number 5.

7^2 , denotes that the number 7 is to be squared.

8^3 , denotes that the number 8 is to be cubed.

&c.

OF ADDITION.

ADDITION is the collecting or putting of several numbers together, in order to find their *sum*, or the total amount of the whole. This is done as follows :

Set or place the numbers under each other, so that each figure may stand exactly under the figures of the same value,
 that

that is, units under units, tens under tens, hundreds under hundreds, &c. and draw a line under the lowest number, to separate the given numbers from their sum, when it is found.—Then add up the figures in the column or row of units, and find how many tens are contained in that sum.—Set down exactly below what remains more than those tens, or if nothing remains, a cipher, and carry as many ones to the next row as there are tens.—Next add up the second row, together with the number carried, in the same manner as the first. And thus proceed till the whole is finished, setting down the total amount of the last row.

TO PROVE ADDITION.

First Method.—Begin at the top, and add together all the rows of numbers downwards; in the same manner as they were before added upwards; then if the two sums agree, it may be presumed the work is right. This method of proof is only doing the same work twice over, a little varied.

Second Method.—Draw a line below the uppermost number, and suppose it cut off. Then add all the rest of the numbers together in the usual way, and set their sum under the number to be proved.—Lastly, add this last found number and the uppermost line together; then if their sum be the same as that found by the first addition, it may be presumed the work is right.—This method of proof is founded on the plain axiom, that “The whole is equal to all its parts taken together.”

Third Method.—Add the figures in the uppermost line together, and find how many nines are contained in their sum.—Reject those nines, and set down the remainder towards the right-hand directly even with the figures in the line, as in the annexed example.—Do the same with each of the proposed lines of numbers, setting all these excesses of nines in a column on the right-hand, as here 5, 5, 6.

EXAMPLE I.

3497	Excess of nines.	5
6512		5
8295		6
———		—
18304		7
———		—

Then, if the excess of 9's in this sum, found as before, be equal to the excess of 9's in the total sum 18304, the work is probably right.—Thus, the sum of the right-hand column, 5, 5, 6, is 16, the excess of which above 9 is 7. Also the sum of the figures in

the sum total 18304, is 16, the excess of which above 9 is also 7, the same as the former*.

OTHER EXAMPLES.

2.	3.	4.
12345	12345	12345
67890	67890	876
98765	9876	9087
43210	543	56
12345	21	234
67890	9	1012
302445	90684	23610
290100	78339	11265
302445	90684	23610

* This method of proof depends on a property of the number 9, which except the number 3, belongs to no other digit whatever; namely, that "any number divided by 9, will leave the same remainder as the sum of its figures or digits divided by 9;" which may be demonstrated in this manner.

Demonstration. Let there be any number proposed as 4658. This, separated into its several parts, becomes $4000 + 600 + 50 + 8$. But $4000 = 4 \times 1000 = 4 \times (999 + 1) = 4 \times 999 + 4$. In like manner $600 = 6 \times 99 + 6$; and $50 = 5 \times 9 + 5$. Therefore the given number $4658 = 4 \times 999 + 4 + 6 \times 99 + 6 + 5 \times 9 + 5 + 8 = 4 \times 999 + 6 \times 99 + 5 \times 9 + 4 + 6 + 5 + 8$; and $4658 \div 9 = (4 \times 999 + 6 \times 99 + 5 \times 9 + 4 + 6 + 5 + 8) \div 9$. But $4 \times 999 + 6 \times 99 + 5 \times 9$ is evidently divisible by 9, without a remainder; therefore if the given number 4658 be divided by nine, it will leave the same remainder as $4 + 6 + 5 + 8$ divided by 9. And the same, it is evident, will hold for any other number whatever.

In like manner, the same property may be shown to belong to the number 3; but the preference is usually given to the number 9, on account of its being more convenient in practice.

Now from the demonstration above given, the reason of the rule itself is evident; for the excess of 9's in two or more numbers being taken separately, and the excess of 9's taken also out of the sum of the former excesses, it is plain that this last excess must be equal to the excess of 9's contained in the total sum of all these numbers; all the parts taken together being equal to the whole.—This rule was first given by Doctor Wallis in his Arithmetic, published in the year 1657.

Ex.

- Ex. 5. Add 3426 ; 9024 ; 5106 ; 8890 ; 1204, together.
 Ans. 27650.
6. Add 509267 ; 235809 ; 72920 ; 8392 ; 420 ; 21 ; and
 9, together. Ans. 826838.
7. Add 2 ; 19 ; 817 ; 4298 ; 50916 ; 730205 ; 9180634,
 together. Ans. 9966891.
8. How many days are in the twelve calendar months ?
 Ans. 365.
9. How many days are there from the 15th day of April to
 the 24th day of November, both days included ? Ans. 224.
10. An army consisting of 52714 infantry*, or foot, 5110
 horse, 6250 dragoons, 3927 light horse, 928 artillery, or
 gunners, 1410 pioneers, 250 sappers, and 406 miners : what
 is the whole number of men ; Ans. 70995.

OF SUBTRACTION.

SUBTRACTION teaches to find how much one number exceeds another called their *difference*, or the *remainder*, by taking the less from the greater. The method of doing which is as follows :

Place the less number under the greater, in the same manner as in addition, that is, units under units, tens under tens, and so on ; and draw a line below them.—Begin at the right-hand, and take each figure in the lower line, or number, from the figure above it, setting down the remainder below it.—But if the figure in the lower line be greater than that above it, first borrow, or add, 10 to the upper one, and then take the lower figure from that sum, setting down the remainder, and carrying 1, for what was borrowed, to the next lower figure, with which proceed as before ; and so on till the whole is finished.

* The whole body of foot soldiers is denoted by the word *Infantry* ; and all those that charge on horseback by the word *Cavalry*.—Some authors conjecture that the term infantry is derived from a certain Infanta of Spain, who finding that the army commanded by the king her father had been defeated by the Moors, assembled a body of the people together on foot, with which she engaged and totally routed the enemy. In honour of this event, and to distinguish the foot soldiers, who were not before held in much estimation, they received the name of Infantry, from her own title of Infanta.

TO PROVE SUBTRACTION.

ADD the remainder to the less number, or that which is just above it; and if the sum be equal to the greater or uppermost number, the work is right*.

EXAMPLES.

1.
From 5386427
Take 2164315

Rem. 3222112

Proof. 5386427

2.
From 5386427
Take 4258792

Rem. 1127635

Proof. 5386427

3.
From 1234567
Take 702973

Rem. 531594

Proof. 1234567

4. From 5331806 take 5073918.

Ans. 257888.

5. From 7020974 take 2766809.

Ans. 4254165.

6. From 8503602 take 574271.

Ans. 7929131.

7. Sir Isaac Newton was born in the year 1642, and he died in 1727: how old was he at the time of his decease?

Ans. 85 years.

8. Homer was born 2543 years ago, and Christ 1810 years ago: then how long before Christ was the birth of Homer?

Ans. 733 years.

9. Noah's flood happened about the year of the world 1656, and the birth of Christ about the year 4000: then how long was the flood before Christ?

Ans. 2344 years.

10. The Arabian or Indian method of notation was first known in England about the year 1150; then how long is it since to this present year 1810?

Ans. 660 years.

11. Gunpowder was invented in the year 1330: then how long was this before the invention of printing, which was in 1441?

Ans. 111 years.

12. The mariner's compass was invented in Europe in the year 1302: then how long was that before the discovery of America by Columbus, which happened in 1492?

Ans. 190 years.

* The reason of this method of proof is evident; for if the difference of two numbers be added to the less, it must manifestly make up a sum equal to the greater.

OF MULTIPLICATION.

MULTIPLICATION is a compendious method of Addition, teaching how to find the amount of any given number when repeated a certain number of times; as, 4 times 6, which is 24.

The number to be multiplied, or repeated, is called the *Multiplicand*.—The number you multiply by, or the number of repetitions, is the *Multiplier*.—And the number found, being the total amount, is called the *Product*.—Also, both the multiplier and multiplicand are, in general named the *Terms* or *Factors*.

Before proceeding to any operations in this rule, it is necessary to learn off very perfectly the following Table, of all the products of the first 12 numbers, commonly called the Multiplication Table, or sometimes Pythagoras's Table, from its inventor.

MULTIPLICATION TABLE.

1	2	3	4	5	6	7	8	9	10	11	12
2	4	6	8	10	12	14	16	18	20	22	24
3	6	9	12	15	18	21	24	27	30	33	36
4	8	12	16	20	24	28	32	36	40	44	48
5	10	15	20	25	30	35	40	45	50	55	60
6	12	18	24	30	36	42	48	54	60	66	72
7	14	21	28	35	42	49	56	63	70	77	84
8	16	24	32	40	48	56	64	72	80	88	96
9	18	27	36	45	54	63	72	81	90	99	108
10	20	30	40	50	60	70	80	90	100	110	120
11	22	33	44	55	66	77	88	99	110	121	132
12	24	36	48	60	72	84	96	108	120	132	144

To multiply any Given Number by a Single Figure, or by any Number not more than 12.

* Set the multiplier under the units figure, or right-hand place, of the multiplicand, and draw a line below it.—Then beginning at the right-hand, multiply every figure in this by the multiplier.—Count how many tens there are in the product of every single figure, and set down the remainder directly under the figure that is multiplied; and if nothing remains, set down a cipher.—Carry as many units or ones as there are tens counted, to the product of the next figures; and proceed in the same manner till the whole is finished.

EXAMPLE.

Multiply 9876543210 the Multiplicand.

By - - - - 2 the Multiplier.

19753086420 the Product.

To multiply by a Number consisting of Several Figures.

† Set the multiplier below the multiplicand, placing them as in Addition, namely, units under units, tens under tens, &c. drawing a line below it. — Multiply the whole of the multiplicand by each figure of the multiplier, as in the last article;

* The reason of this rule is the same as for the process in Addition, in which 1 is carried for every 10, to the next place, gradually as the several products are produced one after another, instead of setting them all down one below each other, as in the annexed example.

$$\begin{array}{r}
 5678 \\
 \underline{\quad} 4 \\
 \hline
 32 = 8 \times 4 \\
 230 = 70 \times 4 \\
 2400 = 600 \times 4 \\
 20000 = 5000 \times 4 \\
 \hline
 22712 = 5678 \times 4
 \end{array}$$

† After having found the produce of the multiplicand by the first figure of the multiplier, as in the former case, the multiplier is supposed to be divided into parts, and the product is found for the second figure in the same manner: but as this figure stands in the place of tens, the product must be ten times its simple value; and therefore the first figure of this product must be set in the place of setting

setting down a line of products for each figure in the multiplier, so as that the first figure of each line may stand straight under the figure multiplying by.—Add all the lines of products together in the order as they stand, and their sum will be the answer or whole product required.

TO PROVE MULTIPLICATION.

THERE are three different ways of proving Multiplication, which are as below.

First Method.—Make the multiplicand and multiplier change places, and multiply the latter by the former in the same manner as before. Then if the product found in this way be the same as the former, the number is right.

Second Method.—*Cast all the 9's out of the sum of the figures in each of the two factors, as in Addition, and set down the remainders. Multiply these two remainders together, and cast the 9's out of the product, as also out of

tens; or, which is the same thing, directly under the figure multiplied by. And proceeding in this manner separately with all the figures of the multiplier, it is evident that we shall multiply all the parts of the multiplicand by all the parts of the multiplier, or the whole of the multiplicand, by the whole of the multiplier: therefore, these several products being added together, will be equal to the whole required product; as in the example annexed.

1234567	the multiplicand.	
. 4567		
8641969	=	7 times the mult.
7407402	=	60 times ditto. *
6172835	=	500 times ditto.
4938268	=	4000 times ditto.
563827489	=	4567 times ditto.

* This method of proof is derived from the peculiar property of the number 9, mentioned in the proof of Addition, and the reason for the one may serve for that of the other. Another more ample demonstration of this rule may be as follows:—Let P and Q denote the number of 9's in the factors to be multiplied, and a and b what remain; then $9P + a$ and $9Q + b$ will be the numbers themselves, and their product is $(9P \times 9Q) + (9P \times b) + (9Q \times a) + (a \times b)$; but the first three of these products are each a precise number of 9's because their factors are so, either one or both: these therefore being cast away, there remains only $a \times b$; and if the 9's also be cast out of this, the excess is the excess of 9's in the total product: but a and b are the excesses in the factors themselves, and $a \times b$ is their product; therefore the rule is true.

the whole product or answer of the question, reserving the remainders of these last two, which remainders must be equal when the work is right.*—*Note*, It is common to set the four remainders within the four angular spaces of a cross, as in the example below.

Third Method.—Multiplication is also very naturally proved by Division; for the product divided by either of the factors, will evidently give the other. But this cannot be practised till the rule of Division is learned.

EXAMPLES.

Mult. 3542
by 6196

21252
31878
3542
21252

21946232 Product.

Proof.

~~2
5 4
2~~

or Mult. 6196
by 3542

12392
24784
30980
18588

21946232 Proof.

OTHER EXAMPLES.

Multiply 123456789 by 3.	Ans. 370370367.
Multiply 123456789 by 4.	Ans. 493827156.
Multiply 123456789 by 5.	Ans. 617283945.
Multiply 123456789 by 6.	Ans. 740740734.
Multiply 123456789 by 7.	Ans. 864197523.
Multiply 123456789 by 8.	Ans. 987654312.
Multiply 123456789 by 9.	Ans. 111111101.
Multiply 123456789 by 11.	Ans. 1358024679.
Multiply 123456789 by 12.	Ans. 1481481468.
Multiply 302914603 by 16.	Ans. 4846633648.
Multiply 273580961 by 23.	Ans. 6292362103.
Multiply 402097316 by 195.	Ans. 78408976620.
Multiply 82164973 by 3027.	Ans. 248713373271.
Multiply 7564900 by 579.	Ans. 4380077100.
Multiply 8496427 by 874359.	Ans. 7428927415293.
Multiply 2760325 by 37072.	Ans. 102330768400.

* If the two remainders be equal, it by no means follows that the answer is correct. Thus, if we multiply 13 by 12, the product is 156, and the remainders are each 3: but if we take for the answer or product any of the numbers 165, 561, 516, 246, &c. the remainders are the same as before, and therefore the rule is defective. *Ed.*

CONTRACTIONS IN MULTIPLICATION.

I. *When there are Ciphers in the Factors.*

If the ciphers be at the right-hand of the numbers ; multiply the other figures only, and annex as many ciphers to the right-hand of the whole product, as are in both the factors.—When the ciphers are in the middle parts of the multiplier ; neglect them as before, only taking care to place the first figure of every line of products exactly under the figure multiplying with.

EXAMPLES.

$$\begin{array}{r}
 \text{1.} \\
 \text{Mult. } 9001635 \\
 \text{by } - \quad 70100 \\
 \hline
 9001635 \\
 63011445 \\
 \hline
 631014613500
 \end{array}$$

$$\begin{array}{r}
 \text{2} \\
 \text{Mult. } 390720400 \\
 \text{by } - \quad 406000 \\
 \hline
 23443224 \\
 15628816 \\
 \hline
 158632482400000
 \end{array}$$

Products

3. Multiply 81503600 by 7030. Ans. 572970308000.
 4. Multiply 9030100 by 2100. Ans. 18963210000.
 5. Multiply 8057069 by 70050. Ans. 564397683450.

II. *When the multiplier is the Product of two or more Numbers in the Table ; then,*

* Multiply by each of those parts separately, instead of the whole number at once.

EXAMPLES.

1. Multiply 51307298 by 56, or 7 times 8.

$$\begin{array}{r}
 51307298 \\
 \quad \quad 7 \\
 \hline
 359151086 \\
 \quad \quad 8 \\
 \hline
 2873208688
 \end{array}$$

* The reason of this rule is obvious enough ; for any number multiplied by the component parts of another, must give the same product as if it were multiplied by that number at once. Thus, in the 1st example, 7 times the product of 8 by the given number, makes 56 times the same number, as plainly as 7 times 8 makes 56.

2. Multiply 31704592 by 36. Ans. 1141365312.
 3. Multiply 29753804 by 72. Ans. 2142273888.
 4. Multiply 7128368 by 96. Ans. 684323328.
 5. Multiply 160430800 by 108. Ans. 17326526400.
 6. Multiply 61835720 by 1320. Ans. 81623150400.
 7. There was an army composed of 104 * battalions, each consisting of 500 men ; what was the number of men contained in the whole ? Ans. 52000.
 8. A convoy of ammunition † bread, consisting of 250 waggons, and each waggon containing 320 loaves, having been intercepted and taken by the enemy, what is the number of loaves lost ? Ans. 80000.

OF DIVISION.

DIVISION is a kind of compendious method of Subtraction, teaching to find how often one number is contained in another, or may be taken out of it : which is the same thing.

The number to be divided is called the *Dividend*.—The number to divide by, is the *Divisor*.—And the number of times the dividend contains the divisor, is called the *Quotient*.—Sometimes there is a *Remainder* left, after the division is finished.

The usual manner of placing the terms, is, the dividend in the middle, having the divisor on the left hand, and the quotient on the right, each separated by a curve line ; as, to divide 12 by 4, the quotient is 3,

Divisor 4)	12	(3 Quotient ;	12
showing that the number 4 is 3 times	contained in 12, or may be 3 times	subtracted out of it, as in the margin.	4 subtr.
† <i>Rule</i> —Having placed the divisor	before the dividend, as above directed,	find how often the divisor is contained in as many figures of the dividend as are just necessary, and place the number on the right in the quotient.	—
			8
			4 subtr.
			—
			4
			4 subtr.
			—
			0
			—
			Mul-

* A battalion is a body of foot, consisting of 500, or 600, or 700 men, more or less.
 † The ammunition bread, is that which is provided for, and distributed to, the soldiers ; the usual allowance being a loaf of 6 pounds to every soldier, once in 4 days.

‡ In this way the dividend is resolved into parts, and by trial is found how often the

Multiply the divisor by this number, and set the product under the figures of the dividend before-mentioned.—Subtract this product from that part of the dividend under which it stands, and bring down the next figure of the dividend, or more if necessary, to join on the right of the remainder—Divide this number, so increased, in the same manner as before; and so on till all the figures are brought down and used.

N. B. If it be necessary to bring down more figures than one to any remainder, in order to make it as large as the divisor, or larger, a cipher must be set in the quotient for every figure so brought down more than one.

TO PROVE DIVISION.

* **MULTIPLY** the quotient by the divisor; to this product add the remainder, if there be any; then the sum will be equal to the dividend when the work is right.

the divisor is contained in each of those parts, one after another, arranging the several figures of the quotient one after another, into one number.

When there is no remainder to a division, the quotient is the whole and perfect answer to the question. But when there is a remainder, it goes so much towards another time, as it approaches to the divisor; so, if the remainder be half the divisor, it will go the half of a time more; if the 4th part of the divisor, it will go one fourth of a time more; and so on. Therefore, to complete the quotient, set the remainder at the end of it, above a small line, and the divisor below it, thus forming a fractional part of the whole quotient.

* This method of proof is plain enough; for since the quotient is the number of times the dividend contains the divisor, the quotient multiplied by the divisor must evidently be equal to the dividend.

There are also several other methods sometimes used for proving Division, some of the most useful of which are as follow:

Second Method—Subtract the remainder from the dividend; and divide what is left by the quotient; so shall the new quotient from this last division be equal to the former divisor, when the work is right.

Third Method—Add together the remainder and all the products of the several quotient figures by the divisor, according to the order in which they stand in the work; and the sum will be equal to the dividend when the work is right.

EXAMPLES.

1.	Quot.	2.	Quot.
$\begin{array}{r} 3) 1234567 \\ \underline{12} \\ 3 \\ \underline{3} \\ 4 \\ \underline{3} \\ \text{Proof.} \\ 15 \\ \underline{15} \\ 6 \\ \underline{6} \\ 7 \\ \underline{6} \\ \text{Rem. 1} \end{array}$	$\begin{array}{r} (411522 \\ \text{mult. 3} \\ \underline{1234566} \\ \text{add 1} \\ \underline{1234567} \\ \text{Proof.} \\ 15 \\ \underline{15} \\ 6 \\ \underline{6} \\ 7 \\ \underline{6} \\ \text{Rem. 1} \end{array}$	$\begin{array}{r} 37) 12345678 \\ \underline{111} \\ 124 \\ \underline{111} \\ 135 \\ \underline{111} \\ 246 \\ \underline{222} \\ 247 \\ \underline{222} \\ 258 \\ \underline{222} \\ \text{Rem. 36} \end{array}$	$\begin{array}{r} (333666. \\ \underline{37} \\ 2335662 \\ \underline{1000998} \\ \text{rem. 36} \\ \underline{12345678} \\ \text{Proof.} \\ 246 \\ \underline{222} \\ 247 \\ \underline{222} \\ 258 \\ \underline{222} \\ \text{Rem. 36} \end{array}$

3. Divide 73146085 by 4. Ans. $18286521\frac{1}{4}$.
4. Divide 5317986027 by 7. Ans. $7597122894\frac{1}{7}$.
5. Divide 570196382 by 12. Ans. $47516365\frac{2}{12}$.
6. Divide 74638105 by 37. Ans. $2017246\frac{3}{37}$.
7. Divide 137896254 by 97. Ans. $1421610\frac{24}{97}$.
8. Divide 35821649 by 764. Ans. $468867\frac{45}{764}$.
9. Divide 72091365 by 5201. Ans. $13861\frac{304}{5201}$.
10. Divide 4637064283 by 57606. Ans. $80496\frac{1707}{57606}$.
11. Suppose 471 men are formed into ranks of three deep, what is the number in each rank? Ans. 157.
12. A party at the distance of 378 miles from the head quarters, receive orders to join the corps in 18 days: what number of miles must they march each day to obey their orders? Ans. 21.
13. The annual revenue of a gentleman being 38330*l*; how much per day is that equivalent to, there being 365 days in the year? Ans. 104*l*.

CONTRACTIONS IN DIVISION.

There are certain contractions in Division, by which the operation in particular cases may be performed in a shorter manner, as follows:

1. *Divi-*

I. *Division by any Small Number*, not greater than 12, may be expeditiously performed, by multiplying and subtracting mentally, omitting to set down the work, except only the quotient immediately below the dividend.

EXAMPLES.

3) 56103961 <hr style="border: 0.5px solid black;"/> Quot. 18701320 $\frac{1}{3}$ <hr style="border: 0.5px solid black;"/> 6) 38672940 <hr style="border: 0.5px solid black;"/> 9) 43981962 <hr style="border: 0.5px solid black;"/>	4) 52619675 <hr style="border: 0.5px solid black;"/> 7) 81396627 <hr style="border: 0.5px solid black;"/> 11) 57614230 <hr style="border: 0.5px solid black;"/>	5) 1379192 <hr style="border: 0.5px solid black;"/> 8) 23718920 <hr style="border: 0.5px solid black;"/> 12) 27980373 <hr style="border: 0.5px solid black;"/>
--	--	---

II. * *When Ciphers are annexed to the Divisor*; cut off those ciphers from it, and cut off the same number of figures from the right-hand of the dividend; then divide with the remaining figures, as usual. And if there be any thing remaining after this division, place the figures cut off from the dividend to the right of it, and the whole will be the true remainder; otherwise, the figures cut off only will be the remainder.

EXAMPLES.

1. Divide 3704196 by 20. 2,0) 370419,6 <hr style="border: 0.5px solid black;"/> Quot. 185209 $\frac{1}{2}$ $\frac{6}{0}$	2. Divide 31086901 by 7100. 71,00) 310869,01 (4378 $\frac{3}{7}$ $\frac{1}{1}$ $\frac{0}{0}$ $\frac{1}{0}$. 284 <hr style="border: 0.5px solid black;"/> 268 213 <hr style="border: 0.5px solid black;"/> 556 497 <hr style="border: 0.5px solid black;"/> 599 568 <hr style="border: 0.5px solid black;"/> 31 <hr style="border: 0.5px solid black;"/>
--	--

3. Divide

* This method is only to avoid a needless repetition of ciphers which would happen in the common way. And the truth of the principle on which it is founded

3. Divide 7380964 by 23000.

Ans. $320\frac{2}{3}\frac{2}{3}\frac{2}{3}\frac{2}{3}$.

4. Divide 2304109 by 5800.

Ans. $397\frac{1}{3}\frac{2}{3}\frac{2}{3}$.

III. When the Divisor is the exact Product of two or more of the small Numbers not greater than 12 : * Divide by each of those numbers separately, instead of the whole divisor at once.

N. B. There are commonly several remainders in working by this rule, one to each division ; and to find the true or whole remainder, the same as if the division had been performed all at once, proceed as follows : Multiply the last remainder by the preceding divisor, or last but one, and to the product add the preceding remainder ; multiply this sum by the next preceding divisor, and to the product add the next preceding remainder ; and so on, till you have gone backward through all the divisors and remainders to the first. As in the example following :

EXAMPLES.

1. Divide 31046835 by 56, or 7 times 8.

$\begin{array}{r} 7) \ 31046835 \\ \hline 8) \ 4435262 \text{---} 1 \text{ first rem.} \\ \hline \quad 554407 \text{---} 6 \text{ second rem. add} \\ \hline \text{Ans. } 554407\frac{1}{7}\frac{2}{8} \end{array}$	$\begin{array}{l} 6 \text{ the last rem.} \\ \text{mult. } 7 \text{ preced. divisor.} \\ \hline 42 \\ \hline 1 \text{ the 1st rem.} \\ \hline 43 \text{ whole rem.} \end{array}$
---	--

2. Divide 7014596 by 72.

Ans. $97424\frac{4}{7}\frac{8}{2}$.

3. Divide 5130652 by 132.

Ans. $38868\frac{1}{3}\frac{6}{2}$.

4. Divide 83016572 by 240.

Ans. $345902\frac{2}{2}\frac{2}{4}$.

founded is evident : for cutting off the same number of ciphers, or figures, from each, is the same as dividing each of them by 10, or 100, or 1000, &c. according to the number of ciphers cut off ; and it is evident, that as often as the whole divisor is contained in the whole dividend, so often must any part of the former be contained in a like part of the latter.

* This follows from the second contraction in Multiplication, being only the converse of it ; for the half of the third part of any thing, is evidently the same as the sixth part of the whole ; and so of any other numbers. — The reason of the method of finding the whole remainder from the several particular ones, will best appear from the nature of Vulgar fractions. Thus in the first example above, the first remainder being 1, when the divisor is 7, makes $\frac{1}{7}$ this must be added to the second remainder, 6, making $6\frac{1}{7}$ to the divisor 8, or to be divided by 8.

But $6\frac{1}{7} = \frac{6 \times 7 + 1}{7} = \frac{43}{7}$; and this divided by 8, gives $\frac{43}{7 \times 8} = \frac{43}{56}$.

IV. Common

IV. *Common Division may be performed more concisely, by omitting the several products, and setting down only the remainders; namely, multiply the divisor by the quotient figures as before, and, without setting down the product, subtract each figure of it from the dividend, as it is produced; always remembering to carry as many to the next figure as were borrowed before.*

EXAMPLES.

1. Divide 3104679 by 833.

833)	3104679	(3727	⁹ ₈	⁹ ₃
	6056			
	2257			
	5919			
	88			
2. Divide 79165238 by 238. Ans. 332627 ¹²₂₃₈.
3. Divide 29137062 by 5317. Ans. 5479 ⁵²¹⁹₅₃₁₇.
4. Divide 62015735 by 7803. Ans. 7947 ²⁸⁰⁴₇₈₀₃.

OF REDUCTION.

REDUCTION is the changing of numbers from one name or denomination to another, without altering their value.—This is chiefly concerned in reducing money, weights, and measures.

When the numbers are to be reduced from a higher name to a lower, it is called *Reduction Descending*; but when, contrarywise, from a lower name to a higher, it is *Reduction Ascending*.

Before proceeding to the rules and questions of Reduction, it will be proper to set down the usual Tables of money, weights, and measures, which are as follow :

Of MONEY, WEIGHTS, AND MEASURES.

TABLES OF MONEY.*

2 ¹ Farthings = 1 Halfpenny	$\frac{1}{2}$	<i>grs</i>	<i>d</i>		
4 Farthings = 1 Penny	<i>d</i>	4 =	1	<i>s</i>	
12 Pence = 1 Shilling	<i>s</i>	48 =	12 =	1	£
20 Shillings = 1 Pound	£	960 =	240 =	20 =	1
					PENCE

* £ denotes pounds, *s* shillings, and *d* denotes pence.
 $\frac{1}{4}$ denotes 1 farthing, or one quarter of any thing.
 $\frac{1}{2}$ denotes a halfpenny, or the half of any thing.
 $\frac{3}{4}$ denotes 3 farthings, or three quarters of any thing.

PENCE TABLE.

<i>d</i>		<i>s</i>	<i>d</i>
20	is	1	8
30	—	2	6
40	—	3	4
50	—	4	2
60	—	5	0
70	—	5	10
80	—	6	8
90	—	7	6
100	—	8	4
110	—	9	2
120	—	10	0

SHILLINGS TABLE.

<i>s</i>		<i>d</i>
1	is	12
2	—	24
3	—	36
4	—	48
5	—	60
6	—	72
7	—	84
8	—	96
9	—	108
10	—	120
11	—	132

FEDERAL MONEY.

10 Mills (m)	= 1 Cent		Standard Weight.	dwt	gr
10 Cents	= 1 Dime <i>d</i>		The Cent weighs	6	23 Copper
10 Dimes	= 1 Dollar <i>D</i>		Dollar	17	1 $\frac{3}{4}$ Silver
10 Dollars	= 1 Eagle <i>E</i>		Eagle	11	4 $\frac{2}{3}$ Gold

The standard for Federal Money of Gold and Silver is 11 parts fine, and 1 part alloy.

A Dollar is equal to 4*s* and 8*d* in South Carolina, to 6*s* in the New-England States and Virginia, to 7*s* and 6*d* in New-Jersey, Pennsylvania, Delaware, and Maryland, and to 8*s* in New-York, and North-Carolina.

TROY

The full weight and value of the English gold and silver coin, is as here below :

GOLD.			SILVER.					
Value.		Weight.	Value.		Weight.			
<i>L</i>	<i>s</i>	<i>d</i>	<i>s</i>	<i>d</i>	<i>dwt</i>	<i>gr</i>		
A Guinea	1	1	0	A Crown	5	0	19	8 $\frac{1}{2}$
Half guinea	0	10	6	Half-Crown	2	6	9	16 $\frac{1}{4}$
Seven Shillings	0	7	0	Shilling	1	0	3	21
Quarter-guinea	0	5	3	Sixpence	0	6	1	22 $\frac{1}{2}$

The usual value of gold is nearly 4*l* an ounce, or 2*d* a grain; and that of silver is nearly 5*s* an ounce. Also the value of any quantity of gold, is to the value of the same weight of standard silver, nearly as 15 to 1, or more nearly as 15 and 1-14th to 1.

Pure gold, free from mixture with other metals, usually called fine gold is of so pure a nature, that it will endure the fire without wasting, though it be kept continually

TROY WEIGHT.*

Grains	-	-	marked	<i>gr</i>	<i>gr</i>	<i>dwt</i>
24 Grains	make	1 Pennyweight	<i>dwt</i>		24=	1 <i>oz</i>
20 Pennyweights	1 Ounce		<i>oz</i>		480=	20= 1 <i>lb</i>
12 Ounces	1 Pound		<i>lb</i>		5760=	240=12= 1

By this weight are weighed Gold, Silver, and Jewels.

APOTHECARIES' WEIGHT.

Grains	-	-	marked	<i>gr</i>
20 Grains	make	1 Scruple	<i>sc</i>	or ᶒ
3 Scruples	-	1 Dram	<i>dr</i>	or $\frac{3}{4}$
8 Drams	-	1 Ounce	<i>oz</i>	or $\frac{3}{4}$
12 Ounces	-	1 Pound	<i>lb</i>	or ᶒ

<i>gr</i>	<i>sc</i>			
20 =	1	<i>dr</i>		
60 =	3 =	1	<i>oz</i>	
480 =	24 =	8 =	1	<i>lb</i>
5760 =	288 =	96 =	12 =	1

This is the same as Troy weight, only having some different divisions. Apothecaries make use of this weight in compounding their Medicines; but they buy and sell their Drugs by Avoirdupois weight

AVOIR-

continually melted. But silver, not having the purity of gold, will not endure the fire like it; yet fine silver will waste but a very little by being in the fire any moderate time; whereas copper, tin, lead, &c. will not only waste, but may be calcined, or burnt to a powder.

Both gold and silver, in their purity, are so very soft and flexible (like new lead, &c), that they are not so useful, either in coin or otherwise (except to beat into leaf gold or silver), as when they are allayed, or mixed and hardened with copper or brass. And though most nations differ, more or less, in the quantity of such alloy, as well as in the same place at different times, yet in England the standard for gold and silver coin has been for a long time as follows—viz. That 22 parts of fine gold, and 2 parts of copper, being melted together, shall be esteemed the true standard for gold coin: And that 11 ounces and 2 pennyweights of fine silver, and 18 pennyweights of copper, being melted together, is esteemed the true standard for silver coin, called Sterling silver.

* The original of all weights used in England, was a grain or corn of wheat, gathered out of the middle of the ear, and, being well dried, 32 of them were to make one pennyweight, 20 pennyweights one ounce, and 12 ounces one pound.

But

ARITHMETIC.

AVOIRDUPOIS WEIGHT.

Drms	-	-	-	-	marked	<i>dr</i>
16 Drms	make	1 Ounce	-	-	-	<i>oz</i>
16 Ounces	-	1 Pound	-	-	-	<i>lb</i>
28 Pounds	-	1 Quarter	-	-	-	<i>qr</i>
4 Quarters	-	1 Hundred weight	-	-	-	<i>cwt</i>
20 Hundred Weight	1 Ton	-	-	-	-	<i>ton</i>

<i>dr</i>	<i>oz</i>					
16 =	1	<i>lb</i>				
256 =	16 =	1	<i>qr</i>			
7168 =	448 =	28 =	1	<i>cwt</i>		
28672 =	1792 =	112 =	4 =	1	<i>ton</i>	
573440 =	35840 =	2240 =	80 =	20 =	1	

By this weight are weighed all things of a coarse or drossy nature, as Corn, Bread, Butter, Cheese, Flesh, Grocery Wares, and some Liquids; also all Metals, except Silver and Gold.

Note, that 1*lb* Avoirdupois $\overset{\text{oz dwt gr}}{=} 14\ 11\ 15\frac{1}{2}$ Troy.
 1*oz* - - = 0 18 $5\frac{1}{2}$
 1*dr* - - = 0 1 $3\frac{1}{2}$

Hence it appears that the pound Avoirdupois contains 6999 $\frac{1}{2}$ grains, and the pound Troy 5760; the former of which augmented by half a grain becomes 7000, and its ratio to the latter is therefore very nearly as 700 to 576, that is, as 175 to 144; consequently 144 pounds Avoirdupois are very nearly equal to 175 pounds Troy: and hence we infer that the ounce Avoirdupois is to the ounce Troy as 175 to 192.

LONG MEASURE.

3 Barley-corns	make	1 Inch	-	-	<i>In.</i>
12 Inches	-	1 Foot	-	-	<i>Ft</i>
3 Feet	-	1 Yard	-	-	<i>Yd</i>
6 Feet	-	1 Fathom	-	-	<i>Fth</i>
5 Yards and a half	1 Pole or Rod	-	-	-	<i>Pl</i>
40 Poles	-	1 Furlong	-	-	<i>Fur</i>
8 Furlongs	-	1 Mile	-	-	<i>Mile</i>
3 Miles	-	1 League	-	-	<i>Lea</i>
69 $\frac{1}{5}$ Miles nearly	-	1 Degree	-	-	<i>Deg or °,</i>

But in latter times, it was thought sufficient to divide the same pennyweight into 24 equal parts, still called grains, being the least weight now in common use; and from thence the rest are computed, as in the Tables above.

<i>In</i>	<i>Ft</i>					
12 =	1		<i>Yd</i>			
36 =	3 =		1	<i>Pl</i>		
198 =	$16\frac{1}{2}$ =		$5\frac{1}{2}$ =	1	<i>Fur</i>	
7920 =	660 =		220 =	40 =	1	<i>Mile</i>
63360 =	5280		1760 =	320 =	8 =	1

CLOTH MEASURE.

2 Inches and a quarter make	1 Nail	-	-	-		<i>Nl</i>
4 Nails	-	-	-	1 Quarter of a Yard		<i>Qr</i>
3 Quarters	-	-	-	1 Ell Flemish	-	<i>E F</i>
4 Quarters	-	-	-	1 Yard	-	<i>Yd</i>
5 Quarters	-	-	-	1 Ell English	-	<i>E E</i>
4 Quarters $1\frac{1}{5}$ Inch	-	-	-	1 Ell Scotch	-	<i>E S</i>

SQUARE MEASURE.

144	Square Inches make	1	Sq Foot	-	<i>Ft</i>
9	Square Feet	-	1 Sq Yard	-	<i>Yd</i>
$30\frac{1}{4}$	Square Yards	-	1 Sq Pole	-	<i>Pole</i>
40	Square Poles	-	1 Rood	-	<i>Rd</i>
4	Roods	-	1 Acre	-	<i>Acr</i>

<i>Sq Inch</i>	<i>Sq Ft</i>		<i>Sq Yd</i>			
144 =	1		<i>Sq Yd</i>			
1296 =	9 =		1	<i>Sq Pl</i>		
39204 =	$272\frac{1}{4}$ =		$30\frac{1}{4}$ =	1	<i>Rd</i>	
1568160 =	10890 =		1210 =	40 =	1	<i>Acr</i>
6272640 =	43560 =		4840 =	160 =	4 =	1

By this measure, Land, and Husbandmen and Gardeners' work are measured; also Artificers' work, such as Board, Glass, Pavements, Plastering, Wainscoting Tiling, Flooring, and every dimension of length and breadth only.

When three dimensions are concerned, namely, length, breadth, and depth or thickness, it is called cubic or solid measure, which is used to measure Timber, Stone, &c.

The cubic or solid Foot, which is 12 inches in length and breadth and thickness, contains 1728 cubic or solid inches, and 27 solid feet make one solid yard.

DRY, OR CORN MEASURE.

2 Pints make	1 Quart	-	-	<i>Qt</i>
2 Quarts	- 1 Pottle	-	-	<i>Pot</i>
2 Pottles	- 1 Gallon	-	-	<i>Gal</i>
2 Gallons	- 1 Peck	-	-	<i>Pec</i>
4 Pecks	- 1 Bushel	-	-	<i>Bu</i>
8 Bushels	- 1 Quarter	-	-	<i>Qr</i>
5 Quarters	- 1 Wey, Load, or Ton	-	-	<i>Wey</i>
2 Weys	- 1 Last	-	-	<i>Last</i>

<i>Pts</i>	<i>Gal</i>		<i>Pec</i>				
8 =	1						
16 =	2 =	1		<i>Bu</i>			
64 =	8 =	4 =	1		<i>Qr</i>		
512 =	64 =	32 =	8 =	1		<i>Wey</i>	
2560 =	320 =	160 =	40 =	5 =	1		<i>Last</i>
5120 =	640 =	320 =	80 =	10 =	2 =	1	

By this are measured all dry wares, as, Corn, Seeds, Roots, Fruit, Salt, Coals, Sand, Oysters, &c.

The standard Gallon dry-measure contains $268\frac{4}{5}$ cubic or solid inches, and the Corn or Winchester bushel $2150\frac{2}{5}$ cubic inches; for the dimensions of the Winchester bushel, by the Statute, are 8 inches deep, and $18\frac{1}{2}$ inches wide or in diameter, but the Coal bushel must be $19\frac{1}{2}$ inches in diameter; and 36 bushels, heaped up, make a London chaldron of coals, the weight of which is 3156lb Avoirdupois.

ALE AND BEER MEASURE.

2 Pints make	-	1 Quart	-	<i>Qt</i>
4 Quarts	-	1 Gallon	-	<i>Gal</i>
36 Gallons	-	1 Barrel	-	<i>Bar</i>
1 Barrel and a half		1 Hogshead	-	<i>Hhd</i>
2 Barrels	-	1 Puncheon	-	<i>Pun</i>
2 Hogsheads	-	1 Butt	-	<i>Butt</i>
2 Butts	-	1 Tun	-	<i>Tun</i>

<i>Pts</i>	<i>Qt</i>		<i>Gal</i>			
2 =	1					
8 =	4 =	1		<i>Bar</i>		
288 =	144 =	36 =	1		<i>Hhd</i>	
432 =	216 =	54 =	$1\frac{1}{2}$ =	1		<i>Butt</i>
864 =	432 =	108 =	3 =	2 =	1	

Note, The Ale Gallon contains 282 cubic or solid inches.

WINE MEASURE.

2 Pints	make	-	-	1 Quart	-	Qt
4 Quarts	-	-	-	1 Gallon	-	Gal
42 Gallons	-	-	-	1 Tierce	-	Tier
63 Gallons	or	1½ Tierces		1 Hogshead	-	Hhd
2 Tierces	-	-	-	1 Puncheon	-	Pun
2 Hogsheads	-	-	-	1 Pipe or Butt	-	Pi
2 Pipes	or	4 Hhds	-	1 Tun	-	Tun

<i>Pts</i>		<i>Qt</i>				
2 =		1		<i>Gal</i>		
8 =		4 =		1		<i>Tier</i>
336 =	168 =	42 =		1		<i>Hhd</i>
504 =	252 =	63 =		1½ =		1 <i>Pun</i>
672 =	336 =	84 =		2 =		1½ = 1 <i>P</i>
1008 =	504 =	126 =		3 =		2 = 1½ = 1 <i>Tun</i>
2016 =	1008 =	252 =		6 =		4 = 3 = 2 = 1

Note, By this are measured all Wines, Spirits, Strong-waters, Cider, Mead, Perry, Vinegar, Oil, Honey, &c.

The Wine Gallon contains 231 cubic or solid inches. And it is remarkable, that the Wine and Ale Gallons have the same proportion to each other, as the Troy and Avoirdupois Pounds have; that is, as one Pound Troy is to one Pound Avoirdupois, so is one Wine Gallon to one Ale Gallon.

OF TIME.

60 Seconds	or	60"	make	-	1 Minute	-	Mo'
60 Minutes	-	-	-	-	1 Hour	-	Hr
24 Hours	-	-	-	-	1 Day	-	Day
7 Days	-	-	-	-	1 Week	-	Wk
4 Weeks	-	-	-	-	1 Month	-	Mo
13 Months	1 Day	6 Hours,	}		1 Julian Year		Yr
	or	365 Days		6 Hours			

<i>Sec</i>		<i>Min</i>				
60 =		1		<i>Hr</i>		
3600 =		60 =		1		<i>Day</i>
36400 =		1440 =		24 =		1 <i>Wk</i>
604800 =		10080 =		168 =		7 = 1 <i>Mo</i>
2419200 =		40320 =		672 =		28 = 4 = 1
31557600 =		525960 =		8766 =		365¼ = 1 Year

Or

Wk Da Hr Mo Da Hr
 Or 52 1 6 = 13 1 6 = 1 *Julian Year*
Da Hr M Sec
 But 365 5 48 48 = 1 *Solar Year.*

RULES FOR REDUCTION.

I. *When the Numbers are to be reduced from a Higher Denomination to a Lower :*

MULTIPLY the number in the highest denomination by as many as of the next lower make an integer, or 1, in that higher ; to this product add the number, if any, which was in this lower denomination before, and set down the amount.

Reduce this amount in like manner, by multiplying it by as many as of the next lower, make an integer of this, taking in the odd parts of this lower, as before. And so proceed through all the denominations to the lowest ; so shall the number last found be the value of all the numbers which were in the higher denominations, taken together.*

EXAMPLE.

1. In 1234*l* 15*s* 7*d*, how many farthings ?

<i>l</i>	<i>s</i>	<i>d</i>
1234	15	7
	20	

24695 Shillings

12

296347 Pence

4

Answer 1185388 Farthings.

* The reason of this rule is very evident ; for pounds are brought into shillings by multiplying them by 20 ; shillings into pence, by multiplying them by 12 ; and pence into farthings, by multiplying by 4 ; and the reverse of this rule by Division.—And the same, it is evident, will be true in the reduction of numbers consisting of any denominations whatever.

II. *When*

II. *When the Numbers are to be reduced from a Lower Denomination to a Higher :*

DIVIDE the given number by as many as of that denomination make 1 of the next higher, and set down what remains, as well as the quotient.

Divide the quotient by as many as of this denomination make 1 of the next higher ; setting down the new quotient, and remainder, as before.

Proceed in the same manner through all the denominations, to the highest ; and the quotient last found, together with the several remainders, if any, will be of the same value as the first number proposed.

EXAMPLES.

2. Reduce 1185388 farthings into pounds, shillings, and pence.

$$\begin{array}{r}
 4)1185388 \\
 \hline
 12) 296347 \text{ d} \\
 \hline
 2,0) 2469,5 \text{ s}—7\text{d} \\
 \hline
 \text{Answer } 1234\text{l } 15\text{s } 7\text{d} \\
 \hline
 \hline
 \end{array}$$

3. Reduce 24l to farthings. Ans. 23040.
4. Reduce 337587 farthings to pounds, &c. Ans. 351l 13s 0 $\frac{3}{4}$.
5. How many farthings are in 36 guineas ? Ans. 36288.
6. In 36288 farthings how many guineas ? Ans. 36.
7. In 59lb 13dwts 5gr how many grains ? Ans. 340157.
8. In 8012131 grains how many pounds, &c. ? Ans. 1390 lb 11 oz 18 dwt 19 gr.
9. In 35 ton 17 cwt 1 qr 23 lb 7 oz 13 dr how many drams ? Ans. 20571005.
10. How many barley-corns will reach round the earth, supposing it, according to the best calculations, to be 24877 miles ? Ans. 4728620160.
11. How many seconds are in a solar year, or 365 days 5 hrs 48 min 48 sec ? Ans. 31556928.
12. In a lunar month, or 29 ds 12 hrs 44 min 3 sec, how many seconds ? Ans. 2551443.

COMPOUND ADDITION.

COMPOUND ADDITION shows how to add or collect several numbers of different denominations into one sum.

RULE.—Place the numbers so, that those of the same denomination may stand directly under each other, and draw a line below them. Add up the figures in the lowest denomination, and find, by Reduction, how many units, or ones, of the next higher denomination are contained in their sum.—Set down the remainder below its proper column, and carry those units or ones to the next denomination, which add up in the same manner as before.—Proceed thus through all the denominations, to the highest, whose sum, together with the several remainders, will give the answer sought.

The method of proof is the same as in Simple Addition.

EXAMPLES OF MONEY.

1.			2.			3.			4.		
<i>l</i>	<i>s</i>	<i>d</i>	<i>l</i>	<i>s</i>	<i>d</i>	<i>l</i>	<i>s</i>	<i>d</i>	<i>l</i>	<i>s</i>	<i>d</i>
7	13	3	14	7	5	15	17	10	53	14	8
<hr/>			<hr/>			<hr/>			<hr/>		
3	5	$10\frac{1}{2}$	8	19	$2\frac{1}{4}$	3	14	6	5	10	$2\frac{3}{4}$
6	18	7	7	8	$1\frac{1}{2}$	23	6	$2\frac{3}{4}$	93	11	6
0	2	$5\frac{3}{4}$	21	2	9	14	9	$4\frac{1}{2}$	7	5	0
4	0	3	7	16	$8\frac{1}{2}$	15	6	4	13	2	5
17	15	$4\frac{1}{2}$	0	4	3	6	12	$9\frac{3}{4}$	0	18	7
<hr/>			<hr/>			<hr/>			<hr/>		
39	15	$9\frac{3}{4}$									
<hr/>			<hr/>			<hr/>			<hr/>		
32	2	$6\frac{3}{4}$									
<hr/>			<hr/>			<hr/>			<hr/>		
39	15	$9\frac{3}{4}$									
<hr/>			<hr/>			<hr/>			<hr/>		
<hr/>			<hr/>			<hr/>			<hr/>		
5.			6.			7.			8.		
<i>l</i>	<i>s</i>	<i>d</i>	<i>l</i>	<i>s</i>	<i>d</i>	<i>l</i>	<i>s</i>	<i>d</i>	<i>l</i>	<i>s</i>	<i>d</i>
14	0	$7\frac{1}{4}$	37	15	8	61	3	$2\frac{1}{2}$	472	15	3
8	15	3	14	12	$9\frac{3}{4}$	7	16	8	9	2	$2\frac{1}{2}$
62	4	7	17	14	9	29	13	$10\frac{3}{4}$	27	12	$6\frac{1}{4}$
4	17	8	23	10	$9\frac{1}{4}$	12	16	2	370	16	$2\frac{1}{2}$
23	0	$4\frac{3}{4}$	8	6	0	0	7	$5\frac{1}{4}$	13	7	4
6	6	7	14	0	$5\frac{1}{2}$	24	13	0	6	10	$5\frac{1}{4}$
91	0	$10\frac{1}{4}$	54	2	$7\frac{1}{2}$	5	0	$10\frac{3}{4}$	30	0	$11\frac{3}{4}$
<hr/>			<hr/>			<hr/>			<hr/>		
<hr/>			<hr/>			<hr/>			<hr/>		

EXAM-

EXAM. 9. A nobleman going out of town, is informed by his steward that his butcher's bill comes to $197l\ 13s\ 7\frac{1}{2}d$; his baker's to $59l\ 5s\ 2\frac{3}{4}d$; his brewer's to $85l$; his wine-merchant's to $103l\ 13s$; to his corn-chandler is due $7\cdot l\ 3d$; to his tallow-chandler and cheesemonger, $27l\ 15s\ 11\frac{1}{4}d$; and to his tailor $55l\ 3s\ 5\frac{3}{4}d$; also for rent, servants' wages, and other charges, $127l\ 3s$: Now, supposing he would take $100l$ with him to defray his charges on the road for what sum must he send to his banker? Ans. $830l.\ 14s.\ 6\frac{1}{4}d.$

10. The strength of a regiment of foot, of 10 companies, and the amount of their subsistence*, for a month of 30 days, according to the annexed Table, are required?

Numb.	Rank.	Subsistence for a Month.		
		<i>l</i>	<i>s</i>	<i>d</i>
1	Colonel	27	0	0
1	Lieutenant Colonel	19	10	0
1	Major	17	5	0
7	Captains	78	15	0
11	Lieutenants	57	15	0
9	Ensigns	40	10	0
1	Chaplain	7	10	0
1	Adjutant	4	10	0
1	Quarter-Master	5	5	0
1	Surgeon	4	10	0
1	Surgeon's Mate	4	10	0
30	Serjeants	45	0	0
30	Corporals	30	0	0
20	Drummers	20	0	0
2	Fifers	2	0	0
390	Private Men	292	10	0
507	Total	656	10	0

* Subsistence Money, is the money paid to the soldiers weekly, which is short of their full pay, because their clothes, accoutrements, &c. are to be accounted for. It is likewise the money advanced to officers till their accounts are made up, which is commonly once a year, when they are paid their arrears. The following Table shows the full pay and subsistence of each rank on the English establishment.

DAILY PAY OF COMMISSIONED OFFICERS.

RANK.	Horse Artillery & corps of C. Cm.	D. G. Dr. & F. Cav.	Foot Ar. & Fenc. Militia.	Life Guards.		Horse Guards.		Foot Guards.	
				Subsist.	Full Pay.	Subsist.	Full Pay.	Subsist.	Full Pay.
Colonel (Comm.)	—	1 12 10 2	4 0 1	2 6 1	7 0 1	16 0 1	11 0 0	1 10 0	1 19 0
Colonel (en Second)	1 10 0	—	1 4 0	—	—	—	—	—	—
1st Lieut. Col. . . .	1 6 0	1 3 0	1 0 0	15 11 1	3 3 1	11 0 1	2 6 6	1 1 6	1 8 6
2d Ditto.	—	—	0 17 0	—	—	—	—	—	—
1st Major	1 1 0	0 19 30	15 0 0	14 1 0	19 6 1	6 0 1	1 6 6	0 18 6	1 4 6
2d Ditto.	—	—	—	—	0 18 0	1 4 0	—	—	—
Captain	0 15 0	0 14 70	10 0 0	9 5 0	12 10 16	0 0 16	6 6 6	0 12 6	0 16 6
Capt. Lieut.	0 10 0	0 9 0	7 0 0	5 8 0	8 20 11	0 0 11	6 0 15	0 6 0	0 7 10
1st Lieut.	0 9 0	0 9 0	6 0 0	5 8 0	—	—	—	—	—
2d Ditto.	0 8 0	—	0 5 0	—	—	—	—	—	—
Cornet	—	0 8 0	—	—	—	—	0 11 0	0 14 0	—
Ensign	—	—	—	0 4 8	—	—	—	—	—
Adjutant	—	0 5 0	—	0 5 0	8 6 0	11 0 0	4 6 0	5 0 0	3 0 0
Pay-master	—	—	—	—	—	—	—	—	—
Quarter-master	—	0 5 0	—	0 5 8	—	—	0 6 6	0 8 6	—
Surgeon-major	0 12 0	0 11 40	10 0 0	9 5 5	—	—	—	—	0 12 6
Bat. Surg. or Surg. . .	—	—	—	—	0 6 0	0 8 0	0 9 0	0 12 0	0 7 6
Assist. Surg.	0 6 0	0 5 0	5 0 0	5 0 0	—	—	0 5 0	0 5 0	0 5 0
Veter Surg.	0 8 0	0 8 0	—	—	—	—	—	—	—
Solicitor	—	—	—	—	—	—	—	—	—

N. B. When a *Lieutenant, Ensign, Adjutant, or Quarter-master of Foot, Militia, Fencible Infantry, or Invalids*, holds two commissions, one shilling per day is to be deducted from the above rates for each commission.

EXAMPLES OF WEIGHTS, MEASURES, &c.

TROY WEIGHT.

1.			2.		
lb	oz	dwt	oz	dwt	gr
17	3	15	37	9	3
7	9	4	9	5	3
0	10	7	8	12	12
9	5	0	17	7	8
176	2	17	5	9	0
23	11	12	3	0	19

APOTHECARIES' WEIGHT.

3.				4.			
lb	oz	dr	sc	oz	dr	sc	gr
3	5	7	2	3	5	1	17
13	7	3	0	7	3	2	5
19	10	6	2	16	7	0	12
0	9	1	2	7	3	2	9
36	3	5	0	4	1	2	18
5	8	6	1	36	4	1	14

AVOIRDUPOIS WEIGHT.

5.			6.		
lb	oz	dr	cwt	qr	lb
17	10	13	15	2	15
5	14	8	6	3	24
12	9	18	9	1	14
27	1	6	9	1	17
0	4	0	10	2	6
6	14	10	3	0	3

LONG MEASURE.

7.			8.		
mls	fur	pls	yds	feet	inc
29	3	14	127	1	5
19	6	29	12	2	9
7	0	24	10	0	10
9	1	37	54	1	11
7	0	3	5	2	7
4	5	9	23	0	5

CLOTH MEASURE.

9.			10.		
yds	qr	nls	el	en	qrs nls
26	3	1	270	1	0
13	1	2	57	4	3
9	1	2	18	1	2
217	0	3	0	3	2
9	1	0	10	1	0
55	3	1	4	4	1

LAND MEASURE.

11.			12.		
ac	ro	p	ac	ro	p
225	3	37	19	0	16
16	1	25	270	3	29
7	2	18	6	3	13
4	2	9	23	0	34
42	1	19	7	2	16
7	0	6	75	0	23

WINE MEASURE.

13.			14.		
t	hds	gal	hds	gal	pts
13	3	15	15	61	5
8	1	37	17	14	13
14	1	20	29	23	7
25	0	12	3	15	1
3	1	9	16	8	0
72	3	21	4	36	6

ALE AND BEER MEASURE.

15.			16.		
hds	gal	pts	hds	gal	pts
17	37	3	29	43	5
9	10	15	12	19	7
3	6	2	14	16	6
5	14	0	6	8	1
12	9	6	57	13	4
8	42	4	5	6	0

COMPOUND SUBTRACTION.

COMPOUND SUBTRACTION shows how to find the difference between any two numbers of different denominations. To perform which, observe the following Rule :

* PLACE the less number below the greater, so that the parts of the same denomination may stand directly under each other; and draw a line below them.—Begin at the right-hand, and subtract each number or part in the lower line, from the one just above it, and set the remainder straight below it.—But if any number in the lower line be greater than that above it, add as many to the upper number as make 1 of the next higher denomination; then take the lower number from the upper one thus increased, and set down the remainder. Carry the unit borrowed to the next number in the lower line; after which subtract this number from the one above it, as before; and so proceed till the whole is finished. Then the several remainders, taken together, will be the whole difference sought.

The method of proof is the same as in Simple Subtraction.

EXAMPLES OF MONEY.

	1.			2.			3.			4.		
	<i>l</i>	<i>s</i>	<i>d</i>	<i>l</i>	<i>s</i>	<i>d</i>	<i>l</i>	<i>s</i>	<i>d</i>	<i>l</i>	<i>s</i>	<i>d</i>
From	79	17	8 $\frac{3}{4}$	103	3	2 $\frac{1}{2}$	81	10	11	254	12	0
Take	35	12	4 $\frac{1}{4}$	71	12	5 $\frac{3}{4}$	29	13	3 $\frac{1}{4}$	37	9	4 $\frac{3}{4}$
	<hr/>			<hr/>			<hr/>			<hr/>		
Rem.	44	5	4 $\frac{1}{2}$	31	10	8 $\frac{3}{4}$						
	<hr/>			<hr/>			<hr/>			<hr/>		
Proof.	79	17	8 $\frac{3}{4}$	103	3	2 $\frac{1}{2}$						
	<hr/>			<hr/>			<hr/>			<hr/>		

5. What is the difference between 73*l* 5 $\frac{1}{4}$ *d* and 19*l* 13*s* 10*d*?
 Ans. 53*l* 6*s* 7 $\frac{1}{4}$ *d*.

* The reason of this Rule will easily appear from what has been said in Simple Subtraction; for the borrowing depends on the same principle, and is only different as the numbers to be subtracted are of different denominations.

Ex. 6.

Ex. 6. A lends to B 100*l*, how much is B in debt after A has taken goods of him to the amount of 73*l* 12*s* 4³/₄*d*?

Ans. 26*l* 7*s* 7¹/₄*d*.

7. Suppose that my rent for half a year is 20*l* 12*s*, and that I have laid out for the land-tax 14*s* 6*d*, and for several repairs 1*l* 3*s* 3¹/₄*d*, what have I to pay of my half-year's rent?

Ans. 18*l* 14*s* 2³/₄*d*.

8. A trader failing, owes to A 35*l* 7*s* 6*d* to B 91*l* 13*s* ¹/₂*d*. to C 53*l* 7¹/₄*d*, to D 87*l* 5*s*, and to E 111*l* 3*s* 5³/₄*d*. When this happened, he had by him in cash 2*l* 7*s* 5*d*, in wares 53*l* 11*s* 10¹/₄*d*, in household furniture 63*l* 17*s* 7³/₄*d*, and in recoverable book-debts 25*l* 7*s* 5*d*. What will his creditors lose by him, suppose these things delivered to them? Ans. 212*l* 5*s* 3¹/₂*d*.

EXAMPLES OF WEIGHTS, MEASURES, &c.

	TROY WEIGHT.				APOTHECARIES' WEIGHT.								
	1.				2.				3.				
	lb	oz	dwt	gr	lb	oz	dwt	gr	lb	oz	dr	scr	gr
From	9	2	12	10	7	10	4	17	73	4	7	0	14
Take	5	4	6	17	3	7	16	12	29	5	3	4	19
	<hr/>				<hr/>				<hr/>				
Rem.	<hr/>				<hr/>				<hr/>				
Proof	<hr/>				<hr/>				<hr/>				

	AVOIRDUPOIS WEIGHT.			LONG MEASURE.								
	4.			5.			6.			7.		
	c	qrs	lb	lb	oz	dr	m	fu	pl	yd	ft	in
From	5	0	17	71	5	9	14	3	17	96	0	4
Take	2	3	10	17	9	18	7	6	11	72	2	9
	<hr/>			<hr/>			<hr/>			<hr/>		
Rem.	<hr/>			<hr/>			<hr/>			<hr/>		
Proof	<hr/>			<hr/>			<hr/>			<hr/>		

	CLOTH MEASURE.			LAND MEASURE.								
	8.			9.			10.			11.		
	yd	qr	nl	yd	qr	nl	ac	ro	p	ac	ro	p
From	17	2	1	9	0	2	17	1	14	57	1	16
Take	9	0	2	7	2	1	16	2	8	22	3	29
	<hr/>			<hr/>			<hr/>			<hr/>		
Rem.	<hr/>			<hr/>			<hr/>			<hr/>		
Proof	<hr/>			<hr/>			<hr/>			<hr/>		

WINE MEASURE.

	12.			13.		
	t	hd gal		hd gal	pt	
From	17	2	23	5	0	4
Take	9	1	36	2	12	6
Rem.	_____			_____		
Proof	_____			_____		

ALE AND BEER MEASURE.

	14.			15.		
	hd gal	pt		hd gal	pt	
	14	29	3	71	16	5
	9	35	7	19	7	1
Rem.	_____			_____		
Proof	_____			_____		

DRY MEASURE.

	16.			17.		
	la	qr	bu	bu	gal	pt
From	9	4	7	13	7	1
Take	6	3	5	9	2	7
Rem.	_____			_____		
Proof	_____			_____		

TIME.

	18.			19.		
	mo	we	da	ds	hrs	min
	71	2	5	114	17	26
	17	1	6	72	10	37
Rem.	_____			_____		
Proof	_____			_____		

20. The line of defence in a certain polygon being 236 yards, and that part of it which is terminated by the curtain and shoulder being 146 yards 1 foot 4 inches; what then was the length of the face of the bastion? Ans. 89 yds 1 ft 8 in.

COMPOUND MULTIPLICATION.

COMPOUND MULTIPLICATION shows how to find the amount of any given number of different denominations repeated a certain proposed number of times; which is performed by the following rule.

SET the multiplier under the lowest number of the multiplicand, and draw a line below it.—Multiply the number in the lowest denomination by the multiplier, and find how many units of the next higher denomination are contained in the product, setting down what remains.—In like manner, multiply the number in the next denomination, and to the product carry or add the units, before found, and find how many units of the next higher denomination are in this amount,

amount, which carry in like manner to the next product, setting down the overplus.—Proceed thus to the highest denomination proposed: so shall the last product, with the several remainders, taken as one compound number, be the whole amount required.—The method of Proof, and the reason of the Rule, are the same as in Simple Multiplication.

EXAMPLES OF MONEY.

1. To find the amount of 8lb of Tea, at 5s 8½*d* per lb.

$$\begin{array}{r}
 \text{s} \quad \text{d} \\
 5 \quad 8\frac{1}{2} \\
 \hline
 \text{£}2 \quad 5 \quad 8 \text{ Answer.} \\
 \hline
 \end{array}$$

- | | | <i>l</i> | <i>s</i> | <i>d</i> |
|---|------|----------|----------|----------|
| 2. 4lb of Tea, at 7s 8 <i>d</i> per lb. | Ans. | 1 | 10 | 8 |
| 3. 6lb of Butter, at 9½ <i>d</i> per lb. | Ans. | 0 | 4 | 9 |
| 4. 7lb of Tobacco, at 1s 8½ <i>d</i> per lb. | Ans. | 0 | 11 | 11½ |
| 5. 9 Stone of Beef, at 2s 7½ <i>d</i> per st. | Ans. | 1 | 1 | 0 |
| 6. 10 cwt of Cheese, at 2 <i>l</i> 17s 10 <i>d</i> per cwt. | Ans. | 28 | 18 | 4 |
| 7. 12 cwt of Sugar, at 3 <i>l</i> 7s 4 <i>d</i> per cwt. | Ans. | 40 | 8 | 0 |

CONTRACTIONS.

I. If the multiplier exceed 12, multiply successively by its component parts, instead of the whole number at once.

EXAMPLES.

1. 15 cwt of Cheese, at 17s 6*d* per cwt.

$$\begin{array}{r}
 \text{l} \quad \text{s} \quad \text{d} \\
 0 \quad 17 \quad 6 \\
 \quad \quad \quad 3 \\
 \hline
 2 \quad 12 \quad 6 \\
 \quad \quad \quad 5 \\
 \hline
 13 \quad 2 \quad 6 \text{ Answer.} \\
 \hline
 \end{array}$$

- | | | | | |
|---|------|----|---|---|
| 2. 20 cwt of Hops, at 4 <i>l</i> 7s 2 <i>d</i> per cwt. | Ans. | 87 | 3 | 4 |
| 3. 24 tons of Hay, at 3 <i>l</i> 7s 6 <i>d</i> per ton. | Ans. | 81 | 0 | 0 |
| 4. 45 ells of Cloth, at 1s 6 <i>d</i> per ell. | Ans. | 3 | 7 | 6 |
- Ex. 5.

	<i>l</i>	<i>s</i>	<i>d</i>
Ex. 5. 63 gallons of Oil, at 2s 3d per gall.	Ans. 7	1	9
6. 70 barrels of Ale, at 1l 4s per barrel	Ans. 84	0	0
7. 84 quarters of Oats, at 1l 12s 8d per qr.	Ans. 137	4	0
8. 96 quarters of Barley, at 1l 3s 4d per qr.	Ans. 112	0	0
9. 120 days' Wages, at 5s 9d per day.	Ans. 34	10	0
10. 144 reams of Paper, at 13s 4d per ream.	Ans. 96	0	0

II. If the multiplier cannot be exactly produced by the multiplication of simple numbers, take the nearest number to it, either greater or less, which can be so produced, and multiply by its parts, as before.—Then multiply the given multiplicand by the difference between this assumed number and the multiplier, and add the product to that before found, when the assumed number is less than the multiplier, but subtract the same when it is greater.

EXAMPLES.

1. 26 yards of Cloth, at 3s 0
- $\frac{3}{4}$
- d per yard.

<i>l</i>	<i>s</i>	<i>d</i>
0	3	0 $\frac{3}{4}$
		5
<hr/>		
0	15	3 $\frac{3}{4}$
		5
<hr/>		
3	16	6 $\frac{3}{4}$
	3	0 $\frac{3}{4}$
<hr/>		
£3	19	7 $\frac{1}{2}$

Answer.

	<i>l</i>	<i>s</i>	<i>d</i>
2. 29 quarters of Corn, at 2l 5s 3 $\frac{1}{4}$ d per qr.	Ans. 65	12	10 $\frac{1}{4}$
3. 53 loads of Hay, at 3l 15s 2d per load.	Ans. 199	3	10
4. 79 bushels of Wheat, at 11s 5 $\frac{3}{4}$ d per bus.	Ans. 45	6	10 $\frac{1}{4}$
5. 97 casks of Beer, at 12s 2d per cask.	Ans. 59	0	2
6. 114 stone of Meat, at 15s 3 $\frac{1}{4}$ d per stone.	Ans. 87	5	7 $\frac{1}{2}$

EXAMPLES OF WEIGHTS AND MEASURES.

1.				2.					3.			
lb	oz	dwt	gr	lb	oz	dr	sc	gr	cwt	qr	lb	oz
28	7	14	10	2	6	3	2	10	29	2	16	14
			5					8				12
<hr/>				<hr/>					<hr/>			

COMPOUND DIVISION.

41

<p style="text-align: center;">4.</p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 25%;">m^{ls}</td> <td style="width: 25%;">f^u</td> <td style="width: 25%;">p^{ls}</td> <td style="width: 25%;">y^{ds}</td> </tr> <tr> <td>22</td> <td>5</td> <td>29</td> <td>6</td> </tr> <tr> <td colspan="3"></td> <td style="border-top: 1px solid black;">4</td> </tr> </table> <hr/>	m ^{ls}	f ^u	p ^{ls}	y ^{ds}	22	5	29	6				4	<p style="text-align: center;">5.</p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 33%;">y^{ds}</td> <td style="width: 33%;">q^{rs}</td> <td style="width: 33%;">n^a</td> </tr> <tr> <td>126</td> <td>3</td> <td>1</td> </tr> <tr> <td colspan="2"></td> <td style="border-top: 1px solid black;">7</td> </tr> </table> <hr/>	y ^{ds}	q ^{rs}	n ^a	126	3	1			7	<p style="text-align: center;">6.</p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 33%;">ac</td> <td style="width: 33%;">ro</td> <td style="width: 33%;">p^o</td> </tr> <tr> <td>28</td> <td>3</td> <td>27</td> </tr> <tr> <td colspan="2"></td> <td style="border-top: 1px solid black;">9</td> </tr> </table> <hr/>	ac	ro	p ^o	28	3	27			9									
m ^{ls}	f ^u	p ^{ls}	y ^{ds}																																						
22	5	29	6																																						
			4																																						
y ^{ds}	q ^{rs}	n ^a																																							
126	3	1																																							
		7																																							
ac	ro	p ^o																																							
28	3	27																																							
		9																																							
<p style="text-align: center;">7.</p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 25%;">t^{uns}</td> <td style="width: 25%;">h^{hd}</td> <td style="width: 25%;">g^{al}</td> <td style="width: 25%;">p^{ts}</td> </tr> <tr> <td>20</td> <td>2</td> <td>26</td> <td>2</td> </tr> <tr> <td colspan="3"></td> <td style="border-top: 1px solid black;">3</td> </tr> </table> <hr/>	t ^{uns}	h ^{hd}	g ^{al}	p ^{ts}	20	2	26	2				3	<p style="text-align: center;">8.</p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 25%;">w^e</td> <td style="width: 25%;">q^r</td> <td style="width: 25%;">b^u</td> <td style="width: 25%;">p^e</td> </tr> <tr> <td>24</td> <td>2</td> <td>5</td> <td>3</td> </tr> <tr> <td colspan="3"></td> <td style="border-top: 1px solid black;">6</td> </tr> </table> <hr/>	w ^e	q ^r	b ^u	p ^e	24	2	5	3				6	<p style="text-align: center;">9.</p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 25%;">m^o</td> <td style="width: 25%;">w^e</td> <td style="width: 25%;">d^a</td> <td style="width: 25%;">h^o</td> <td style="width: 25%;">mⁱⁿ</td> </tr> <tr> <td>172</td> <td>3</td> <td>5</td> <td>16</td> <td>49</td> </tr> <tr> <td colspan="4"></td> <td style="border-top: 1px solid black;">10</td> </tr> </table> <hr/>	m ^o	w ^e	d ^a	h ^o	m ⁱⁿ	172	3	5	16	49					10
t ^{uns}	h ^{hd}	g ^{al}	p ^{ts}																																						
20	2	26	2																																						
			3																																						
w ^e	q ^r	b ^u	p ^e																																						
24	2	5	3																																						
			6																																						
m ^o	w ^e	d ^a	h ^o	m ⁱⁿ																																					
172	3	5	16	49																																					
				10																																					

COMPOUND DIVISION.

COMPOUND DIVISION teaches how to divide a number of several denominations by any given number, or into any number of equal parts ; as follows :

PLACE the divisor on the left of the dividend, as in Simple Division.—Begin at the left-hand, and divide the number of the highest denomination by the divisor, setting down the quotient in its proper place.—If there be any remainder after this division, reduce it to the next lower denomination, which add to the number, if any, belonging to that denomination, and divide the sum by the divisor.—Set down again this quotient, reduce its remainder to the next lower denomination again, and so on through all the denominations to the last.

EXAMPLES OF MONEY.

1. Divide 237l 8s 6d by 2

$$\begin{array}{r} \\ 2 237 \\ \hline \end{array}$$

£118 14 3 the Quotient.

2. Divide

	<i>l</i>	<i>s</i>	<i>d</i>			<i>l</i>	<i>s</i>	<i>d</i>
2. Divide	432	12	$1\frac{3}{4}$	by 3.	Ans.	144	4	$0\frac{1}{2}$
3. Divide	507	3	5	by 4.	Ans.	126	15	$10\frac{1}{4}$
4. Divide	632	7	$6\frac{1}{2}$	by 5.	Ans.	126	9	6
5. Divide	690	14	$3\frac{1}{4}$	by 6.	Ans.	115	2	$4\frac{1}{2}$
6. Divide	705	10	2	by 7.	Ans.	100	15	$8\frac{3}{4}$
7. Divide	760	5	6	by 8.	Ans.	95	0	$8\frac{1}{4}$
8. Divide	761	5	$7\frac{3}{4}$	by 9.	Ans.	84	11	$8\frac{3}{4}$
9. Divide	829	17	10	by 10.	Ans.	82	19	$9\frac{1}{4}$
10. Divide	937	8	$8\frac{3}{4}$	by 11.	Ans.	85	4	5
11. Divide	1145	11	$4\frac{1}{4}$	by 12.	Ans.	95	9	$3\frac{1}{4}$

CONTRACTIONS.

I. If the divisor exceed 12, find what simple numbers, multiplied together, will produce it, and divide by them separately, as in Simple Division, as below.

EXAMPLES.

1. What is Cheese per cwt, if 16 cwt cost 25*l* 14*s* 8*d*?

$$\begin{array}{r} \\ 25 \\ \hline \end{array}$$

$$\begin{array}{r} \\ 6 \\ \hline \end{array}$$

£1 12 2 the Answer.

- | | | | | |
|---|------|----|----------------|----------------|
| 2. If 20 cwt of Tobacco come to }
150 <i>l</i> 6 <i>s</i> 8 <i>d</i> , what is that per cwt? } | Ans. | 7 | 10 | 4 |
| 3. Divide 98 <i>l</i> 8 <i>s</i> by 36 | Ans. | 2 | 14 | 8 |
| 4. Divide 71 <i>l</i> 13 <i>s</i> 10 <i>d</i> by 56. | Ans. | 15 | $7\frac{1}{4}$ | |
| 5. Divide 44 <i>l</i> 4 <i>s</i> by 96. | Ans. | 0 | 9 | $2\frac{1}{2}$ |
| 6. At 31 <i>l</i> 10 <i>s</i> per cwt, how much per lb? | Ans. | 0 | 5 | $7\frac{1}{2}$ |

II. If the divisor cannot be produced by the multiplication of small numbers, divide by the whole divisor at once, after the manner of Long Division, as follows.

EXAMPLES.

1. Divide 59l 6s 3 $\frac{3}{4}$ d by 19.

	<i>l</i>	<i>s</i>	<i>d</i>		<i>l</i>	<i>s</i>	<i>d</i>
19)	59	6	3 $\frac{3}{4}$		(3	2	5 $\frac{1}{4}$ Ans.
	57						
	<hr/>						
	2						
	20						
	<hr/>						
	46	(2					
	38						
	<hr/>						
	8						
	12						
	<hr/>						
	99	(5					
	95						
	<hr/>						
	4						
	4						
	<hr/>						
	19	(1					

- | | <i>l</i> | <i>s</i> | <i>d</i> | | <i>l</i> | <i>s</i> | <i>d</i> |
|--|----------|----------|----------|--|----------|----------|------------------|
| 2. Divide 39 14 5 $\frac{1}{4}$ by 57. | | | | | Ans. 0 | 13 | 11 $\frac{1}{4}$ |
| 3. Divide 125 4 9 by 43. | | | | | Ans. 2 | 18 | 3 |
| 4. Divide 542 7 10 by 97. | | | | | Ans. 5 | 11 | 10 |
| 5. Divide 123 11 2 $\frac{1}{2}$ by 127. | | | | | Ans. 0 | 19 | 5 $\frac{1}{2}$ |

EXAMPLES OF WEIGHTS AND MEASURES.

1. Divide 17 lb 9 oz 0 dwts 2 gr by 7.
Ans. 2 lb 6 oz 8 dwts 14 gr.
2. Divide 17 lb 5 oz 2 dr 1 scr 4 gr by 12.
Ans. 1 lb 5 oz 3 dr 1 scr 12 gr.
3. Divide 178 cwt 3 qrs 14 lb by 53. Ans. 3 cwt 1 qr 14 lb.
4. Divide 144 mi 4 fur 2 po 1 yd 2 ft 0 in by 39.
Ans. 3 mi 5 fur 26 po 0 yds 2 ft 8 in.
5. Divide 534 yds 2 qrs 2 na by 47. Ans. 11 yds 1 qr 2 na.
6. Divide 71 ac 1 ro 33 po by 51. Ans. 1 ac 2 ro 3 po.
7. Divide 7 tu 0 hbds 47 gal 7 pi by 65. Ans. 27 gal 7 pi.
8. Divide 387 la 9 qr by 72. Ans. 5 la 3 qrs 7 bu.
9. Divide 206 mo 4 da by 26. Ans. 7 mo 3 we 5 ds.

THE GOLDEN RULE, OR RULE OF THREE.

THE RULE OF THREE teaches how to find a fourth proportional to three numbers given: for which reason it is sometimes called the Rule of proportion. It is called the Rule of Three, because three terms or numbers are given, to find a fourth. And because of its great and extensive usefulness, it is often called the Golden Rule. This Rule is usually considered as of two kinds, namely, Direct, and Inverse.

The Rule of Three Direct is that in which more requires more, or less requires less. As in this; if 3 men dig 21 yards of trench in a certain time, how much will 6 men dig in the same time? Here more requires more, that is, 6 men, which are more than 3 men, will also perform more work in the same time. Or when it is thus: if 6 men dig 42 yards, how much will 3 men dig in the same time? Here then less requires less, or 3 men will perform proportionably less work than 6 men, in the same time. In both these cases then, the Rule, or the Proportion, is Direct; and the stating must be

thus, As 3 : 21 :: 6 : 42,
or thus, As 6 : 42 :: 3 : 21.

But the Rule of Three Inverse, is when more requires less, or less requires more. As in this: if 3 men dig a certain quantity of trench in 14 hours, in how many hours will 6 men dig the like quantity? Here it is evident that 6 men, being more than 3, will perform an equal quantity of work in less time or fewer hours. Or thus: if 6 men perform a certain quantity of work in 7 hours, in how many hours will 3 men perform the same? Here less requires more, for 3 men will take more hours than 6 to perform the same work. In both these cases then the Rule, or the Proportion, is Inverse; and the stating must be

thus, As 6 : 14 :: 3 : 7,
or thus, As 3 : 7 :: 6 : 14.

And in all these statings, the fourth term is found, by multiplying the 2d and 3d terms together, and dividing the product by the 1st term.

Of the three given numbers; two of them contain the supposition, and the third a demand. And for stating and working questions of these kinds observe the following general Rule:

STATE the question, by setting down in a straight line the three given numbers, in the following manner, viz. so that the 2d term be that number of supposition, which is of the same kind that the answer or fourth term is to be ; making the other number of supposition the 1st term, and the demanding number the 3d term, when the question is in direct proportion ; but contrariwise, the other number of supposition the 3d term, and the demanding number the 1st term, when the question has inverse proportion.

Then, in both cases, multiply the 2d and 3d terms together, and divide the product by the 1st, which will give the answer, or 4th term sought, viz. of the same denomination as the second term.

Note, If the first and third terms consist of different denominations, reduce them both to the same : and if the second term be a compound number, it is mostly convenient to reduce it to the lowest denomination mentioned.—If, after division, there be any remainder, reduce it to the next lower denomination, and divide by the same divisor as before, and the quotient will be of this last denomination. Proceed in the same manner with all the remainders, till they be reduced to the lowest denomination which the second admits of, and the several quotients taken together will be the answer required.

Note also, The reason for the foregoing Rules will appear, when we come to treat of the nature of proportions.—Sometimes two or more statings are necessary, which may always be known from the nature of the question.

EXAMPLES.

1. If 8 yards of Cloth cost 1l 4s, what will 96 yards cost ?

yds	l	s	yds	l	s
As 8	:	1 4	::	96	: 14 8 the Answer
		20			
		—			
		24			
		96			
		—			
		144			
		216			
		—			
	3)	2304			
		—			
		2,0		28,8s	
		—			
		£14 8		Answer.	

Ex. 2. An engineer having raised 100 yards of a certain work in 24 days with 5 men ; how many men must he employ to finish a like quantity of work in 15 days ?

$$\begin{array}{r} \text{ds men} \quad \text{ds men} \\ \text{As } 15 : 5 :: 24 : 8 \text{ Ans.} \\ \quad \quad \quad 5 \\ \quad \quad \quad \text{---} \\ \quad \quad \quad 15) 120 \text{ (8 Answer.} \\ \quad \quad \quad 120 \\ \quad \quad \quad \text{---} \end{array}$$

3. What will 72 yards of cloth cost, at the rate of 9 yards for 5*l* 12*s* ? Ans. 44*l* 16*s*.

4. A person's annual income being 146*l* ; how much is that per day ? Ans. 8*s*.

5. If 3 paces or common steps of a certain person be equal to 2 yards, how many yards will 160 of his paces make ? Ans. 106 yds 2 ft.

6. What length must be cut off a board, that is 9 inches broad, to make a square foot, or as much as 12 inches in length and 12 in breadth contains ? Ans. 16 inches.

7. If 700 men require 22500 rations of bread for a month ; how many rations will a garrison of 1200 men require ? Ans. 36000.

8. If 7 cwt 1 qr of sugar cost 26*l* 10*s* 4*d* ; what will be the price of 4 cwt 2 qrs ? Ans. 159*l* 2*s*.

9. The clothing of a regiment of foot of 750 men amounting to 2831*l* 5*s* ; what will the clothing of a body of 3500 men amount to ? Ans. 13212*l* 10*s*.

10. How many yards of matting, that is 3 ft broad, will cover a floor that is 27 feet long and 20 feet broad ? Ans. 60 yards.

11. What is the value of 6 bushels of coals, at the rate of 1*l* 14*s* 6*d* the chaldron ? Ans. 5*s* 9*d*.

12. If 6352 stones of 3 feet long complete a certain quantity of walling ; how many stones of 2 feet long will raise a like quantity ? Ans. 9528.

13. What must be given for a piece of silver weighing 73 lb 5 oz 15 dwts, at the rate of 5*s* 9*d* per ounce ? Ans. 253*l* 10*s* 0 $\frac{3}{4}$ *d*.

14. A garrison of 536 men having provision for 12 months ; how long will those provisions last, if the garrison be increased to 1124 men ? Ans. 174 days and $\frac{64}{1124}$.

15. What will be the tax upon 763*l* 15*s*, at the rate of 3*s* 6*d* per pound sterling ? Ans. 133*l* 13*s* 1 $\frac{1}{2}$ *d*.

16. A certain work being raised in 12 days, by working 4 hours each day; how long would it have been in raising by working 6 hours per day? Ans. 8 days.

17. What quantity of corn can I buy for 90 guineas, at the rate of 6s the bushel? Ans. 39 qrs 3 bu.

18. A person, failing in trade, owes in all 977*l*; at which time he has, in money, goods, and recoverable debts, 420*l* 6s 3¼*d*; now supposing these things delivered to his creditors, how much will they get per pound? Ans. 8s 7¼*d*.

19. A plain of a certain extent having supplied a body of 3000 horse with forage for 18 days; then how many days would the same plain have supplied a body of 2000 horse? Ans. 27 days.

20. Suppose a gentleman's income is 600 guineas a year, and that he spends 25s 6*d* per day, one day with another; how much will he have saved at the year's end? Ans. 164*l* 12s 6*d*.

21. What cost 30 pieces of lead, each weighing 1 cwt 12lb, at the rate of 16s 4*d* the cwt? Ans. 27*l* 2s 6*d*.

22. The governor of a besieged place having provision for 54 days, at the rate of 1½lb of bread; but being desirous to prolong the siege to 80 days, in expectation of succour, in that case what must the ration of bread be? Ans. 1⅙ lb.

23. At half a guinea per week, how long can I be boarded for 20 pounds? Ans. 38⅙ wks.

24. How much will 75 chaldrons 7 bushels of coals come to, at the rate of 1*l* 13s 6*d* per chaldron? Ans. 125*l* 19s 0⅙*d*.

25. If the penny loaf weigh 8 ounces when the bushel of wheat costs 7s 3*d*, what ought the penny loaf to weigh when the wheat is at 8s 4*d*? Ans. 6 oz 15⅔ dr.

26. How much a year will 173 acres 2 roods 14 poles of land give, at the rate of 1*l* 7s 8*d* per acre? Ans. 240*l* 2s 7⅔*d*.

27. To how much amounts 73 pieces of lead, each weighing 1 cwt 3 qrs 7 lb, at 10*l* 4s per fother of 19½ cwt? Ans. 69*l* 4s 2*d* 1⅙ q.

28. How many yards of stuff, of 3 qrs wide, will line a cloak that is 1¾ yards in length and 3½ yards wide? Ans. 8 yds 0 qrs 2⅔ al.

29. If 5 yards of cloth cost 14s 2*d*, what must be given for 9 pieces, containing each 21 yards 1 quarter? Ans. 27*l* 1s 10½*d*.

30. If a gentleman's estate be worth 2107*l* 12s a year; what may he spend per day, to save 500*l* in the year? Ans. 4*l* 8s 1⅓⅔*d*.

31. Wanting

31. Wanting just an acre of land cut off from a piece which is $13\frac{1}{2}$ poles in breadth, what length must the piece be?

Ans. 11 po. 4 yds. 2 ft. $0\frac{1}{2}\frac{3}{7}$ in.

32. At $7s\ 9\frac{1}{2}d$ per yard, what is the value of a piece of cloth containing 53 ells English 1 qr. Ans. $25l\ 18s\ 1\frac{3}{4}d$.

33. If the carriage of 5 cwt 14 lb for 96 miles be $1l\ 12s\ 6d$; how far may I have 3 cwt 1 qr carried for the same money?

Ans. 151 m 3 fur $3\frac{1}{3}$ pol.

34. Bought a silver tankard, weighing 1 lb 7 oz 14 dwts; what did it cost me at $6s\ 4d$ the ounce? Ans. $6l\ 4s\ 9\frac{1}{5}d$.

35. What is the half year's rent of 547 acres of land, at $15s\ 6d$ the acre? Ans. $211l\ 19s\ 3d$.

36. A wall that is to be built to the height of 36 feet, was raised 9 feet high by 16 men in 6 days; then how many men must be employed to finish the wall in 4 days, at the same rate of working? Ans 72 men.

37. What will be the charge of keeping 20 horses for a year, at the rate of $14\frac{1}{2}d$ per day for each horse?

Ans. $441l\ 0s\ 10d$.

38. If 18 ells of stuff that is $\frac{3}{4}$ yard wide, cost $39s\ 6d$; what will 50 ells, of the same goodness, cost, being yard wide?

Ans. $7l\ 6s\ 3\frac{3}{4}\frac{1}{4}d$.

39. How many yards of paper that is 30 inches wide, will hang a room that is 20 yards in circuit and 9 feet high.

Ans. 72 yards.

40. If a gentleman's estate be worth $384l\ 16s$ a year, and the land-tax be assessed at $2s\ 9\frac{1}{2}d$ per pound, what is his net annual income? Ans. $351l\ 1s\ 9\frac{1}{5}d$.

41. The circumference of the earth is about 24877 miles; at what rate per hour is a person at the middle of its surface carried round, one whole rotation being made in 23 hours 56 minutes? Ans. $1039\frac{1}{3}\frac{5}{9}\frac{4}{9}$ miles.

42. If a person drink 20 bottles of wine per month, when it costs $8s$ a gallon; how many bottles per month may he drink, without increasing the expence, when wine costs $10s$ the gallon? Ans. 16 bottles.

43. What cost 43 qrs 5 bushels of corn, at $1l\ 8s\ 6d$ the quarter. Ans. $62l\ 3s\ 3\frac{1}{4}d$.

44. How many yards of canvas that is ell wide will line 50 yards of say that is 3 quarters wide? Ans. 30 yds.

45. If an ounce of gold cost 4 guineas, what is the value of a grain? Ans. $2\frac{1}{10}d$.

46. If 3 cwt of tea cost $40l\ 12s$; at how much a pound must it be retailed, to gain $10l$ by the whole? Ans. $3\frac{4}{3}\frac{4}{8}s$.

COMPOUND PROPORTION.

COMPOUND PROPORTION shows how to resolve such questions as require two or more statings by Simple Proportion ; and these may be either Direct or Inverse.

In these questions, there is always given an odd number of terms, either five or seven, or nine, &c. These are distinguished into terms of supposition, and terms of demand, there being always one term more of the former than of the latter, which is of the same kind with the answer sought. The method is thus :

SET down in the middle place that term of supposition which is of the same kind with the answer sought.—Take one of the other terms of supposition, and one of the demanding terms which is of the same kind with it ; then place one of them for a first term, and the other for a third, according to the directions given in the Rule of Three.—Do the same with another term of supposition, and its corresponding demanding term ; and so on if there be more terms of each kind ; setting the numbers under each other which fall all on the left-hand side of the middle term, and the same for the others on the right-hand side.—Then, to work

By several Operations.—Take the two upper terms and the middle term, in the same order as they stand, for the first Rule-of-Three question to be worked, whence will be found a fourth term. Then take this fourth number, so found, for the middle term of a second Rule-of-Three question, and the next two under terms in the general stating, in the same order as they stand, finding a fourth term for them. And so on, as far as there are any numbers in the general stating, making always the fourth number, resulting from each simple stating, to be the second term in the next following one. So shall the last resulting number be the answer to the question.

By one Operation.—Multiply together all the terms standing under each other, on the left-hand side of the middle term ; and, in like manner, multiply together all those on the right-hand side of it. Then multiply the middle term by the latter product, and divide the result by the former product ; so shall the quotient be the answer sought.

EXAMPLES.

1. How many men can complete a trench of 135 yards long in 8 days, when 16 men can dig 54 yards in 6 days ?

General Stating.

yds	54	:	16	::	135	yds
days	8				6	days

432

810

16

4860

81 men

432) 12960 (30 Ans. by one operation.

1296

0

The same by two Operations.

1st.
As 54 : 16 :: 135 : 40

16

810

135

54) 2160 (40

216

0

2d.
As 8 : 40 :: 6 : 30

6

8) 240 (30 Ans.

24

0

2. If 100*l* in one year gain 5*l* interest, what will be the interest of 750*l* for 7 years ? Ans. 262*l* 10*s*.

3. If a family of 8 persons expend 200*l* in 9 months ; how much will serve a family of 18 people 12 months ? Ans. 300*l*.

4. If 27*s* be the wages of 4 men for 7 days ; what will be the wages of 14 men for 10 days ? Ans. 6*l* 15*s*.

5. If a footman travel 130 miles in 3 days, when the days are 12 hours long ; in how many days, of 10 hours each, may he travel 360 miles ? Ans. 9 $\frac{2}{3}$ days.

Ex. 6.

Ex. 6. If 120 bushels of corn can serve 14 horses 56 days ; how many days will 94 bushels serve 6 horses ?

Ans. $102\frac{1}{5}$ days.

7. If 3000 lb. of beef serve 340 men 15 days ; how many lbs will serve 120 men for 25 days ? Ans. 1764 lb $11\frac{1}{5}$ oz.

8. If a barrel of beer be sufficient to last a family of 8 persons 12 days ; how many barrels will be drank by 16 persons in the space of a year ? Ans. $60\frac{5}{6}$ barrels.

9. If 180 men, in 6 days of 10 hours each, can dig a trench 200 yards long, 3 wide, and 2 deep ; in how many days, of 8 hours long, will 100 men dig a trench of 360 yards long, 4 wide, and 3 deep ? Ans. 15 days.

OF VULGAR FRACTIONS.

A FRACTION, or broken number, is an expression of a part, or some parts, of something considered as a whole.

It is denoted by two numbers, placed one below the other, with a line between them :

Thus, -	3 numerator	}	which is named 3-fourths.
	4 denominator		

The denominator, or number placed below the line, shows how many equal parts the whole quantity is divided into ; and it represents the Divisor in Division.—And the Numerator, or number set above the line, shows how many of these parts are expressed by the Fraction ; being the remainder after division.—Also, both these numbers are, in general, named the Terms of the Fraction.

Fractions are either Proper, Improper, Simple, Compound, or Mixed.

A Proper Fraction, is when the numerator is less than the denominator ; as, $\frac{1}{2}$, or $\frac{2}{3}$, or $\frac{3}{5}$, &c.

An Improper Fraction, is when the numerator is equal to, or exceeds, the denominator ; as, $\frac{3}{3}$, or $\frac{5}{4}$, or $\frac{7}{5}$, &c.

A Simple Fraction, is a single expression, denoting any number of parts of the integer ; as, $\frac{2}{3}$, or $\frac{3}{4}$.

A Compound Fraction, is the fraction of a fraction, or several fractions connected with the word *of* between them ; as, $\frac{1}{2}$ of $\frac{2}{3}$, or $\frac{3}{5}$ of $\frac{5}{6}$ of 3, &c.

A Mixed Number, is composed of a whole number and a fraction together ; as, $3\frac{1}{4}$, or $12\frac{2}{3}$, &c.

A whole

A whole or integer number may be expressed like a fraction, by writing 1 below it, as a denominator ; so 3 is $\frac{3}{1}$, or 4 is $\frac{4}{1}$, &c.

A fraction denotes division ; and its value is equal to the quotient obtained by dividing the numerator by the denominator ; so $\frac{12}{4}$ is equal to 3, and $\frac{20}{5}$ is equal to 4.

Hence then, if the numerator be less than the denominator, the value of the fraction is less than 1. But if the numerator be the same as the denominator, the fraction is just equal to 1. And if the numerator be greater than the denominator, the fraction is greater than 1.

REDUCTION OF VULGAR FRACTIONS.

REDUCTION of Vulgar Fractions, is the bringing them out of one form or denomination into another ; commonly to prepare them for the operations of Addition, Subtraction, &c. of which there are several cases.

PROBLEM.

To find the Greatest Common Measure of Two or more Numbers.

THE Common Measure of two or more numbers, is that number which will divide them both without remainder ; so, 3 is a common measure of 18 and 24 ; the quotient of the former being 6, and of the latter 8. And the greatest number that will do this, is the greatest common measure ; so 6 is the greatest common measure of 18 and 24 ; the quotient of the former being 3, and of the latter 4, which will not both divide further.

RULE.

If there be two numbers only ; divide the greater by the less ; then divide the divisor by the remainder ; and so on, dividing always the last divisor by the last remainder, till nothing remains ; so shall the last divisor of all be the greatest common measure sought.

When there are more than two numbers, find the greatest common measure of two of them, as before ; then do the same for that common measure and another of the numbers ; and so
on,

on, through all the numbers ; so will the greatest common measure last found be the answer.

EXAMPLES.

1. To find the greatest common measure of 1908, 936, and 630.

$$\begin{array}{r} 936) 1908 \text{ (2} \\ \underline{1872} \end{array}$$

So that 36 is the greatest common measure of 1908 and 936.

$$\begin{array}{r} 36) 936 \text{ (26} \\ \underline{72} \end{array}$$

$$\begin{array}{r} 216 \\ \underline{216} \end{array}$$

$$\begin{array}{r} 36) 630 \text{ (17} \\ \underline{36} \end{array}$$

$$\begin{array}{r} 270 \\ \underline{252} \end{array}$$

$$\begin{array}{r} 18) 36 \text{ (2} \\ \underline{36} \end{array}$$

Hence then 18 is the answer required.

2. What is the greatest common measure of 246 and 372 ?

Ans. 6.

3. What is the greatest common measure of 324, 612, and 1032 ?

Ans. 12.

CASE. I.

To Abbreviate or Reduce Fractions to their Lowest Terms.

*DIVIDE the terms of the given fraction by any number that will divide them without a remainder ; then divide these quotients

* That dividing both the terms of the fraction by the same number, whatever it be, will give another fraction equal to the former, is evident. And when these divisions are performed as often as can be done, or when the common divisor is the greatest possible, the terms of the resulting fraction must be the least possible.

Note 1. Any number ending with an even number, or a cipher, is divisible, or can be divided, by 2.

2. Any number ending with 5, or 0, is divisible by 5.

3. If

quotients again in the same manner ; and so on, till it appears that there is no number greater than 1 which will divide them ; then the fraction will be in its lowest terms.

Or, divide both the terms of the Fraction by their greatest common measure at once, and the quotients will be the terms of the fraction required, of the same value as at first.

EXAMPLES.

1. Reduce $\frac{216}{288}$ to its least terms.

$$\frac{216}{288} = \frac{72}{96} = \frac{36}{48} = \frac{12}{16} = \frac{6}{8} = \frac{3}{4}, \text{ the answer.}$$

Or thus :

$$\begin{array}{r} 216) 288 \quad (1 \\ \underline{216} \end{array}$$

$$\begin{array}{r} 72) 216 \quad (3 \\ \underline{216} \end{array}$$

Therefore 72 is the greatest common measure ; and 72) $\frac{216}{288} = \frac{3}{4}$ the Answer, the same as before.

2. Reduce

3. If the right hand place of any number be 0, the whole is divisible by 10 ; if there be two ciphers, it is divisible by 100 ; if three ciphers by 1000 : and so on ; which is only cutting off those ciphers.

4. If the two right hand figures of any number be divisible by 4, the whole is divisible by 4. And if the three right hand figures be divisible by 8, the whole is divisible by 8. And so on.

5. If the sum of the digits in any number be divisible by 3, or by 9, the whole is divisible by 3, or by 9.

6. If the right hand digit be even, and the sum of all the digits be divisible by 6, then the whole is divisible by 6.

7. A number is divisible by 11, when the sum of the 1st, 3d, 5th, &c. or all the odd places, is equal to the sum of the 2d, 4th, 6th, &c. or of all the even places of digits.

8. If a number cannot be divided by some quantity less than the square root of the same, that number is a prime, or cannot be divided by any number whatever.

9. All prime numbers, except 2 and 5, have either 1, 3, 7, or 9, in the place of units ; and all other numbers are composite or can be divided.

10. When

2. Reduce $1\frac{2}{8}\frac{5}{0}$ to its lowest terms. Ans. $\frac{1}{4}$
 3. Reduce $1\frac{3}{2}\frac{6}{4}$ to its lowest terms. Ans. $\frac{2}{3}$
 4. Reduce $5\frac{2}{6}\frac{5}{0}$ to its lowest terms. Ans. $\frac{5}{6}$

CASE II.

To Reduce a Mixed Number to its Equivalent Improper Fraction.

* MULTIPLY the integer or whole number by the denominator of the fraction, and to the product add the numerator; then set that sum above the denominator for the fraction required.

EXAMPLES.

1. Reduce $23\frac{2}{5}$ to a fraction.

$$\begin{array}{r} 23 \\ 5 \\ \hline \end{array}$$

$$\begin{array}{r} 115 \\ 2 \\ \hline \end{array}$$

$$\begin{array}{r} 117 \\ 5 \\ \hline \end{array}$$

Or,

$$\frac{(23 \times 5) + 2}{5} = \frac{117}{5}, \text{ the Answer.}$$

2. Reduce $12\frac{7}{6}$ to a fraction. Ans. $1\frac{1}{2}$
 3. Reduce $14\frac{7}{10}$ to a fraction. Ans. $1\frac{7}{10}$
 4. Reduce $183\frac{5}{21}$ to a fraction. Ans. $3\frac{84}{21}$

10. When numbers, with the sign of addition or subtraction between them, are to be divided by any number, then each of those numbers must be divided by it. Thus $\frac{10+8-4}{2} = 5+4-2 = 7$.

11. But if the numbers have the sign of multiplication between them, only one of them must be divided. Thus,

$$\frac{10 \times 8 \times 3}{6 \times 2} = \frac{10 \times 4 \times 3}{6 \times 1} = \frac{10 \times 4 \times 1}{2 \times 1} = \frac{10 \times 2 \times 1}{1 \times 1} = \frac{20}{1} = 20$$

* This is no more than first multiplying a quantity by some number, and then dividing the result back again by the same; which it is evident does not alter the value; for any fraction represents a division of the numerator by the denominator.

CASE III.

To Reduce an Improper Fraction to its Equivalent Whole or Mixed Number.

*DIVIDE the numerator by the denominator, and the quotient will be the whole or mixed number sought.

EXAMPLES.

1. Reduce $\frac{12}{3}$ to its equivalent number.

Here $\frac{12}{3}$ or $12 \div 3 = 4$, the Answer.

2. Reduce $\frac{15}{7}$ to its equivalent number.

Here $\frac{15}{7}$ or $15 \div 7 = 2\frac{1}{7}$ the Answer.

3. Reduce $\frac{749}{17}$ to its equivalent number.

Thus $17 \overline{) 749} (44\frac{1}{17}$

68

—

69

68

—

1

—

So that $\frac{749}{17} = 44\frac{1}{17}$, the Answer.

4. Reduce $\frac{56}{7}$ to its equivalent number.

Ans. 8.

5. Reduce $\frac{1282}{5}$ to its equivalent number.

Ans. $54\frac{2}{5}$.

6. Reduce $\frac{218}{17}$ to its equivalent number.

Ans. $17\frac{1}{17}$.

CASE. IV.

To Reduce a Whole number to an Equivalent Fraction, having a Given denominator.

† MULTIPLY the whole number by the given denominator ; then set the product over the said denominator, and it will form the fraction required.

* This rule is evidently the reverse of the former ; and the reason of it is manifest from the nature of Common Division.

† Multiplication and Division being here equally used, the result must be the same as the quantity first proposed.

EXAMPLES.

EXAMPLES.

1. Reduce 9 to a fraction whose denominator shall be 7.
Here $9 \times 7 = 63$: then $\frac{63}{7}$ is the Answer ;
For $\frac{63}{7} = 63 \div 7 = 9$, the Proof.
2. Reduce 12 to a fraction whose denominator shall be 13.
Ans. $\frac{156}{13}$.
3. Reduce 27 to a fraction whose denominator shall be 11.
Ans. $\frac{297}{11}$.

CASE V.

To Reduce a Compound Fraction to an Equivalent Simple One.

* MULTIPLY all the numerators together for a numerator, and all the denominators together for a denominator, and they will form the simple fraction sought.

When part of the compound fraction is a whole or mixed number, it must first be reduced to a fraction by one of the former cases.

And, when it can be done, any two terms of the fraction may be divided by the same number, and the quotients used instead of them. Or, when there are terms that are common, they may be omitted, or cancelled.

EXAMPLES.

1. Reduce $\frac{1}{2}$ of $\frac{2}{3}$ of $\frac{3}{4}$ to a simple fraction.

$$1 \times 2 \times 3 = 6 \quad \frac{1}{4}$$

Here $\frac{6}{24} = \frac{1}{4}$, the Answer

$$\frac{2 \times 3 \times 4}{24} = \frac{1}{4}$$

$$\frac{1 \times 2 \times 3}{6} = \frac{1}{4}$$

Or, $\frac{2 \times 3 \times 4}{2 \times 3 \times 4} = \frac{1}{4}$, by cancelling the 2's and 3's.

2. Reduce $\frac{2}{3}$ of $\frac{3}{5}$ of $\frac{10}{11}$ to a simple fraction.

$$2 \times 3 \times 10 = 60 \quad \frac{12}{11}$$

Here $\frac{60}{660} = \frac{12}{11}$, the Answer.

$$\frac{3 \times 5 \times 11}{165} = \frac{12}{11}$$

* The truth of this Rule may be shown as follows: Let the compound fraction be $\frac{2}{3}$ of $\frac{3}{5}$. Now $\frac{1}{3}$ of $\frac{3}{5}$ is $\frac{3}{5} \div 3$, which is $\frac{1}{5}$; consequently $\frac{2}{3}$ of $\frac{3}{5}$ will be $\frac{2}{5}$ or $\frac{2}{5}$; that is the numerators are multiplied together, and also the denominators, as in the Rule. When the compound fraction consists of more than two single ones; having first reduced two of them as above, then the resulting fraction and a third will be the same as a compound fraction of two parts; and so on to the last of all.

2. Reduce

Or, $\frac{2 \times 3 \times 10}{3 \times 5 \times 11} = \frac{4}{11}$, the same as before, by cancelling

the 3's, and dividing by 5's.

- | | |
|--|------------------------|
| 3. Reduce $\frac{3}{4}$ of $\frac{4}{5}$ to a simple fraction. | Ans. $\frac{12}{20}$. |
| 4. Reduce $\frac{2}{3}$ of $\frac{3}{5}$ of $\frac{5}{6}$ to a simple fraction. | Ans. $\frac{2}{6}$. |
| 5. Reduce $\frac{2}{5}$ of $\frac{5}{8}$ of $3\frac{1}{2}$ to a simple fraction. | Ans. $\frac{7}{8}$. |
| 6. Reduce $\frac{2}{4}$ of $\frac{5}{8}$ of $\frac{7}{2}$ of 4 to a simple fraction. | Ans. $\frac{5}{2}$. |
| 7. Reduce 2 and $\frac{2}{5}$ of $\frac{5}{6}$ to a fraction. | Ans. $\frac{7}{3}$. |

CASE VI.

To Reduce Fractions of Different Denominators, to Equivalent Fractions having a Common Denominator.

* MULTIPLY each numerator by all the denominators except its own, for the new numerators : and multiply all the denominators together for a common denominator.

Note. It is evident that in this and several other operations, when any of the proposed quantities are integers, or mixed numbers, or compound fractions, they must first be reduced, by their proper Rules, to the form of simple fractions.

EXAMPLES.

1. Reduce $\frac{1}{2}$, $\frac{2}{3}$, and $\frac{3}{4}$, to a common denominator.
- | | | |
|----------------------------|-------------------------|---------------|
| $1 \times 3 \times 4 = 12$ | the new numerator for | $\frac{1}{2}$ |
| $2 \times 2 \times 4 = 16$ | ditto | $\frac{2}{3}$ |
| $3 \times 2 \times 3 = 18$ | ditto | $\frac{3}{4}$ |
| $2 \times 3 \times 4 = 24$ | the common denominator. | |

Therefore the equivalent fractions are $\frac{12}{24}$, $\frac{16}{24}$, and $\frac{18}{24}$.

Or the whole operation of multiplying may be best performed mentally, only setting down the results and given fractions thus : $\frac{1}{2}, \frac{2}{3}, \frac{3}{4} = \frac{12}{24}, \frac{16}{24}, \frac{18}{24} = \frac{6}{12}, \frac{8}{12}, \frac{9}{12}$ by abbreviation.

2. Reduce $\frac{2}{3}$ and $\frac{5}{6}$ to fractions of a common denominator.
- Ans. $\frac{16}{24}, \frac{20}{24}$.

3. Reduce $\frac{2}{3}$, $\frac{3}{5}$, and $\frac{3}{4}$, to a common denominator.
- Ans. $\frac{40}{60}, \frac{36}{60}, \frac{45}{60}$.

4. Reduce $\frac{5}{6}$, $2\frac{2}{3}$ and 4, to a common denominator.
- Ans. $\frac{25}{30}, \frac{78}{30}, \frac{120}{30}$.

Note I. When the denominators of two given fractions have a common measure, let them be divided by it; then

* This is evidently no more than multiplying each numerator and its denominator by the same quantity, and consequently the value of the fraction is not altered.

multiply the terms of each given fraction by the quotient arising from the other's denominator.

Ex. $\frac{2}{5}$, and $\frac{4}{35} = \frac{2 \cdot 1}{175}$ and $\frac{20}{175}$, by multiplying the former by 7, and the latter by 5.

2. When the less denominator of two fractions exactly divides the greater, multiply the terms of that which has the less denominator by the quotient.

Ex. $\frac{3}{7}$ and $\frac{5}{14} = \frac{6}{14}$ and $\frac{5}{14}$, by mult. the former by 2.

3. When more than two fractions are proposed, it is sometimes convenient, first to reduce two of them to a common denominator; then these and a third; and so on till they be all reduced to their least common denominator.

Ex. $\frac{2}{3}$ and $\frac{3}{4}$ and $\frac{7}{8} = \frac{2}{8}$ and $\frac{6}{8}$ and $\frac{7}{8} = \frac{16}{24}$ and $\frac{18}{24}$ and $\frac{21}{24}$.

CASE VII.

To find the value of a Fraction in parts of the Integer.

MULTIPLY the integer by the numerator, and divide the product by the denominator, by Compound Multiplication and Division, if the integer be a compound quantity.

Or, if it be a single integer, multiply the numerator by the parts in the next inferior denomination, and divide the product by the denominator. Then, if any thing remains, multiply it by the parts in the next inferior denomination, and divide by the denominator as before; and so on as far as necessary; so shall the quotients, placed in order, be the value of the fraction required.*

* The numerator of a fraction being considered as a remainder, in Division, and the denominator as the divisor, this rule is of the same nature as Compound Division, or the valuation of remainders in the Rule of Three, before explained.

Note, by the Editor.—Fractions may be reduced to their least common denominator as follows.

Let 24, 27, 30, 32, 36, 40, 45, 48 be the denominators: reduce each denominator into the product of the powers of its prime factors, and the given numbers become $2^3 \times 3$, 3^3 , $2 \times 3 \times 5$, 2^5 , $2^2 \times 3^2$, $2^3 \times 5$, $3^2 \times 5$, $2^4 \times 3$: now take the highest power of each prime factor and we have 2^5 , 3^3 , 5 ; the product of which $2^5 \times 3^3 \times 5 = 32 \times 27 \times 5 = 4320$, is the least common denominator required. Again, let 2, 3, 4, 5, 6, 7, 8, 9, 10 be the denominators. In this case the powers of the primes in each number are 2, 3, 2^2 , 5, 2×3 , 7, 2^3 , 3^2 , 2×5 ; and the highest powers of the primes are 2^3 , 3^2 , 5, 7, of which the product is $2^3 \times 3^2 \times 5 \times 7 = 8 \times 9 \times 5 \times 7 = 63 \times 40 = 2520$, which is the least common denominator.

This method is advantageous when the prime factors are easily discovered, in other cases we may proceed in the following manner. Find the greatest common divisor of the first and second given numbers; divide the product of the first and second given numbers by this greatest common divisor, and call the quotient c: in like manner divide the product of c and the third given number by their greatest common divisor, and call the quotient d: proceed in like manner with d and the fourth given number, and the last number thus found will be the

EXAMPLES.

1. What is the $\frac{1}{3}$ of 2l 6s ? By the former part of the Rule

$$\begin{array}{r} 2l \ 6s \\ \quad 4 \\ \hline 5) \ 9 \ 4 \\ \hline \text{Ans.} \ 1l \ 16s \ 9d \ 2\frac{2}{3}q. \end{array}$$

2. What is the value of $\frac{2}{3}$ of 1l ? By the 2d part of the Rule,

$$\begin{array}{r} 2 \\ 20 \\ \hline 3) \ 40 \ (13s \ 4d \ \text{Ans.} \\ \hline 1 \\ 12 \\ \hline 3) \ 12 \ (4d \\ \hline \end{array}$$

3. Find the value of $\frac{3}{8}$ of a pound sterling. Ans. 7s 6d.
 4. What is the value of $\frac{2}{3}$ of a guinea ? Ans. 4s 8d
 5. What is the value of $\frac{3}{4}$ of a half crown ? Ans. 1s 10 $\frac{1}{2}$ d.
 6. What is the value of $\frac{2}{5}$ of 4s 10d ? Ans. 1s 4 $\frac{1}{5}$ d.
 7. What is the value of $\frac{4}{5}$ lb troy ? Ans. 9 oz 12 dwts.
 8. What is the value of $\frac{5}{16}$ of a cwt ? Ans. 1 qr 7 lb.

the least common multiple of the given numbers; that is, the least common denominator of the given fractions.

Ex. 1. Let 24, 27, 30, 32, 36, 40, 45, 48 be the given numbers. The greatest common divisor of 24 and 27 is found by the common rule to be 3, then $\frac{24 \times 27}{3} = 216 = c$. Again, the greatest common divisor of 216 and 30 is found

to be 6, and therefore $d = \frac{216 \times 30}{6} = 1080$. Again the greatest common divisor of 1080 and 32 is 3, therefore $e = \frac{1080 \times 32}{3} = 4320$. Farther, the greatest

common divisor of 4320 and 36 is 36, whence $f = \frac{4320 \times 36}{36} = 4320$. In like

manner $g = \frac{4320 \times 40}{40} = 4320$, and $h = \frac{4320 \times 45}{45} = 4320$, and lastly $k =$

$\frac{4320 \times 48}{48} = 4320 =$ the least common multiple of the given numbers.

Ex. 2. Let 2, 3, 4, 5, 6, 7, 8, 9, 10 be the given numbers. Here $c = \frac{2 \times 3}{1} = 6$, $d = \frac{6 \times 4}{2} = 12$, $e = \frac{12 \times 5}{1} = 60$, $f = \frac{60 \times 6}{6} = 60$, $g = \frac{60 \times 7}{1} = 420$, $h = \frac{420 \times 8}{4} = 840$, $k = \frac{840 \times 9}{3} = 2520$, and lastly $L = \frac{2520 \times 10}{10} = 2520 =$ the least common multiple required.

This general rule may be expressed as follows.

Divide the first by the greatest common measure of the first and second, and multiply the quotient by the second, and call the product c : divide c by the greatest common measure of c and the third given number, and multiply the quotient by the third, call this product d : in like manner proceed with d and the fourth given number, and the last product will be the least common multiple required.

9. What

9. What is the value of $\frac{7}{8}$ of an acre? Ans. 3 ro. 20 po.
 10. What is the value of $\frac{3}{10}$ of a day? Ans. 7 hrs 12 min.

CASE VIII.

To Reduce a Fraction from one Denomination to another.

* CONSIDER how many of the less denomination make one of the greater; then multiply the numerator by that number, if the reduction be to a less name, but multiply the denominator, if to a greater.

EXAMPLES.

1. Reduce $\frac{3}{8}$ of a pound to the fraction of a penny.
 $\frac{3}{8} \times \frac{20}{1} \times \frac{12}{1} = 4\frac{8}{8} = 1\frac{6}{8}$, the Answer.
2. Reduce $\frac{5}{7}$ of a penny to the fraction of a pound.
 $\frac{5}{7} \times \frac{1}{12} \times \frac{1}{20} = \frac{1}{336}$ the answer.
3. Reduce $\frac{1}{15}l$ to the fraction of a penny. Ans. $\frac{33}{1}d$.
4. Reduce $\frac{2}{5}q$ to the fraction of a pound Ans. $\frac{1}{240}$.
5. Reduce $\frac{2}{7}$ cwt to the fraction of a lb. Ans. $\frac{32}{1}$.
6. Reduce $\frac{3}{5}$ dwt to the fraction of a lb troy. Ans. $\frac{1}{40}$.
7. Reduce $\frac{3}{8}$ crown to the fraction of a guinea Ans. $\frac{5}{8}$.
8. Reduce $\frac{5}{8}$ half-crown to the fract. of a shilling. Ans. $\frac{2}{1}$.
9. Reduce 2s 6d to the fraction of a £. Ans. $\frac{1}{8}$.
10. Reduce 17s 7d $3\frac{3}{5}q$ to the fraction of a £.

ADDITION OF VULGAR FRACTIONS.

If the fractions have a common denominator; add all the numerators together, then place the sum over the common denominator, and that will be the sum of the fractions required.

† If the proposed fractions have not a common denominator, they must be reduced to one. Also compound fractions

* This is the same as the Rule of Reduction in whole numbers from one denomination to another.

† Before fractions are reduced to a common denominator, they are quite dissimilar, as much as shillings and pence are, and therefore cannot be incorporated with one another, any more than these can. But when they are reduced to a common denominator, and made parts of the same thing, their sum, or difference, may then be as properly expressed by the sum or difference of the numerators, as the sum or difference of any two quantities whatever, by the sum or difference of their individuals. Whence the reason of the Rule is manifest, both for Addition and Subtraction.

When

must be reduced to simple ones, and fractions of different denominations to those of the same denomination. Then add the numerators as before. As to mixed numbers, they may either be reduced to improper fractions, and so added with the others; or else the fractional parts only added, and the integers united afterwards.

EXAMPLES.

1. To add $\frac{3}{5}$ and $\frac{4}{5}$ together.
Here $\frac{3}{5} + \frac{4}{5} = \frac{7}{5} = 1\frac{2}{5}$, the Answer.
2. To add $\frac{3}{5}$ and $\frac{5}{6}$ together.
 $\frac{3}{5} + \frac{5}{6} = \frac{13}{30} + \frac{25}{30} = \frac{38}{30} = 1\frac{13}{30}$, the Answer.
3. To add $\frac{5}{8}$ and $7\frac{1}{2}$ and $\frac{1}{3}$ of $\frac{3}{4}$ together.
 $\frac{5}{8} + 7\frac{1}{2} + \frac{1}{3}$ of $\frac{3}{4} = \frac{5}{8} + 1\frac{1}{2} + \frac{1}{4} = \frac{5}{8} + \frac{6}{8} + \frac{2}{8} = \frac{13}{8} = 1\frac{5}{8}$.
4. To add $\frac{3}{7}$ and $\frac{6}{7}$ together. Ans. $1\frac{9}{7}$.
5. To add $\frac{3}{4}$ and $\frac{5}{9}$ together. Ans. $1\frac{11}{36}$.
6. Add $\frac{2}{7}$ and $\frac{3}{14}$ together. Ans. $\frac{9}{14}$.
7. What is the sum of $\frac{2}{3}$ and $\frac{3}{5}$ and $\frac{5}{7}$? Ans. $1\frac{110}{105}$.
8. What is the sum of $\frac{5}{9}$ and $\frac{3}{5}$ and $2\frac{1}{6}$? Ans. $3\frac{29}{90}$.
9. What is the sum of $\frac{3}{5}$ and $\frac{4}{5}$ of $\frac{1}{3}$ and $9\frac{3}{6}$? Ans. $10\frac{1}{6}$.
10. What is the sum of $\frac{2}{3}$ of a pound and $\frac{5}{9}$ of a shilling?
Ans. $12\frac{5}{9}s$ or $13s$ $10d$ $2\frac{2}{3}q$.
11. What is the sum of $\frac{3}{5}$ of a shilling and $\frac{4}{15}$ of a penny?
Ans. $\frac{11}{15}d$ or $7d$ $1\frac{1}{3}q$.
12. What is the sum of $\frac{1}{7}$ of a pound, and $\frac{2}{9}$ of a shilling, and $\frac{5}{12}$ of penny?
Ans. $\frac{216}{1080}s$ or $3s$ $1d$ $1\frac{1}{2}q$.

SUBTRACTION OF VULGAR FRACTIONS.

PREPARE the fractions the same as for Addition, when necessary; then subtract the one numerator from the other, and set the remainder over the common denominator, for the difference of the fractions sought.

EXAMPLES

1. To find the difference between $\frac{5}{6}$ and $\frac{1}{6}$.
Here $\frac{5}{6} - \frac{1}{6} = \frac{4}{6} = \frac{2}{3}$, the Answer.
2. To find the difference between $\frac{3}{4}$ and $\frac{5}{9}$.
 $\frac{3}{4} - \frac{5}{9} = \frac{27}{36} - \frac{20}{36} = \frac{7}{36}$, the Answer.

When several fractions are to be collected, it is commonly best first to add two of them together that most easily reduce to a common denominator; then add their sum and a third, and so on.

3. What

MULTIPLICATION OF VULGAR FRACTIONS. 63

3. What is the difference between $\frac{5}{12}$ and $\frac{7}{12}$? Ans. $\frac{1}{6}$.
4. What is the difference between $\frac{2}{13}$ and $\frac{4}{39}$? Ans. $\frac{2}{39}$.
5. What is the difference between $\frac{2}{12}$ and $\frac{7}{13}$? Ans. $\frac{19}{156}$.
6. What is the diff. between $5\frac{2}{3}$ and $\frac{2}{7}$ of $4\frac{1}{6}$? Ans. $4\frac{2}{105}$.
7. What is the difference between $\frac{5}{6}$ of a pound, and $\frac{2}{3}$ of $\frac{2}{4}$ of a shilling? Ans. $\frac{19}{91}$ s or 10s 7d $1\frac{1}{2}$ q.
8. What is the difference between $\frac{2}{7}$ of $5\frac{1}{6}$ of a pound, and $\frac{2}{5}$ of a shilling? Ans. $\frac{2}{21}$ or 1l 8s $11\frac{2}{3}$ d.

MULTIPLICATION OF VULGAR FRACTIONS.

* REDUCE mixed numbers, if there be any, to equivalent fractions; then multiply all the numerators together for a numerator, and all the denominators together for a denominator, which will give the product required.

EXAMPLES:

1. Required the product of $\frac{3}{4}$ and $\frac{2}{6}$.

Here $\frac{3}{4} \times \frac{2}{6} = \frac{6}{24} = \frac{1}{4}$, the Answer.

Or $\frac{3}{4} \times \frac{2}{6} = \frac{1}{2} \times \frac{1}{3} = \frac{1}{6}$.

2. Required the continual product of $\frac{2}{3}$, $3\frac{1}{2}$, 5, and $\frac{3}{4}$ of $\frac{2}{5}$.

Here $\frac{2}{3} \times 13 \frac{5}{2} \times 3 \times 13 \times 3 \frac{39}{4} = 4\frac{7}{4}$ Ans.

3. Required the product of $\frac{2}{7}$ and $\frac{5}{8}$. Ans. $\frac{5}{28}$.

4. Required the product of $\frac{4}{15}$ and $\frac{5}{24}$. Ans. $\frac{1}{18}$.

5. Required the product of $\frac{3}{7}$, $\frac{4}{9}$, and $\frac{14}{15}$. Ans. $\frac{8}{45}$.

* Multiplication of any thing by a fraction, implies the taking some part of parts of the thing; it may therefore be truly expressed by a compound fraction, which is resolved by multiplying together the numerators and denominators.

Note, A Fraction is best multiplied by an integer, by dividing the denominator by it; but if it will not exactly divide, then multiply the numerator by it.

6. Required

6. Required the product of $\frac{1}{2}$, $\frac{2}{3}$, and 3.
 7. Required the product of $\frac{7}{9}$, $\frac{3}{5}$, and $4\frac{5}{14}$. Ans. $2\frac{1}{30}$.
 8. Required the product of $\frac{5}{6}$, and $\frac{2}{3}$ of $\frac{4}{7}$. Ans. $\frac{10}{21}$.
 9. Required the product of 6, and $\frac{2}{3}$ of 5. Ans. 20.
 10. Required the product of $\frac{2}{9}$ of $\frac{3}{5}$, and $\frac{5}{6}$ of $3\frac{2}{7}$. Ans. $\frac{23}{84}$.
 11. Required the product of $3\frac{2}{7}$ and $4\frac{1}{3}$. Ans. $14\frac{11}{21}$.
 12. Required the product of 5, $\frac{2}{3}$, $\frac{2}{7}$ of $\frac{3}{5}$, and $4\frac{1}{6}$. Ans. $2\frac{2}{21}$.

DIVISION OF VULGAR FRACTIONS.

* PREPARE the fractions as before in multiplication; then divide the numerator by the numerator, and the denominator by the denominator, if they will exactly divide: but if not, then invert the terms of the divisor, and multiply the dividend by it, as in multiplication.

EXAMPLES.

1. Divide $2\frac{5}{9}$ by $\frac{5}{3}$.
 Here $2\frac{5}{9} \div \frac{5}{3} = \frac{5}{3} = 1\frac{2}{3}$, by the first method.
2. Divide $\frac{2}{6}$ by $\frac{2}{15}$.
 Here $\frac{5}{9} \div \frac{2}{15} = \frac{5}{9} \times \frac{15}{2} = \frac{5}{3} \times \frac{5}{2} = 2\frac{5}{6} = 4\frac{1}{6}$.
3. It is required to divide $\frac{1}{2}$ by $\frac{4}{5}$. Ans. $\frac{5}{8}$.
4. It is required to divide $\frac{7}{16}$ by $\frac{3}{4}$. Ans. $\frac{7}{12}$.
5. It is required to divide $\frac{1}{6}$ by $\frac{7}{8}$. Ans. $1\frac{1}{3}$.
6. It is required to divide $\frac{5}{6}$ by $\frac{1}{7}$. Ans. $7\frac{5}{6}$.
7. It is required to divide $\frac{1}{3}$ by $\frac{2}{5}$. Ans. $\frac{5}{6}$.
8. It is required to divide $\frac{2}{7}$ by $\frac{3}{4}$. Ans. $\frac{8}{21}$.

* Division being the reverse of Multiplication, the reason of the Rule is evident.

Note. A fraction is best divided by an integer, by dividing the numerator by it; but if it will not exactly divide, then multiply the denominator by it.

9. It

RULE OF THREE IN VULGAR FRACTIONS. 65

9. It is required to divide $\frac{9}{16}$ by 3. Ans. $\frac{3}{16}$.
 10. It is required to divide $\frac{3}{5}$ by 2. Ans. $\frac{3}{10}$.
 11. It is required to divide $7\frac{1}{3}$ by $9\frac{5}{8}$. Ans. $\frac{32}{43}$.
 12. It is required to divide $\frac{1}{3}$ of $\frac{1}{2}$ by $\frac{2}{7}$ of $7\frac{2}{5}$. Ans. $\frac{7}{171}$.

RULE OF THREE IN VULGAR FRACTIONS.

MAKE the necessary preparations as before directed ; then multiply continually together, the second and the third terms, and the first with its parts inverted as in Division, for the answer.*

EXAMPLES.

1. If $\frac{2}{3}$ of a yard of velvet cost $\frac{2}{5}$ of a pound sterling ; what will $\frac{5}{16}$ of a yard cost ?

$$\frac{3}{8} : \frac{2}{5} : : \frac{5}{16} : \frac{8}{3} \times \frac{2}{5} \times \frac{3}{8} = \frac{1}{3}l = 6s \ 8d, \text{ Answer.}$$

$$\frac{8}{3} \times \frac{5}{2} \times \frac{16}{5} = 16$$

2. What will $3\frac{3}{8}$ oz of silver cost, at 6s 4d an ounce ?

Ans. 11 1s $4\frac{1}{2}d$.

3. If $\frac{3}{8}$ of a ship be worth 273l 2s 6d ; what are $\frac{6}{32}$ of her worth ?

Ans. 227l 12s 1d.

4. What is the purchase of 1230l bank-stock, at $108\frac{5}{8}$ per cent. ?

Ans. 1336l 1s 9d.

5. What is the interest of 273l 15s for a year at $3\frac{1}{4}$ per cent. ?

Ans. 8l 17s $1\frac{1}{4}d$.

6. If $\frac{1}{8}$ of a ship be worth 73l 1s 3d ; what part of her is worth 250l 10s ?

Ans. $\frac{3}{7}$.

7. What length must be cut off a board that is $7\frac{3}{4}$ inches broad, to contain a square foot, or as much as another piece of 12 inches long and 12 broad ?

Ans. $18\frac{1}{3}\frac{1}{4}$ inches.

8. What quantity of shalloon that is $\frac{3}{4}$ of a yard wide, will line $9\frac{1}{2}$ yards of cloth, that is $2\frac{1}{2}$ yards wide ?

Ans. $31\frac{2}{3}$ yds.

* This is only multiplying the 2nd and 3rd terms together, and dividing the product by the first, as in the Rule of Three in whole numbers.

9. If the penny loaf weighs $6\frac{9}{10}$ oz, when the price of wheat is 5s the bushel; what ought it to weigh when the wheat is 8s 6d the bushel? Ans. $4\frac{1}{7}$ oz.

10. How much in length, of a piece of land that is $11\frac{1}{2}$ poles broad, will make an acre of land, or as much as 40 poles in length and 4 in breadth? Ans. $13\frac{6}{14}\frac{1}{2}$ poles.

11. If a courier perform a certain journey in $35\frac{1}{2}$ days, travelling $13\frac{5}{8}$ hours a day; how long would he be in performing the same, travelling only $11\frac{9}{10}$ hours a day? Ans. $40\frac{8}{9}\frac{1}{5}\frac{1}{2}$ days.

12. A regiment of soldiers, consisting of 976 men, are to be new clothed; each coat to contain $2\frac{1}{2}$ yards of cloth that is $1\frac{5}{8}$ yard wide, and lined with shalloon $\frac{7}{8}$ yard wide: how many yards of shalloon will line them? Ans. 4531 yds 1 qr $2\frac{6}{7}$ nails.

DECIMAL FRACTIONS.

A DECIMAL FRACTION, is that which has for its denominator an unit (1), with as many ciphers annexed as the numerator has places; and it is usually expressed by setting down the numerator only, with a point before it, on the left hand. Thus, $\frac{4}{10}$ is $\cdot 4$, and $\frac{24}{100}$ is $\cdot 24$, and $\frac{74}{1000}$ is $\cdot 074$, and $\frac{124}{10000}$ is $\cdot 00124$; where ciphers are prefixed to make up as many places as are ciphers in the denominator, when there is a deficiency of figures.

A mixed number is made up of a whole number with some decimal fraction, the one being separated from the other by a point. Thus, $3\cdot 25$ is the same as $3\frac{25}{100}$, or $3\frac{5}{20}$.

Ciphers on the right hand of decimals make no alteration in their value; for $\cdot 4$ or 40 , or 400 , are decimals having all the same value, each being $= \frac{4}{10}$ or $\frac{2}{5}$. But when they are placed on the left hand they decrease the value in a ten-fold proportion: Thus, $\cdot 4$ is $\frac{4}{10}$, or 4 tenths: but $\cdot 04$ is only $\frac{4}{100}$, or 4 hundredths, and $\cdot 004$ is only $\frac{4}{1000}$, or 4 thousandths.

The 1st place of decimals, counted from the left hand towards the right, is called the place of primes, or 10ths; the 2d is the place of seconds, or 100ths; the 3d is the place of thirds, or 1000ths; and so on. For in decimals, as well as in whole numbers, the values of the places increase towards the left hand, and decrease towards the right, both in the same

same ten-fold proportion ; as in the following Scale or Table of Notation.

3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
millions	hundred thousands	ten thousands	thousands	hundreds	tens	units	tenth parts	hundredth parts	thousandth parts	ten thousandth parts	hundred thousandth parts	millionth parts		

—◆—

ADDITION OF DECIMALS.

SET the numbers under each other according to the value of their places, like as in whole numbers ; in which state the decimal separating points will stand all exactly under each other. Then, beginning at the right-hand, add up all the columns of numbers as in integers ; and point off as many places, for decimals, as are in the greatest number of decimal places in any of the lines that are added ; or place the point directly below all the other points.

EXAMPLES.

1. To add together 29·0146, and 3146·5, and 2109, and ·62417, and 14·16.

$$\begin{array}{r}
 29\cdot0146 \\
 3146\cdot5 \\
 2109\cdot \\
 \quad\cdot62417 \\
 14\cdot16 \\
 \hline
 5299\cdot29877 \text{ the Sum.} \\
 \hline
 \end{array}$$

Ex. 2. What is the sum of 276, 39·213, 72014·9, 417, and 5032 ?

3. What is the sum of 7530, 16·201, 3·0142, 957·13, 6·72119 and ·03014.

4. What is the sum of 312·09, 3·5711, 7195·6, 71·498, 9739·215, 179, and ·0027 ?

SUBTRACTION

SUBTRACTION OF DECIMALS.

PLACE the numbers under each other according to the value of their places, as in the last Rule. Then, beginning at the right hand, subtract as in whole numbers, and point off the decimals as in Addition.

EXAMPLES.

1. To find the difference between 91.73 and 2.138.

$$\begin{array}{r} 91.73 \\ 2.138 \\ \hline \end{array}$$

Ans. 89.592

2. Find the diff. between 1.9185 and 2.73 Ans. 0.8115.
 3. To subtract 4.90142 from 214.81. Ans. 209.90858.
 4. Find the diff. between 2714 and .916. Ans. 2713.084.

MULTIPLICATION OF DECIMALS.

* PLACE the factors, and multiply them together the same as if they were whole numbers.—Then point off in the product just as many places of decimals as there are decimals in both the factors. But if there be not so many figures in the product, then supply the defect by prefixing ciphers.

* The Rule will be evident from this example:—Let it be required to multiply .12 by .361; these numbers are equivalent to $\frac{12}{100}$ and $\frac{361}{1000}$; the product of which is $\frac{4332}{100000} = .04332$, by the nature of Notation, which consists of as many places as there are ciphers, that is, of as many places as there are in both numbers. And in like manner for any other numbers.

EXAMPLES.

EXAMPLES.

1. Multiply $\cdot 321096$
by $\cdot 2465$

1605480
1926576
1284384
642192

Ans. $\cdot 0791501640$ the Product.

2. Multiply $79\cdot 347$ by $23\cdot 15$. Ans. $1836\cdot 88305$.
3. Multiply $\cdot 63478$ by 8204 . Ans. $\cdot 520773512$.
4. Multiply $\cdot 385746$ by $\cdot 00464$. Ans. $\cdot 00178986144$.

CONTRACTION I.

To multiply Decimals by 1 with any number of Ciphers, as by 10, or 100, or 1000, &c.

THIS is done by only removing the decimal point so many places farther to the right hand, as there are ciphers in the multiplier ; and subjoining ciphers if need be.

EXAMPLES.

1. The product of $51\cdot 3$ and 1000 is 51300 .
2. The product of $2\cdot 714$ and 100 is
3. The product of $\cdot 916$ and 1000 is
4. The product of $21\cdot 31$ and 10000 is

CONTRACTION II.

To Contract the Operation, so as to retain only as many Decimals in the Product as may be thought Necessary, when the Product would naturally contain several more Places.

SET the units' place of the multiplier under that figure of the multiplicand whose place is the same as is to be retained for the last in the product ; and dispose of the rest of the figures in the inverted or contrary order to what they are usually placed in.—Then, in multiplying, reject all the figures that are more to the right hand than each multiplying figure, and set down the products, so that their right hand figures may

may fall in a column straight below each other; but observing to increase the first figure of every line with what would arise from the figures omitted, in this manner, namely 1 from 5 to 14, 2 from 15 to 24, 3 from 25 to 34, &c.; and the sum of all the lines will be the product as required, commonly to the nearest unit in the last figure.

EXAMPLES.

1. To multiply 27·14986 by 92·41035, so as to retain only four places of decimals in the product.

Contracted Way.

$$\begin{array}{r}
 27\cdot14986 \\
 53014\cdot29 \\
 \hline
 24434874 \\
 542997 \\
 108599 \\
 2715 \\
 81 \\
 14 \\
 \hline
 2508\cdot9280 \\
 \hline
 \end{array}$$

Common Way.

$$\begin{array}{r}
 27\cdot14986 \\
 92\cdot41035 \\
 \hline
 13574930 \\
 8144958 \\
 2714986 \\
 10859944 \\
 5429972 \\
 24434874 \\
 \hline
 2508\cdot9280 \quad 650510 \\
 \hline
 \end{array}$$

2. Multiply 480·14936 by 2·72416, retaining only four decimals in the product.

3. Multiply 2490·3048 by ·573286, retaining only five decimals in the product.

4. Multiply 325·701428 by ·7218393, retaining only three decimals in the product.

DIVISION OF DECIMALS.

DIVIDE as in whole numbers; and point off in the quotient as many places for decimals, as the decimal places in the dividend exceed those in the divisor.*

* The reason of this Rule is evident; for, since the divisor multiplied by the quotient gives the dividend, therefore the number of decimal places in the dividend, is equal to those in the divisor and quotient, taken together, by the nature of Multiplication; and consequently the quotient itself must contain as many as the dividend exceeds the divisor.

Another

Another way to know the place for the decimal point, is this : The first figure of the quotient must be made to occupy the same place, of integers or decimals, as doth that figure of the dividend which stands over the unit's figure of the first product.

When the places of the quotient are not so many as the Rule requires, the defect is to be supplied by prefixing ciphers.

When there happens to be a remainder after the division, or when the decimal places in the divisor are more than those in the dividend ; then ciphers may be annexed to the dividend, and the quotient carried on as far as required.

EXAMPLES.

<p style="text-align: center;">1.</p> $ \begin{array}{r} 178) \cdot 48520928 \text{ (} \cdot 00272589 \\ \underline{1292} \\ 460 \\ \underline{1049} \\ 1599 \\ \underline{1758} \\ 156 \\ \hline \end{array} $		<p style="text-align: center;">2.</p> $ \begin{array}{r} \cdot 2639) 27\cdot 00000 \text{ (} 102\cdot 3114 \\ \underline{6100} \\ 8220 \\ \underline{3030} \\ 3910 \\ \underline{12710} \\ 2154 \\ \hline \end{array} $
--	--	--

- | | |
|-------------------------------|---------------|
| 3. Divide 123·70536 by 54·25. | Ans. 2·2802. |
| 4. Divide 12 by 7854. | Ans. 15·278. |
| 5. Divide 4195·68 by 100. | Ans. 41·9568. |
| 6. Divide ·8297592 by ·153. | Ans. 5·4232. |

CONTRACTION I.

WHEN the divisor is an integer, with any number of ciphers annexed : cut off those ciphers, and remove the decimal point in the dividend as many places farther to the left as there are ciphers cut off, prefixing ciphers if need be ; then proceed as before.*

* This is no more than dividing both divisor and dividend by the same number, either 10, or 100, or 1000, &c. according to the number of ciphers cut off, which, leaving them in the same proportion, does not affect the quotient. And, in the same way, the decimal point may be moved the same number of places in both the divisor and dividend, either to the right or left, whether they have ciphers or not.

EXAMPLES.

EXAMPLES.

1. Divide 45.5 by 2100.

$$\begin{array}{r} 21 \cdot 00 \) \cdot 455 \ (\cdot 0216, \ \&c. \\ \underline{35} \\ 140 \\ \underline{14} \\ \hline \end{array}$$

2. Divide 41020 by 32000.
 3. Divide 953 by 21600.
 4. Divide 61 by 79000.

CONTRACTION II.

Hence, if the divisor be 1 with ciphers, as 10, 100, or 1000, &c. : then the quotient will be found by merely moving the decimal point in the dividend, so many places farther to the left, as the divisor has ciphers ; prefixing ciphers if need be.

EXAMPLES.

$$\begin{array}{ll} \text{So, } 217.3 \div 100 = 2.173 & \text{And } 419 \div 10 = \\ \text{And } 5.16 \div 100 = & \text{And } 21 \div 1000 = \end{array}$$

CONTRACTION III.

WHEN there are many figures in the divisor ; or when only a certain number of decimals are necessary to be retained in the quotient ; then take only as many figures of the divisor as will be equal to the number of figures, both integers and decimals, to be in the quotient, and find how many times they may be contained in the first figures of the dividend, as usual.

Let each remainder be a new dividend ; and for every such dividend, leave out one figure more on the right-hand side of the divisor ; remembering to carry for the increase of the figures cut off, as in the 2d contraction in Multiplication.

Note. When there are not so many figures in the divisor, as are required to be in the quotient, begin the operation with all the figures, and continue it as usual till the number of figures in the divisor be equal to those remaining to be found in the quotient : after which begin the contraction.

EXAMPLES.

1. Divide 2508.92806 by 92.41035, so as to have only four decimals in the quotient, in which case the quotient will contain six figures.

Contracted

<i>Contracted.</i>	<i>Common.</i>
92·4103,5)2508·928,06(27·1498	92·4103,5)2508·928,06(27·1498
660721	66072106
13849	13848610
4608	46075750
912	91116100
80	79467850
6	5539570

2. Divide 4109·2351 by 230·409, so that the quotient may contain only four decimals. Ans. 17·8345.

3. Divide 37·10438 by 5713·96, that the quotient may contain only five decimals. Ans. 00649,

4. Divide 913·08 by 2137·2, that the quotient may contain only three decimals.

—◆—

REDUCTION OF DECIMALS.

CASE I.

To reduce a Vulgar Fraction to its equivalent Decimal.

DIVIDE the numerator by the denominator as in Division of Decimals, annexing ciphers to the numerator as far as necessary ; so shall the quotient be the decimal required.

EXAMPLES.

1. Reduce $\frac{7}{24}$ to a decimal.
 $24 = 4 \times 6$. Then $4 \overline{) 7}$
 $6 \overline{) 1.750000}$
·291666 &c.

2. Reduce $\frac{1}{4}$, and $\frac{1}{2}$, and $\frac{3}{4}$, to decimals. Ans. ·25, and ·5, and ·75.

3. Reduce $\frac{5}{8}$ to a decimal. Ans. ·625.

4. Reduce $\frac{3}{25}$ to a decimal. Ans. ·12.

5. Reduce $\frac{6}{1002}$ to a decimal. Ans. 031350.

6. Reduce $\frac{550}{3842}$ to a decimal. Ans. ·143155 &c.

CASE II.

To find the Value of a Decimal in terms of the Inferior Denominations.

MULTIPLY the decimal by the number of parts in the next lower denomination ; and cut off as many places for a remainder to the right hand, as there are places in the given decimal.

Multiply that remainder by the parts in the next lower denomination again, cutting off for another remainder as before.

Proceed in the same manner through all the parts of the integer ; then the several denominations separated on the left hand, will make up the answer.

Note, This operation is the same as Reduction Descending in whole numbers.

EXAMPLES.

1. Required to find the value of $\cdot 775$ pounds sterling.

$$\begin{array}{r}
 \cdot 775 \\
 20 \\
 \hline
 s. 15.500 \\
 12 \\
 \hline
 d. 6.000
 \end{array}
 \qquad
 \text{Ans. } 15s \ 6d.$$

2. What is the value of $\cdot 625$ shil ? Ans. $7\frac{1}{2}d.$
 3. What is the value of $\cdot 8635l$? Ans. $17s. 3.24d.$
 4. What is the value of $\cdot 0125$ lb troy ? Ans. 3 dwts.
 5. What is the value of $\cdot 4694$ lb troy ?
Ans. 5 oz 12 dwts 15.744 gr.
 6. What is the value of $\cdot 625$ cwt ? Ans. 2 qr 14 lb.
 7. What is the value of $\cdot 009943$ miles ?
Ans. 17 yd 1 ft 5.98848 inc.
 8. What is the value of $\cdot 6875$ yd ? Ans. 2 qr 3 nls.
 9. What is the value of $\cdot 3375$ acr ? Ans. 1 rd 14 poles.
 10. What is the value of $\cdot 2083$ hhd of wine ?
Ans. 13.1229 gal.

CASE III.

To reduce Integers or Decimals to Equivalent Decimals of Higher Denominations.

DIVIDE by the number of parts in the next higher denomination; continuing the operation to as many higher denominations as may be necessary, the same as in Reduction Ascending of whole numbers.

EXAMPLES.

1. Reduce 1 dwt to the decimal of a pound troy.

20	1 dwt
12	0.05 oz
	0.004166 &c. lb. Ans.
2. Reduce 9d to the decimal of a pound. Ans. .0375*l*.
3. Reduce 7 drams to the decimal of a pound avoird.
Ans. .02734375*lb*.
4. Reduce 26d to the decimal of a *l*. Ans. .0010833 &c. *l*.
5. Reduce 2.15 lb to the decimal of cwt.
Ans. .019196 † cwt.
6. Reduce 24 yards to the decimal of a mile.
Ans. .013636 &c. mile.
7. Reduce .056 pole to the decimal of an acre.
Ans. .00035 ac.
8. Reduce 1.2 pint of wine to the decimal of a hhd.
Ans. .00238 † hhd.
9. Reduce 14 minutes to the decimal of a day.
Ans. .009722 &c. da.
10. Reduce .21 pint to the decimal of a peck.
Ans. .013125 pec.
11. Reduce 28" 12'" to the decimal of a minute.

NOTE, *When there are several numbers, to be reduced all to the decimal of the highest:*

Set the given numbers directly under each other, for dividends, proceeding orderly from the lowest denomination to the highest.

Opposite to each dividend, on the left hand, set such a number for a divisor as will bring it to the next higher name; drawing a perpendicular line between all the divisors and dividends.

Begin at the uppermost, and perform all the divisions: only observing to set the quotient of each division, as decimal parts,

parts, on the right hand of the dividend next below it; so shall the last quotient be the decimal required.

EXAMPLES.

1. Reduce $17s\ 9\frac{3}{4}d$ to the decimal of a pound.

$$\begin{array}{r|l} 4 & 3 \\ 12 & 9.75 \\ 20 & 17.8125 \\ \hline & \text{£ } 0.890625 \text{ Ans.} \end{array}$$

2. Reduce $19l\ 17s\ 3\frac{1}{4}d$ to l . Ans. 19.86354166 &c. l .
 3. Reduce $15s\ 6d$ to the decimal of a l . Ans. $.775l$.
 4. Reduce $7\frac{1}{2}d$ to the decimal of a shilling. Ans. $.625s$.
 5. Reduce $5\text{ oz } 12\text{ dwts } 16\text{ gr}$ to lb . Ans. $.46944$ &c. lb .



RULE OF THREE IN DECIMALS.

PREPARE the terms by reducing the vulgar fractions to decimals, and any compound numbers either to decimals of the higher denominations, or to integers of the lower, also the first and third terms to the same name: Then multiply and divide as in whole numbers.

Note. Any of the convenient Examples in the Rule of Three or Rule of Five in integers, or Vulgar Fractions, may be taken as proper examples to the same rules in Decimals.—The following Example, which is the first in Vulgar Fractions, is wrought out here, to show the method.

If $\frac{3}{8}$ of a yard of velvet cost $\frac{2}{5}l$, what will $\frac{5}{16}$ yd cost?

$\frac{3}{8} = .375$	yd l	yd l	s d
	$.375 : .4 ::$	$.3125 : .333$	&c. or 6 8
		$.4$	
$\frac{2}{5} = .4$.375	.12500
		1250	20
		125	6.66666 &c.
$\frac{5}{16} = .3125$			12
	Ans. 6s 8d.		d 7.99999 &c. = 8d.

DUODECIMALS.

DUODECIMALS OF CROSS MULTIPLICATION, is a rule used by workmen and artificers, in computing the contents of their works.

Dimensions are usually taken in feet, inches, and quarters; any parts smaller than these being neglected as of no consequence. And the same in multiplying them together, or casting up the contents. The method is as follows.

SET down the two dimensions to be multiplied together, one under the other, so that feet may stand under feet, inches under inches, &c.

Multiply each term in the multiplicand, beginning at the lowest, by the feet in the multiplier, and set the result of each straight under its corresponding term, observing to carry 1 for every 12, from the inches to the feet.

In like manner, multiply all the multiplicand by the inches and parts of the multiplier, and set the result of each term one place removed to the right hand of those in the multiplicand; omitting, however, what is below parts of inches, only carrying to these the proper number of units from the lowest denomination.

Or, instead of multiplying by the inches, take such parts of the multiplicand as there are of a foot.

Then add the two lines together after the manner of Compound Addition, carrying 1 to the feet for 12 inches, when these come to so many.

EXAMPLES.

$$\begin{array}{r} 1. \text{ Multiply } 4 \text{ f } 7 \text{ inc.} \\ \text{by } 6 \quad 4 \\ \hline \end{array}$$

$$\begin{array}{r} 27 \quad 6 \\ 1 \quad 6\frac{1}{3} \\ \hline \end{array}$$

$$\text{Ans. } 29 \quad 0\frac{1}{3}$$

$$\begin{array}{r} 2. \text{ Multiply } 14 \text{ f } 9 \text{ inc.} \\ \text{by } 4 \quad 6 \\ \hline \end{array}$$

$$\begin{array}{r} 59 \quad 0 \\ 7 \quad 4\frac{1}{2} \\ \hline \end{array}$$

$$\text{Ans. } 66 \quad 4\frac{1}{2}$$

- | | |
|--|-------------------------------|
| 3. Multiply 4 feet 7 inches by 9 f 6 inc. | Ans. 43 f 6 $\frac{1}{2}$ inc |
| 4. Multiply 12 f 5 inc by 6 f 8 inc. | Ans. 32 9 $\frac{1}{3}$ |
| 5. Multiply 35 f 4 $\frac{1}{2}$ inc by 12 f 3 inc | Ans. 433 4 $\frac{1}{8}$ |
| 6. Multiply 64 f 6 inc by 8 f 9 $\frac{1}{4}$ inc. | Ans. 565 8 $\frac{3}{4}$ |

INVOLUTION.

INVOLUTION.

INVOLUTION is the raising of Powers from any given number, as a root.

A Power is a quantity produced by multiplying any given number, called the Root, a certain number of times continually by itself. Thus,

$$\begin{aligned} 2 &= 2 \text{ is the root, or 1st power of } 2. \\ 2 \times 2 &= 4 \text{ is the 2d power, or square of } 2. \\ 2 \times 2 \times 2 &= 8 \text{ is the 3d power, or cube of } 2. \\ 2 \times 2 \times 2 \times 2 &= 16 \text{ is the 4th power of } 2, \text{ \&c.} \end{aligned}$$

And in this manner may be calculated the following Table of the first nine powers of the first 9 numbers.

TABLE OF THE FIRST NINE POWERS OF NUMBERS.

1st	2d	3d	4th	5th	6th	7th	8th	9th
1	1	1	1	1	1	1	1	1
2	4	8	16	32	64	128	256	512
3	9	27	81	243	729	2187	6561	19683
4	16	64	256	1024	4096	16384	65536	262144
5	25	125	625	3125	15625	78125	390625	1953125
6	36	216	1296	7776	46656	279936	1679616	10077696
7	49	343	2401	16807	117649	823543	5764801	40353607
8	64	512	4096	32768	262144	2097152	16777216	134217728
9	81	729	6561	59049	531441	4782969	43046721	387420489

The index or Exponent of a Power, is the number denoting the height or degree of that power; and it is 1 more than the number of multiplications used in producing the same. So 1 is the index or exponent of the first power or root, two of the 2d power or square, 3 of the third power or cube, 4 of the 4th power, and so on.

Powers, that are to be raised, are usually denoted by placing the index above the root or first power.

So $2^2=4$ is the 2d power of 2.
 $2^3=8$ is the 3d power of 2.
 $2^4=16$ is the 4th power of 2.
 540^4 is the 4th power of 540, &c.

When two or more powers are multiplied together, their product is that power whose index is the sum of the exponents of the factors or powers multiplied. Or the multiplication of the powers, answers to the addition of the indices. Thus, in the following powers of 2,

	1st	2d	3d	4th	5th	6th	7th	8th	9th	10th
	2	4	8	16	32	64	128	256	512	1024
or	2^1	2^2	2^3	2^4	2^5	2^6	2^7	2^8	2^9	2^{10}

Here, $4 \times 4 = 16$, and $2 + 2 = 4$ its index;
 and $8 \times 16 = 128$, and $3 + 4 = 7$ its index;
 also $16 \times 64 = 1024$, and $4 + 6 = 10$ its index;

OTHER EXAMPLES.

- | | |
|---|--------------------------|
| 1. What is the 2d power of 45? | Ans. 2025. |
| 2. What is the square of 4.16? | Ans. 17.3056. |
| 3. What is the 3d power of 3.5 | Ans. 42.875. |
| 4. What is the 5th power of .029? | Ans. .00000020511149. |
| 5. What is the square of $\frac{2}{3}$? | Ans. $\frac{4}{9}$. |
| 6. What is the 3d power of $\frac{5}{6}$? | Ans. $\frac{125}{216}$. |
| 7. What is the 4th power of $\frac{3}{4}$? | Ans. $\frac{81}{256}$. |

EVOLUTION.

EVOLUTION, or the reverse of Involution, is the extracting or finding the roots of any given powers.

The root of any number, or power, is such a number, as being multiplied into itself a certain number of times, will produce that power. Thus, 2 is the square root or 2d root of 4, because $2^2 = 2 \times 2 = 4$; and 3 is the cube root or 3d root of 27, because $3^3 = 3 \times 3 \times 3 = 27$.

Any power of a given number or root may be found exactly, namely, by multiplying the number continually into itself. But there are many numbers of which a proposed root can never be exactly found. Yet by means of decimals, we may approximate or approach towards the root, to any degree of exactness.

Those roots which only approximate, are called Surd roots; but those which can be found quite exact, are called Rational Roots. Thus, the square root of 3 is a surd root; but the square root of 4 is a rational root, being equal to 2: also the cube root of 8 is rational, being equal to 2; but the cube root of 9 is surd or irrational.

Roots are sometimes denoted by writing the character $\sqrt{\quad}$ before the power, with the index of the root against it. Thus, the third root of 20 is expressed $\sqrt[3]{20}$; and the square root or 2d root of it is $\sqrt{20}$, the index 2 being always omitted, when only the square root is designed.

When the power is expressed by several numbers, with the sign $+$ or $-$ between them, a line is drawn from the top of the sign over all the parts of it: thus the third root of $45 - 12$ is $\sqrt[3]{45 - 12}$, or thus $\sqrt[3]{(45 - 12)}$, inclosing the numbers in parenthesis.

But all roots are now often designed like powers, with fractional indices: thus, the square root of 8 is $8^{\frac{1}{2}}$, the cube root of 25 is $25^{\frac{1}{3}}$, and the fourth root of $45 - 18$ is $(45 - 18)^{\frac{1}{4}}$, or $(45 - 18)^{\frac{1}{4}}$.

TO EXTRACT THE SQUARE ROOT.

* **DIVIDE** the given number into periods of two figures each, by setting a point over the place of units, another over the place of hundreds, and so on, over every second figure, both to the left-hand in integers, and to the right in decimals.

Find the greatest square in the first period on the left-hand, and set its root on the right-hand of the given number, after the manner of a quotient figure in Division.

* The reason for separating the figures of the dividend into periods or portions of two places each, is, that the square of any single figure never consists of more than two places; the square of a number of two figures, of not more than four places, and so on. So that there will be as many figures in the root as the given number contains periods so divided or parted off.

And the reason of the several steps in the operation appears from the algebraic form of the square of any number of terms, whether two or three or more. Thus,

$(a + b)^2 = a^2 + 2ab + b^2 = a^2 + (2a + b)b$, the square of two terms; where it appears that a is the first term of the root, and b the second term; also a the first divisor, and the new divisor is $2a + b$, or double the first term increased by the second. And hence the manner of extraction is thus:

$$\text{1st divisor } a \begin{array}{r} a^2 + 2ab + b^2 \\ \underline{a^2} \end{array} (a + b \text{ the root.})$$

$$\text{2d divisor } \begin{array}{r} 2a + b \\ b \end{array} \begin{array}{r} 2ab + b^2 \\ \underline{2ab + b^2} \end{array}$$

Again, for a root of three parts, a, b, c , thus:

$$(a + b + c)^2 = a^2 + 2ab + b^2 + 2ac + 2bc + c^2 =$$

$a^2 + (2a + b)b + (2a + 2b + c)c$, the square of three terms, where a is the first term of the root b , the second, and c the third term; also a the first divisor, $2a + b$ the second, and $2a + 2b + c$ the third, each consisting of the double of the root increased by the next term of the same. And the mode of extraction is thus:

$$\text{1st divisor } a \begin{array}{r} a^2 + 2ab + b^2 + 2ac + 2bc + c^2 \\ \underline{a^2} \end{array} (a + b + c \text{ the root.})$$

$$\text{2d divisor } \begin{array}{r} 2a + b \\ b \end{array} \begin{array}{r} 2ab + b^2 \\ \underline{2ab + b^2} \end{array}$$

$$\text{3d divisor } \begin{array}{r} 2a + 2b + c \\ c \end{array} \begin{array}{r} 2ac + 2bc + c^2 \\ \underline{2ac + 2bc + c^2} \end{array}$$

Subtract the square thus found from the said period, and to the remainder annex the two figures of the next following period, for a dividend.

Double the root above mentioned for a divisor ; and find how often it is contained in the said dividend, exclusive of its right-hand figure ; and set that quotient figure both in the quotient and divisor.

Multiply the whole augmented divisor by this last quotient figure, and subtract the product from the said dividend, bringing down to it the next period of the given number, for a new dividend.

Repeat the same process over again, viz. find another new divisor, by doubling all the figures now found in the root ; from which, and the last dividend, find the next figure of the root as before ; and so on through all the periods, to the last.

Note, The best way of doubling the root, to form the new divisors, is by adding the last figure always to the last divisor, as appears in the following examples.—Also, after the figures belonging to the given number are all exhausted, the operation may be continued into decimals at pleasure, by adding any number of periods of ciphers, two in each period.

EXAMPLES.

1. To find the square root of 29506624.

$$\begin{array}{r}
 29506624(5432 \text{ the root.} \\
 25 \\
 \hline
 104 \mid 450 \\
 4 \mid 416 \\
 \hline
 1083 \mid 3466 \\
 3 \mid 3249 \\
 \hline
 10862 \mid 21724 \\
 2 \mid 21724 \\
 \hline
 \end{array}$$

NOTE, When the root is to be extracted to many places of figures, the work may be considerably shortened, thus :

Having proceeded in the extraction after the common method, till there be found half the required number of figures
in

in the root; or one figure more ; then, for the rest, divide the last remainder by its corresponding divisor, after the manner of the third contraction in Division of Decimals ; thus,

2. To find the root of 2 to nine places of figures.

2 (1.41421356 the root.

1

$$\begin{array}{r|l} 24 & 100 \\ 4 & 96 \end{array}$$

$$\begin{array}{r|l} 281 & 400 \\ 1 & 281 \end{array}$$

$$\begin{array}{r|l} 2824 & 11900 \\ 4 & 11296 \end{array}$$

$$\begin{array}{r|l} 28282 & 60400 \\ 2 & 56564 \end{array}$$

$$\begin{array}{r} 28284) 3836 (1356 \\ \dots \quad 1008 \\ \quad \quad 160 \\ \quad \quad \quad 19 \\ \quad \quad \quad \quad 2 \end{array}$$

- | | |
|---|----------------|
| 3. What is the square root of 2025 ? | Ans. 45. |
| 4. What is the square root of 17.3056 ? | Ans. 4.16. |
| 5. What is the square root of .000729 ? | Ans. .027. |
| 6. What is the square root of 3 ? | Ans. 1.732050. |
| 7. What is the square root of 5 ? | Ans. 2.236068. |
| 8. What is the square root of 6 ? | Ans. 2.449489. |
| 9. What is the square root of 7 ? | Ans. 2.645751. |
| 10. What is the square root of 10 ? | Ans. 3.162277. |
| 11. What is the square root of 11 ? | Ans. 3.316624. |
| 12. What is the square root of 12 ? | Ans. 3.464101. |

RULES FOR THE SQUARE ROOTS OF VULGAR FRACTIONS AND MIXED NUMBERS.

First prepare all vulgar fractions, by reducing them to their least terms, both for this and all other roots. Then

1. Take the root of the numerator and of the denominator for the respective terms of the root required. And this is the best way if the denominator be a complete power : but if it be not, then

2. Multiply the numerator and denominator together ; take the root of the product : this root being made the numerator

rator to the denominator of the given fraction, or made the denominator to the numerator of it, will form the fractional root required.

$$\text{That is, } \sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}} = \frac{\sqrt{ab}}{b} = \frac{a}{\sqrt{ab}}.$$

And this rule will serve, whether the root be finite or infinite.

3. Or reduce the vulgar fraction to a decimal, and extract its root.

4. Mixed numbers may be either reduced to improper fractions, and extracted by the first or second rule, or the vulgar fraction may be reduced to a decimal, then joined to the integer, and the root of the whole extracted.

EXAMPLES.

- | | |
|---|----------------------|
| 1. What is the root of $\frac{25}{36}$? | Ans. $\frac{5}{6}$. |
| 2. What is the root of $\frac{27}{147}$? | Ans. $\frac{3}{7}$. |
| 3. What is the root of $\frac{9}{12}$? | Ans. 0·866025. |
| 4. What is the root of $\frac{5}{12}$? | Ans. 0·645497. |
| 5. What is the root of $17\frac{3}{8}$? | Ans. 4·168333. |

By means of the square root also may readily be found the 4th root, or the 8th root, or the 16th root, &c. that is, the root of any power whose index is some power of the number 2; namely, by extracting so often the square root as is denoted by that power of 2; that is, two extractions for the 4th root, three for the 8th root, and so on.

So, to find the 4th root of the number 21035·8, extract the square root two times as follows:

$\begin{array}{r} \overset{\cdot}{2}\overset{\cdot}{1}\overset{\cdot}{0}\overset{\cdot}{3}\overset{\cdot}{5}\overset{\cdot}{8}\overset{\cdot}{0}\overset{\cdot}{0}\overset{\cdot}{0} \\ \underline{1} \end{array}$	$\begin{array}{r} \overset{\cdot}{1}\overset{\cdot}{4}\overset{\cdot}{5}\overset{\cdot}{0}\overset{\cdot}{3}\overset{\cdot}{7}\overset{\cdot}{2}\overset{\cdot}{3}\overset{\cdot}{7} \\ \underline{1} \end{array}$	(12·0431407 the 4th root.)
$\begin{array}{r} 24 \overline{) 110} \\ \underline{4 \quad 96} \end{array}$	$\begin{array}{r} 22 \overline{) 45} \\ \underline{2 \quad 44} \end{array}$	
$\begin{array}{r} 285 \overline{) 1435} \\ \underline{5 \quad 1425} \end{array}$	$\begin{array}{r} 2404 \overline{) 10372} \\ \underline{4 \quad 9616} \end{array}$	
$\begin{array}{r} 29003 \overline{) 108000} \\ \underline{3 \quad 87009} \end{array}$	$\begin{array}{r} 24083 \overline{) 75637} \\ \underline{3 \quad 72249} \end{array}$	
$\begin{array}{r} 20991 \\ \underline{687} \\ 107 \end{array}$	$\begin{array}{r} 7237 \\ \underline{3388} \\ 960 \\ \underline{17} \end{array}$	(1407

Ex. 2. What is the 4th root of 97·41 ?

TO EXTRACT THE CUBE ROOT.

I. *By the Common Rule.**

1. HAVING divided the given number into periods of three figures each, (by setting a point over the place of units, and also over every third figure, from thence, to the left hand in whole numbers, and to the right in decimals), find the nearest less cube to the first period; set its root in the quotient, and subtract the said cube from the first period; to the remainder bring down the second period, and call this the resolvend.

2. To three times the square of the root, just found, add three times the root itself, setting this one place more to the right than the former, and call this sum the divisor. Then divide the resolvend, wanting the last figure, by the divisor, for the next figure of the root, which annex to the former; calling this last figure e , and the part of the root before found let be called a .

Add all together these three products, namely, thrice a square multiplied by e , thrice a multiplied by e square, and e cube, setting each of them one place more to the right than the former, and call the sum the subtrahend; which must not exceed the resolvend; but if it does, then make the last figure e less, and repeat the operation for finding the subtrahend, till it be less than the resolvend.

4. From the resolvend take the subtrahend, and to the remainder join the next period of the given number for a new resolvend; to which form a new divisor from the whole root now found; and from thence another figure of the root, as directed in Article 2, and so on.

* The reason for pointing the given number into periods of three figures each; is because the cube of one figure never amounts to more than three places. And, for a similar reason, a given number is pointed into periods of four figures for the 4th root, of five figures for the 5th root, and so on.

And the reason for the other parts of the rule depends on the algebraic formation of a cube: for if the root consists of the two parts $a+b$, then its cube is as follows: $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$; where a is the root of the first part a^3 ; the resolvend is $3a^2b + 3ab^2 + b^3$, which is also the same as the three parts of the subtrahend; also the divisor is $3a^2 + 3a$, by which dividing the first two terms of the resolvend $3a^2b + ab^2$, gives b for the second part of the root; and so on.

EXAMPLE.

To extract the cube root of 48228·544.

$$\begin{array}{r|l}
 3 \times 3^2 = 27 & 48228 \cdot 544 \text{ (36} \cdot 4 \text{ root.} \\
 3 \times 3 = 09 & 27 \\
 \hline
 \text{Divisor } 279 & 21228 \text{ resolvend.} \\
 \hline
 3 \times 3^2 \times 6 = 162 & \\
 3 \times 3 \times 6^2 = 324 & \left. \begin{array}{l} \\ \\ \end{array} \right\} \text{ add} \\
 6^3 = 216 & \\
 \hline
 3 \times 36^2 = 3888 & 19656 \text{ subtrahend.} \\
 3 \times 36 = 108 & \\
 \hline
 38988 & 1572544 \text{ resolvend.} \\
 \hline
 3 \times 36^2 \times 4 = 15552 & \\
 3 \times 36 \times 4^2 = 1728 & \left. \begin{array}{l} \\ \\ \end{array} \right\} \text{ add} \\
 4^3 = 64 & \\
 \hline
 & 1572544 \text{ subtrahend.} \\
 \hline
 & 000000 \text{ remainder.} \\
 \hline
 \end{array}$$

Ex. 2. Extract the cube root of 571482·19.

Ex. 3. Extract the cube root of 1628·1582.

Ex. 4. Extract the cube root of 1332.

II. To extract the Cube Root by a short Way.*

1. By trials, or by the table of roots at p. 90, &c. take the nearest rational cube to the given number, whether it be greater or less; and call it the assumed cube.

2. Then

* The method usually given for extracting the cube root, is so exceedingly tedious, and difficult to be remembered, that various other approximating rules have been invented; viz, by Newton, Raphson, Halley, De Lagny, Simpson, Emerson, and several other mathematicians; but no one that I have yet seen, is so simple in its form, or seems so well adapted for general use, as that above given.

2. Then say, by the Rule of Three, As the sum of the given number and double the assumed cube, is to the sum of the assumed cube and double the given number, so is the root of the assumed cube, to the root required, nearly. Or, As the first sum is to the difference of the given and assumed cube, so is the assumed root to the difference of the roots nearly.

3. Again, by using in like manner, the cube of the root last found as a new assumed cube, another root will be obtained still nearer. And so on as far as we please; using always the cube of the last found root, for the assumed cube.

EXAMPLE.

To find the cube root of 21035·8.

Here we soon find that the root lies between 20 and 30, and then between 27 and 28. Taking therefore 27, its cube is 19683, which is the assumed cube. Then

19683	21035·8
2	2
39366	42071·6
21035·8	19683

As 60401·8:61754·6 :: 27 : 27·6047.

4322822
1235092

60401·8) 1667374·2 (27·6047 the root nearly.
 459338
 36525
 284
 42

en. This rule is the same in effect as Dr. Halley's rational formula, but more commodiously expressed; and the first investigation of it was given in my Tracts, p. 49. The algebraic form of it is this:

$$\begin{aligned} & ASP + 2A : A + 2P :: r : R. \quad \text{Or,} \\ & ASP + 2A : P \omega A :: r : R \omega r; \end{aligned}$$

where P is the given number, A the assumed nearest cube, r the cube root of A, and a the root of P sought.

Again,

EXAMPLE.

To extract the 5th root of 21035.8.

Here it appears that the 5th root is between 7.3 and 7.4. Taking 7.3. its 5th power is 20730.71593. Hence we have $P = 21035.8$, $n = 5$, $r = 7.3$ and $A = 20730.71593$; then

$$n + 1. \frac{1}{2} A + n - 1. \frac{1}{2} P : P \omega A :: r : R \omega r, \text{ that is}$$

$$3 \times 20730.71593 + 2 \times 21035.8 : 305.084 :: 7.3 :$$

62192.14779	42071.6	915252
42071.6		2135588
104263.74779	2227.1132	(.0213605 = R ω r 7.3 = r, add.
		7.321360 = R, true to the last figure.

OTHER EXAMPLES.

- | | |
|---------------------------------------|-----------------|
| 1. What is the 3d root of 2 ? | Ans. 1.259921. |
| 2. What is the 3d root of 3214 ? | Ans. 14.75758. |
| 3. What is the 4th root of 2 ? | Ans. 1.189207. |
| 4. What is the 4th root of 97.41 ? | Ans. 3.1415999. |
| 5. What is the 5th root of 2 ? | Ans. 1.148699. |
| 6. What is the 6th root of 21035.8 ? | Ans. 5.254037. |
| 7. What is the 6th root of 2 ? | Ans. 1.122462. |
| 8. What is the 7th root of 21035.8 ? | Ans. 4.145392. |
| 9. What is the 7th root of 2 ? | Ans. 1.104089. |
| 10. What is the 8th root of 21035.8 ? | Ans. 3.470323. |
| 11. What is the 8th root of 2 ? | Ans. 1.090508. |
| 12. What is the 9th root of 21035.8 ? | Ans. 3.022239. |
| 13. What is the 9th root of 2 ? | Ans. 1.080059. |

The following is a table of squares and cubes, as also the square roots and cube roots, of all numbers from 1 to 1000 which will be found very useful on many occasions, in numerical calculations, when roots or powers are concerned.

90 A TABLE OF SQUARES, CUBES, AND ROOTS.

Number.	Square.	Cube.	Square Root.	Cube Root.
1	1	1	1.000000	1.000000
2	4	8	1.4142136	1.259921
3	9	27	1.7320508	1.442250
4	16	64	2.0000000	1.587401
5	25	125	2.2360680	1.709976
6	36	216	2.4494897	1.817121
7	49	343	2.6457513	1.912933
8	64	512	2.8284271	2.000000
9	81	729	3.0000000	2.080084
10	100	1000	3.1622777	2.154435
11	121	1331	3.3166248	2.223980
12	144	1728	3.4641016	2.289428
13	169	2197	3.6055513	2.351335
14	196	2744	3.7416574	2.410142
15	225	3375	3.8729833	2.466212
16	256	4096	4.0000000	2.519842
17	289	4913	4.1231056	2.571282
18	324	5832	4.2426407	2.620741
19	361	6859	4.3588989	2.668402
20	400	8000	4.4721360	2.714418
21	441	9261	4.5825757	2.758923
22	484	10648	4.6904158	2.802039
23	529	12167	4.7958315	2.843867
24	576	13824	4.8989795	2.884499
25	625	15625	5.0000000	2.924018
26	676	17576	5.0990195	2.962496
27	729	19683	5.1961524	3.000000
28	784	21952	5.2915026	3.036589
29	841	24389	5.3851648	3.072317
30	900	27000	5.4772256	3.107232
31	961	29791	5.5677644	3.141381
32	1024	32768	5.6568542	3.174802
33	1089	35937	5.7445626	3.207534
34	1156	39304	5.8309519	3.239612
35	1225	42875	5.9160798	3.271066
36	1296	46656	6.0000000	3.301927
37	1369	50653	6.0827625	3.332222
38	1444	54872	6.1644140	3.361975
39	1521	59319	6.2449980	3.391211
40	1600	64000	6.3245553	3.419952
41	1681	68921	6.4031242	3.448217
42	1764	74088	6.4807407	3.476027
43	1849	79507	6.5574385	3.503398
44	1939	85184	6.6332496	3.530348
45	2025	91125	6.7082039	3.556893
46	2116	97336	6.7823300	3.583048
47	2209	103823	6.8556546	3.608826
48	2304	110592	6.9282032	3.634241
49	2401	117649	7.0000000	3.659306
50	2500	125000	7.0710678	3.684031

Number.	Square.	Cube.	Square Root.	Cube Root.
51	2601	132651	7.1414284	3.708430
52	2704	140608	7.2111026	3.732511
53	2809	148877	7.2801099	3.756286
54	2916	157464	7.3484692	3.779763
55	3025	166375	7.4161985	3.802953
56	3136	175616	7.4833148	3.825862
57	3249	185193	7.5498344	3.848501
58	3364	195112	7.6157731	3.870877
59	3481	205379	7.6811457	3.892996
60	3600	216000	7.7459667	3.914867
61	3721	226981	7.8102497	3.936497
62	3844	238328	7.8740079	3.957892
63	3969	250047	7.9372539	3.979057
64	4096	262144	8.0000000	4.000000
65	4225	274625	8.0622577	4.020726
66	4356	287496	8.1240384	4.041240
67	4489	300763	8.1853528	4.061548
68	4624	314432	8.2462113	4.081656
69	4761	328509	8.3066239	4.101566
70	4900	343000	8.3666003	4.121285
71	5041	357911	8.4261498	4.140818
72	5184	373248	8.4852814	4.160168
73	5329	389017	8.5440037	4.179339
74	5476	405224	8.6023253	4.198336
75	5625	421875	8.6602540	4.217183
76	5776	438976	8.7177979	4.235824
77	5929	456533	8.7749644	4.254321
78	6084	474552	8.8317609	4.272659
79	6241	493039	8.8881944	4.290841
80	6400	512000	8.9442719	4.308870
81	6561	531441	9.0000000	4.326749
82	6724	551368	9.0553851	4.344481
83	6889	571787	9.1104336	4.362071
84	7056	592704	9.1651514	4.379519
85	7225	614125	9.2195445	4.396830
86	7396	636056	9.2736185	4.414005
87	7569	658503	9.3273791	4.431047
88	7744	681472	9.3808315	4.447960
89	7921	704969	9.4339811	4.464745
90	8100	720000	9.4868330	4.481405
91	8281	753571	9.5393920	4.497942
92	8464	778688	9.5916630	4.514357
93	8649	804357	9.6436508	4.530655
94	8836	830584	9.6953597	4.546836
95	9025	857375	9.7467943	4.562903
96	9216	884736	9.7979590	4.578857
97	9409	912673	9.8488578	4.594701
98	9604	941192	9.8994949	4.610436
99	9801	970299	9.9498744	4.626065
100	10000	1000000	10.0000000	4.641589

Numb.	Square.	Cube.	Square Root.	Cube Root.
101	10201	1030301	10.0498756	4.657010
102	10404	1061208	10.0995049	4.672330
103	10609	1092727	10.1488916	4.687548
104	10816	1124864	10.1980390	4.702669
105	11025	1157625	10.2469508	4.717694
106	11236	1191016	10.2456301	4.732624
107	11449	1225043	10.3940804	4.747459
108	11664	1259712	10.3923048	4.762203
109	11881	1295029	10.4403065	4.776856
110	12100	1331090	10.4880885	4.791420
111	12321	1367631	10.5356538	4.805896
112	12544	1404928	10.5830052	4.820284
113	12769	1442897	10.6301458	4.834588
114	12996	1481544	10.6770783	4.848808
115	13225	1520875	10.7238053	4.862944
116	13456	1560896	10.7703296	4.876999
117	13689	1601613	10.8166538	4.890973
118	13924	1643032	10.8627805	4.904868
119	14161	1685159	10.9087121	4.918685
120	14400	1728000	10.9544512	4.932424
121	14641	1771561	11.0000000	4.946088
122	14884	1815848	11.0453610	4.959675
123	15129	1860867	11.0905365	4.973190
124	15376	1906624	11.1355287	4.986631
125	15625	1953125	11.1803399	5.000000
126	15876	2000376	11.2249722	5.013298
127	16129	2048383	11.2694277	5.026526
128	16384	2097152	11.3137085	5.039684
129	16641	2146689	11.3578167	5.052774
130	16900	2197000	11.4017543	5.065797
131	17161	2248091	11.4455231	5.078753
132	17424	2299968	11.4891253	5.091643
133	17689	2352637	11.5325626	5.104469
134	17956	2406104	11.5758369	5.117230
135	18225	2460375	11.6189500	5.129928
136	18496	2515456	11.6619038	5.142563
137	18769	2571353	11.7046999	5.155137
138	19044	2628072	11.7473444	5.167649
139	19321	2685619	11.7898261	5.180101
140	19600	2744000	11.8321596	5.192494
141	19881	2803221	11.8743421	5.204828
142	20164	2863288	11.9163753	5.217103
143	20449	2924207	11.9582607	5.229321
144	20736	2985984	12.0000000	5.241482
145	21025	3048625	12.0415946	5.253588
146	21316	3112136	12.0830460	5.265637
147	21609	3176523	12.1243557	5.277632
148	21904	3241792	12.1655251	5.289572
149	22201	3307949	12.2065556	5.301459
150	22500	3375000	12.2474487	5.313293

Numb.	Square.	Cube.	Square Root.	Cube Root.
151	22801	3442951	12.2882057	5.325074
152	23104	3511808	12.3288280	5.336803
153	23409	3581577	12.3693169	5.348481
154	23716	3652264	12.4096736	5.360108
155	24025	3723875	12.4498996	5.371685
156	24336	3796416	12.4899960	5.383213
157	24649	3869893	12.5299641	5.394690
158	24964	3944312	12.5698051	5.406120
159	25281	4019679	12.6095202	5.417501
160	25600	4096000	12.6491106	5.428835
161	25921	4173281	12.6885775	5.440122
162	26244	4251528	12.7279221	5.451362
163	26569	4330747	12.7671453	5.462556
164	26896	4410944	12.8062485	5.473703
165	27225	4492125	12.8452326	5.484806
166	27556	4574296	12.8840987	5.495865
167	27889	4657463	12.9228480	5.506879
168	28224	4741632	12.9614814	5.517848
169	28561	4826809	13.0000000	5.528775
170	28900	4913000	13.0384048	5.539658
171	29241	5000211	13.0766968	5.550499
172	29584	5088448	13.1148770	5.561298
173	29929	5177717	13.1529464	5.572054
174	30276	5268024	13.1909060	5.582770
175	30625	5359375	13.2287566	5.593445
176	30976	5451776	13.2664992	5.604079
177	31329	5545233	13.3041347	5.614673
178	31684	5639752	13.3416641	5.625226
179	32041	5735339	13.3790882	5.635741
180	32400	5832000	13.4164079	5.646216
181	32761	5929741	13.4536240	5.656652
182	33124	6028568	13.4907376	5.667051
183	33480	6128487	13.5277493	5.677411
184	33836	6229504	13.5646600	5.687734
185	34225	6331625	13.6014705	5.698019
186	34596	6434856	13.6381817	5.708267
187	34969	6539203	13.6747943	5.718479
188	35344	6644672	13.7113092	5.728654
189	35721	6751269	13.7477271	5.738794
190	36100	6859000	13.7840488	5.748897
191	36481	6967871	13.8202750	5.758965
192	36864	7077888	13.8564065	5.768998
193	37249	7189057	13.8924410	5.778996
194	37636	7301384	13.9283883	5.788960
195	38025	7414875	13.9642400	5.798890
196	38416	7529536	14.0000000	5.808786
197	38809	7645373	14.0356688	5.818648
198	39204	7762392	14.0712473	5.828476
199	39601	7880599	14.1067360	5.838272
200	40000	8000000	14.1421356	5.848035

ARITHMETIC.

Numb.	Square.	Cube.	Square Root.	Cube Root.
201	40401	8120601	14.1774469	5.857765
202	40804	8242408	14.2126704	5.867464
203	41209	8365427	14.2478068	5.877130
204	41616	8489664	14.2828569	5.886765
205	42025	8615125	14.3178211	5.896368
206	42436	8741816	14.3527001	5.905941
207	42849	8869743	14.3874946	5.915481
208	43264	8998912	14.4222051	5.924991
209	43681	9123329	14.4568323	5.934473
210	44100	9261000	14.4913767	5.943911
211	44521	9393931	14.5258390	5.953341
212	44944	9528128	14.5602198	5.962731
213	45369	9663597	14.5945195	5.972091
214	45796	9800344	14.6287388	5.981426
215	46225	9938375	14.6628783	5.990927
216	46656	10077696	14.6969385	6.000000
217	47089	10218313	14.7309199	6.009244
218	47524	10360232	14.7648231	6.018463
219	47961	10503459	14.7986486	6.027650
220	48400	10648000	14.8323970	6.036811
221	48841	10793861	14.8660687	6.045943
222	49284	10941048	14.8996644	6.055048
223	49729	11089567	14.9331845	6.064126
224	50176	11239424	14.9666295	6.073177
225	50625	11390625	15.0000000	6.082201
226	51076	11543176	15.0332964	6.091199
227	51529	11697083	15.0665192	6.100170
228	51984	11852352	15.0996689	6.109115
229	52441	12008989	15.1327460	6.118032
230	52900	12167000	15.1657509	6.126925
231	53361	12326391	15.1986842	6.135792
232	53824	12487168	15.2315462	6.144634
233	54289	12649337	15.2643375	6.153449
234	54756	12812904	15.2970585	6.162239
235	55225	12977875	15.3297097	6.171005
236	55696	13144256	15.3622915	6.179747
237	56169	13312053	15.3948043	6.188463
238	56644	13481272	15.4272486	6.197154
239	57121	13651919	15.4596248	6.205821
240	57600	13824000	15.4919334	6.214464
241	58081	13997521	15.5241747	6.223083
242	58564	14172488	15.5563492	6.231678
243	59049	14348907	15.5884573	6.240251
244	59536	14526784	15.6204994	6.248800
245	60025	14706125	15.6524758	6.257324
246	60516	14886936	15.6843871	6.265826
247	61009	15069223	15.7162336	6.274304
248	61504	15252992	15.7480157	6.282760
249	62001	15438249	15.7797338	6.291194
250	62500	15625000	15.8113883	6.299604

Numb.	Square.	Cube.	Square Root.	Cube Root.
251	63001	15813251	15.8429795	6.307992
252	63504	16003008	15.8745079	6.316359
253	64009	16194277	15.9059737	6.324704
254	64516	16387064	15.9373775	6.333025
255	65025	16581375	15.9687194	6.341325
256	65536	16777216	16.0000000	6.349602
257	66049	16974593	16.0312195	6.357859
258	66564	17173512	16.0623784	6.366095
259	67081	17373979	16.0934769	6.374310
260	67600	17576000	16.1245155	6.382504
261	68121	17779581	16.1554944	6.390676
262	68644	17984728	16.1864141	6.398827
263	69169	18191447	16.2172747	6.406958
264	69696	18399744	16.2480768	6.415068
265	70225	18609625	16.2788206	6.423157
266	70756	18821096	16.3095064	6.431226
267	71289	19034163	16.3401346	6.439275
268	71824	19248832	16.3707055	6.447305
269	72361	19465109	16.4012195	6.455314
270	72900	19683000	16.4316767	6.463304
271	73441	19902511	16.4620776	6.471274
272	73984	20123648	16.4924225	6.479224
273	74529	20346417	16.5227116	6.487153
274	75076	20570824	16.5529454	6.495064
275	75625	20796875	16.5831240	6.502956
276	76176	21024576	16.6132477	6.510829
277	76729	21253933	16.6433170	6.518684
278	77284	21484952	16.6733320	6.526519
279	77841	21717639	16.7032931	6.534335
280	78400	21952000	16.7332005	6.542132
281	78961	22188041	16.7630546	6.549911
282	79524	22425768	16.7928556	6.557672
283	80089	22665187	16.8226038	6.565415
284	80656	22906304	16.8522995	6.573139
285	81225	23149125	16.8819430	6.580844
286	81796	23393656	16.9115345	6.588531
287	82369	23639903	16.9410743	6.596202
288	82944	23887872	16.9705627	6.603854
289	83521	24137569	17.0000000	6.611488
290	84100	24389000	17.0295864	6.619106
291	84681	24642171	17.0587221	6.626705
292	85264	24897088	17.0880075	6.634287
293	85849	25153757	17.1172428	6.641851
294	86436	25412184	17.1464282	6.649399
295	87025	25672375	17.1755640	6.656930
296	87616	25934336	17.2046505	6.664443
297	88209	26198073	17.2336879	6.671940
298	88804	26463592	17.2626765	6.679419
299	89401	26730899	17.2916165	6.686882
300	90000	27000000	17.3205081	6.694328

Numb.	Square.	Cube.	Square Root.	Cube Root.
301	90601	27270901	17.3493516	6.701758
302	91204	27543608	17.3781472	6.709172
303	91809	27818127	17.4068952	6.716569
304	92416	28094464	17.4355958	6.723950
305	93025	28372625	17.4642492	6.731316
306	93636	28652616	17.4928557	6.738665
307	94249	28934443	17.5214155	6.745997
308	94864	29218112	17.5499288	6.753313
309	95481	29503629	17.5783958	6.760614
310	96100	29791000	17.6068169	6.767899
311	96721	30080231	17.6351921	6.775168
312	97344	30371328	17.6635217	6.782422
313	97969	30664297	17.6918060	6.789661
314	98596	30959144	17.7200451	6.796884
315	99225	31281875	17.7482393	6.804091
316	99856	31554496	17.7763888	6.811284
317	100489	31855013	17.8044938	6.818461
318	101124	32157432	17.8325545	6.825624
319	101761	32461759	17.8605711	6.832771
320	102400	32768000	17.8885438	6.839903
321	103041	33076161	17.9104729	6.847021
322	103684	33386248	17.9443584	6.854124
323	104329	33698267	17.9722008	6.861211
324	104976	34012224	18.0000000	6.868284
325	105625	34328125	18.0277564	6.875343
326	106276	34645976	18.0554701	6.882388
327	106929	34965783	18.0831413	6.889419
328	107584	35287552	18.1107703	6.896435
329	108241	35611289	18.1383571	6.903436
330	108900	35937000	18.1659021	6.910423
331	109561	36264691	18.1934054	6.917396
332	110224	36594368	18.2208672	6.924355
333	110889	36926037	18.2482876	6.931300
334	111556	37259704	18.2756669	6.938232
335	112225	37595375	18.3030052	6.945149
336	112896	37933056	18.3303028	6.952053
337	113569	38272753	18.3575598	6.958943
338	114244	38614472	18.4847763	6.965819
339	114921	38958219	18.4119526	6.972682
340	115600	39304000	18.4390889	6.979532
341	116281	39651821	18.4661853	6.986369
342	116964	40001688	18.4932420	6.993191
343	117649	40353607	18.5202592	7.000000
344	118336	40707584	18.5472370	7.006796
345	119025	41063625	18.5741756	7.013579
346	119716	41421736	18.6010752	7.020349
347	120409	41781923	18.6279360	7.027106
348	121104	42144192	18.6547581	7.033850
349	121801	42508549	18.6815417	7.040581
350	122500	42875000	18.7082869	7.047208

Numb.	Square.	Cube.	Square Root.	Cube Root.
351	123201	43243551	18.7349940	7.054003
352	123904	43614208	18.7616630	7.060696
353	124609	43986977	18.7882942	7.067376
354	125316	44361864	18.8148877	7.074043
355	126025	44738875	18.8414437	7.080698
356	126736	45118016	18.8679623	7.087341
357	127449	45499293	18.8944436	7.093970
358	128164	45882712	18.9208879	7.100588
359	128881	46268279	18.9472953	7.107193
360	129600	46656000	18.9736660	7.113786
361	130321	47045881	19.0000000	7.120367
362	131044	47437928	19.0262976	7.126935
363	131769	47832147	19.0525589	7.133492
364	132496	48228544	19.0787840	7.140037
365	133225	48627125	19.1049732	7.146569
366	133956	49027896	19.1311265	7.153090
367	134689	49430863	19.1572441	7.159599
368	135424	49836032	19.1833261	7.166095
369	136161	50243409	19.2093727	7.172580
370	136900	50653000	19.2353841	7.179054
371	137641	51064811	19.2613603	7.185516
372	138384	51478848	19.2873015	7.191966
373	139129	51895117	19.3132079	7.198405
374	139876	52313624	19.3390796	7.204832
375	140625	52734375	19.3649167	7.211247
376	141376	53157376	19.3907194	7.217652
377	142129	53582633	19.4164878	7.224045
378	142884	54010152	19.4422221	7.230427
379	143641	54439939	19.4679223	7.236797
380	144400	54872000	19.4935887	7.243156
381	145161	55306341	19.5192213	7.249504
382	145924	55742968	19.5448203	7.255841
383	146689	56181887	19.5703858	7.262167
384	147456	56623104	19.5959179	7.268482
385	148225	57066625	19.6214169	7.274786
386	148996	57512456	19.6468827	7.281079
387	149769	57960603	19.6723146	7.287362
388	150544	58411072	19.6977156	7.293633
389	151321	58863869	19.7230829	7.299893
390	152100	59319000	19.7484177	7.306143
391	152881	59776471	19.7737199	7.312383
392	153664	60236288	19.7989899	7.318611
393	154449	60698457	19.8242276	7.324829
394	155236	61162984	19.8494332	7.331037
395	156025	61629875	19.8746069	7.337234
396	156816	62099136	19.8997487	7.343420
397	157609	62570773	19.9248588	7.349596
398	158404	63044792	19.9499373	7.355762
399	159201	63521199	19.9749844	7.361917
400	160000	64000000	20.0000000	7.368063

Numb.	Square.	Cube.	Square Root.	Cube Root.
401	160801	64481201	20.0249844	7.374198
402	161604	64964808	20.0499377	7.380322
403	162409	65450827	20.0748599	7.386437
404	163216	65939264	20.0997512	7.392542
405	164025	66430125	20.1246118	7.398636
406	164836	66923416	20.1494417	7.404720
407	165649	67419143	20.1742410	7.410794
408	166464	67911312	20.1990099	7.416859
409	167281	68417929	20.2237484	7.422914
410	168100	68921000	20.2484567	7.428958
411	168921	69426531	20.2731349	7.434993
412	169744	69934528	20.2977831	7.441018
413	170569	70444997	20.3224014	7.447033
414	171396	70957944	20.3469899	7.453039
415	172225	71473375	20.3715488	7.459036
416	173056	71991296	20.3960781	7.465022
417	173889	72511713	20.4205779	7.470999
418	174724	73034632	20.4450483	7.476966
419	175561	73560059	20.4694895	7.482924
420	176400	74088000	20.4939015	7.488872
421	177241	74618461	20.5182845	7.494810
422	178084	75151448	20.5426386	7.500740
423	178929	75686967	20.5669638	7.506660
424	179776	76225024	20.5912603	7.512571
425	180625	76765625	20.6155281	7.518473
426	181476	77308776	20.6397674	7.524365
427	182329	77854483	20.6639783	7.530248
428	183184	78402752	20.6881609	7.536121
429	184041	78953589	20.7123152	7.541986
430	184900	79507000	20.7364114	7.547841
431	185761	80062991	20.7605395	7.553688
432	186624	80621568	20.7846097	7.559525
433	187489	81182737	20.8086520	7.565353
434	188356	81746504	20.8326667	7.571173
435	189225	82312875	20.8566536	7.576984
436	190096	82881856	20.8806130	7.582786
437	190969	83453453	20.9045450	7.588579
438	191844	84027672	20.9284495	7.594363
439	192721	84604519	20.9523268	7.600138
440	193600	85184000	20.9761771	7.605905
441	194481	85766121	21.0000000	7.611662
442	195364	86350888	21.0237960	7.617411
443	196249	86938307	21.0475652	7.623151
444	197136	87528384	21.0713075	7.628883
445	198025	88121125	21.0950231	7.634606
446	198916	88716536	21.1187121	7.640321
447	199809	89314623	21.1423745	7.646027
448	200704	89915392	21.1660105	7.651725
449	201601	90518849	21.1896201	7.657414
450	202500	91125000	21.2132034	7.663094

Numb.	Square.	Cube.	Square Root.	Cube Root.
451	203401	91733851	21.2367606	7.668766
452	204304	92345408	21.2602916	7.674430
453	205209	92959677	21.2837967	7.680085
454	206116	93576664	21.3072758	7.685732
455	207025	94196375	21.3307290	7.691371
456	207936	94818816	21.3541565	7.697002
457	208849	95443993	21.3775583	7.702624
458	209764	96071912	21.4009346	7.708238
459	210681	96702579	21.4242853	7.713844
460	211600	97336000	21.4476106	7.719412
461	212521	97972181	21.4709166	7.725032
462	213444	98611128	21.4941853	7.730614
463	214369	99252847	21.5174348	7.736187
464	215296	99897344	21.5406592	7.741753
465	216225	100544625	21.5638587	7.747310
466	217156	101194696	21.5870331	7.752860
467	218089	101847563	21.6101828	7.758402
468	219024	102503232	21.6333077	7.763936
469	219961	103161709	21.6564078	7.769462
470	220900	103823000	21.6794834	7.774980
471	221841	104487111	21.7025344	7.780490
472	222784	105154048	21.7255610	7.785992
473	223729	105823817	21.7485632	7.791487
474	224676	106496424	21.7715411	7.796974
475	225625	107171875	21.7944947	7.802453
476	226576	107850176	21.8174242	7.807925
477	227529	108531333	21.8403297	7.813389
478	228484	109215352	21.8632111	7.818845
479	229441	109902239	21.8860686	7.824294
480	230400	110592000	21.9089023	7.829735
481	231361	111284641	21.9317122	7.835168
482	232324	111980168	21.9544984	7.840594
483	233289	112678587	21.9772610	7.846013
484	234256	113379904	22.0000000	7.851424
485	235225	114084125	22.0227155	7.856828
486	236196	114791256	22.0454077	7.862224
487	237169	115501303	22.0680765	7.867613
488	238144	116214272	22.0907220	7.872994
489	239121	116930169	22.1133444	7.878368
490	240100	117649000	22.1359436	7.883734
491	241081	118370771	22.1585198	7.889094
492	242064	119095488	22.1810730	7.894446
493	243049	119823157	22.2036033	7.899791
494	244036	120553784	22.2261408	7.905129
495	245025	121287375	22.2485955	7.910460
496	246016	122023936	22.2710575	7.915784
497	247009	122763473	22.2934968	7.921100
498	248004	123505992	22.3159136	7.926408
499	249001	124251499	22.3383079	7.931710
500	250000	125000000	22.3606798	7.937005

Numb.	Square.	Cube.	Square Root.	Cube Root.
501	251001	125751501	22.3830293	7.942293
502	252004	126506008	22.4053565	7.947573
503	253009	127263527	22.4276615	7.952847
504	254016	128024064	22.4499443	7.958114
505	255025	128787625	22.4722051	7.963374
506	256036	129554216	22.4944438	7.968627
507	257049	130323843	22.5166605	7.973873
508	258064	131096512	22.5388553	7.979112
509	259081	131872229	22.5610283	7.984344
510	260100	132651000	22.5831796	7.989569
511	261121	133432831	22.6053091	7.994788
512	262144	134217728	22.6274170	8.000000
513	263169	135005697	22.6495033	8.005205
514	264196	135796744	22.6715681	8.010403
515	265225	136590875	22.6936114	8.015595
516	266256	137388096	22.7156334	8.020779
517	267289	138188413	22.7376340	8.025957
518	268324	138991832	22.7596134	8.031129
519	269361	139798359	22.7815715	8.036293
520	270400	140608000	22.8035085	8.041451
521	271441	141420761	22.8254244	8.046603
522	272484	142236648	22.8473193	8.051748
523	273529	143056667	22.8691933	8.056886
524	274576	143877824	22.8910463	8.062018
525	275625	144703125	22.9128785	8.067143
526	276676	145531576	22.9346899	8.072262
527	277729	146363183	22.9564806	8.077374
528	278784	147197952	22.9782506	8.082480
529	279841	148035889	23.0000000	8.087579
530	280900	148877000	23.0217289	8.092672
531	281961	149721291	23.0434372	8.097758
532	283024	150568768	23.0651252	8.102838
533	284089	151419437	23.0867928	8.107912
534	285156	152273304	23.1084400	8.112980
535	286225	153130375	23.1300670	8.118041
536	287296	153990656	23.1516738	8.123096
537	288369	154854153	23.1732605	8.128144
538	289441	155720872	23.1948270	8.133186
539	290521	156590819	23.2163735	8.138223
540	291600	157464000	23.2379001	8.143253
541	292681	158340421	23.2594067	8.148276
542	293764	159220088	23.2808935	8.153293
543	294849	160103007	23.3023604	8.158304
544	295936	160989184	23.3238076	8.163309
545	297025	161878625	23.3452351	8.168308
546	298116	162771336	23.3666429	8.173302
547	299209	163666323	23.3880311	8.178289
548	300304	164566592	23.4093998	8.183269
549	301401	165469149	23.4307490	8.188244
550	302500	166375000	23.4520788	8.193212

Numb.	Square.	Cube.	Square Root.	Cube Root.
551	303601	167284151	23.4733892	8.198175
552	304704	168196608	23.4946802	8.203131
553	305809	169112377	23.5159520	8.208082
554	306916	170031464	23.5372046	8.213027
555	308025	170953875	23.5584380	8.217965
556	309136	171879616	23.5796522	8.222898
557	310249	172808693	23.6008474	8.227825
558	311364	173741112	23.6220236	8.232746
559	312481	174676879	23.6431808	8.237661
560	313600	175616000	23.6643191	8.242570
561	314721	176558481	23.6854386	8.247474
562	315844	177504328	23.7065392	8.252371
563	316969	178453547	23.7276210	8.257263
564	318096	179406144	23.7486842	8.262149
565	319225	180362125	23.7697286	8.267029
566	320356	181321496	23.7907545	8.271903
567	321489	182284263	23.8117618	8.276772
568	322624	183250432	23.8327506	8.281635
569	323761	184220009	23.8537209	8.286493
570	324900	185193000	23.8746728	8.291344
571	326041	186169411	23.8956063	8.296190
572	327184	187149248	23.9165215	8.301030
573	328329	188132517	23.9374184	8.305865
574	329476	189119224	23.9582971	8.310694
575	330625	190109375	23.9791576	8.315517
576	331776	191102976	24.0000000	8.320335
577	332929	192100033	24.0208243	8.325147
578	334084	193100552	24.0416306	8.329954
579	335241	194104539	24.0624188	8.334755
580	336400	195112000	24.0831892	8.339551
581	337561	196122941	24.1039416	8.344341
582	338724	197137368	24.1246762	8.349125
583	339889	198155287	24.1453929	8.353904
584	341056	199176704	24.1660919	8.358678
585	342225	200201625	24.1867732	8.363446
586	343396	201230056	24.2074369	8.368209
587	344569	202262003	24.2280829	8.372966
588	345744	203297472	24.2487113	8.377718
589	346921	204336469	24.2693222	8.382465
590	348100	205379000	24.2899156	8.387206
591	349281	206425071	24.3104916	8.391942
592	350464	207474688	24.3310501	8.396673
593	351649	208527857	24.3515913	8.401398
594	352836	209584584	24.3721152	8.406118
595	354025	210644875	24.3926218	8.410832
596	355216	211708736	24.4131112	8.415541
597	356409	212776173	24.4335834	8.420245
598	357604	213847192	24.4540385	8.424944
599	358801	214921799	24.4744765	8.429638
600	360000	216000000	24.4948974	8.434327

Numb.	Square.	Cube.	Square Root.	Cube Root.
601	361201	217081801	24.5153013	8.439009
602	362404	218167208	24.5356883	8.443687
603	363609	219256227	24.5560583	8.448360
604	364816	220348864	24.5764115	8.453027
605	366025	221445125	24.5967478	8.457689
606	367236	222545016	24.6170673	8.462347
607	368449	223648543	24.6373700	8.466999
608	369664	224755712	24.6576560	8.471647
609	370881	225866529	24.6779254	8.476289
610	372100	226981000	24.6981781	8.480926
611	373321	228099131	24.7184142	8.485557
612	374544	229220928	24.7386338	8.490184
613	375769	230346397	24.7588368	8.494806
614	376996	231475544	24.7790234	8.499423
615	378225	232608375	24.7991935	8.504034
616	379456	233744896	24.8193473	8.508641
617	380689	234885113	24.8394847	8.513243
618	381924	236029032	24.8596058	8.517840
619	383161	237176659	24.8797106	8.522432
620	384400	238328000	24.8997992	8.527018
621	385641	239483061	24.9198716	8.531600
622	386884	240641848	24.9399278	8.536177
623	388129	241804367	24.9599679	8.540749
624	389376	242970624	24.9799920	8.545317
625	390625	244140625	25.0000000	8.549879
626	391876	245314376	25.0199920	8.554437
627	393129	246491883	25.0399681	8.558990
628	394384	247673152	25.0599282	8.563537
629	395641	248858189	25.0798724	8.568080
630	396900	250047000	25.0998008	8.572618
631	398161	251239591	25.1197134	8.577152
632	399424	252435968	25.1396102	8.581680
633	400689	253636137	25.1594913	8.586204
634	401956	254840104	25.1793566	8.590723
635	403225	256047875	25.1992063	8.595238
636	404496	257259456	25.2190404	8.599747
637	405769	258474853	25.2388589	8.604252
638	407044	259694072	25.2586619	8.608752
639	408321	260917119	25.2784493	8.613248
640	409600	262144000	25.2982213	8.617738
641	410881	263374721	25.3179778	8.622224
642	412164	264609288	25.3377189	8.626706
643	413449	265847707	25.3574447	8.631183
644	414736	267089984	25.3771551	8.635655
645	416025	268336125	25.3968502	8.640122
646	417316	269586136	25.4165301	8.644585
647	418609	270840023	25.4361947	8.649043
648	419904	272097792	25.4558441	8.653497
649	421201	273359449	25.4754784	8.657946
650	422500	274625000	25.4950076	8.662301

Numb.	Square.	Cube.	Square Root.	Cube Root.
651	423801	275894451	25.5147016	8.666831
652	425104	277167808	25.5342907	8.671266
653	426409	278445077	25.5538647	8.675697
654	427716	279726264	25.5734237	8.680123
655	429025	281011375	25.5929678	8.684545
656	430336	282300416	25.6124969	8.688963
657	431649	283593393	25.6320112	8.693376
658	432964	284890312	25.6515107	8.697784
659	434281	286191179	25.6709953	8.702188
660	435600	287496000	25.6904652	8.706587
661	436921	288804781	25.7099203	8.710982
662	438244	290117528	25.7203607	8.715373
663	439569	291434247	25.7487864	8.719759
664	440896	292754944	25.7681975	8.724141
665	442225	294079625	25.7875939	8.728518
666	443556	295408296	25.8069758	8.732891
667	444889	296740963	25.8263431	8.737260
668	446224	298077632	25.8456960	8.741624
669	447561	299418309	25.8650343	8.745984
670	448900	300763000	25.8843582	8.750340
671	450241	302111711	25.9036677	8.754691
672	451584	303464448	25.9229628	8.759038
673	452929	304821217	25.9422435	8.763380
674	454276	306182024	25.9615100	8.767719
675	455625	307546875	25.9807621	8.772053
676	456976	308915776	26.0000000	8.776382
677	458329	310288733	26.0192237	8.780708
678	459684	311665752	26.0384331	8.785029
679	461040	313046839	26.0576284	8.789346
680	462400	314432000	26.0768096	8.793659
681	463761	315821241	26.0959767	8.797967
682	465124	317214568	26.1151297	8.802272
683	466489	318611987	26.1342687	8.806572
684	467856	320013504	26.1533937	8.810868
685	469225	321419125	26.1725047	8.815159
686	470596	322828856	26.1916017	8.819447
687	471969	324242703	26.2106848	8.823730
688	473344	325660672	26.2297541	8.828009
689	474721	327082769	26.2488095	8.832285
690	476100	328509000	26.2678511	8.836556
691	477481	329939371	26.2868789	8.840822
692	478864	331373888	26.3058929	8.845085
693	480249	332812557	26.3248932	8.849344
694	481636	334255384	26.3438797	8.853598
695	483025	335702375	26.3628527	8.857849
696	484416	337153536	26.3818119	8.862095
697	485809	338608873	26.4007576	8.866337
698	487204	340068392	26.4196896	8.870575
699	488601	341532099	26.4386081	8.874809
700	490000	343000000	26.4575131	8.879040

Numb.	Square.	Cube.	Square Root.	Cube Root.
701	491401	344472101	26.4764046	8.883266
702	492804	345948008	26.4952826	8.887488
703	494209	347428927	26.5141472	8.891706
704	495616	348913664	26.5329983	8.895920
705	497025	350402625	26.5518361	8.900130
706	498436	351895816	26.5706605	8.904336
707	499849	353393243	26.5894716	8.908538
708	501264	354894912	26.6082694	8.912736
709	502681	356400829	26.6270539	8.916931
710	504100	357911000	26.6458252	8.921121
711	505521	359425431	26.6645833	8.925307
712	506944	360944128	26.6833281	8.929490
713	508369	362467097	26.7020598	8.933668
714	509796	363994344	26.7207784	8.937843
715	511225	365525875	26.7394839	8.942014
716	512656	367061696	26.7581763	8.946180
717	514089	368601813	26.7768557	8.950343
718	515524	370146232	26.7955220	8.954502
719	516961	371694959	26.8141754	8.958658
720	518400	373248000	26.8328157	8.962809
721	519841	374805361	26.8514432	8.966957
722	521284	376367048	26.8700577	8.971100
723	522729	377933067	26.8886593	8.975240
724	524176	379503424	26.9072481	8.979376
725	525625	381078125	26.9258240	8.983508
726	527076	382657176	26.9443872	8.987637
727	528529	384240583	26.9629375	8.991762
728	529984	385828353	26.9814751	9.995883
729	531441	387420489	27.0000000	9.000000
730	532900	389017000	27.0185122	9.004113
731	534361	390617891	27.0370117	9.008223
732	535824	392223168	27.0554985	9.012328
733	537289	393832837	27.0739727	9.016430
734	538756	395446904	27.0924344	9.020529
735	540225	397065375	27.1108834	9.024623
736	541696	398688256	27.1293199	9.028714
737	543169	400315553	27.1477439	9.032802
738	544644	401947272	27.1661554	9.036885
739	546121	403583419	27.1845544	9.040965
740	547600	405224000	27.2029410	9.045041
741	549081	406869021	27.2213152	9.049114
742	550564	408518488	27.2396769	9.053183
743	552049	410172407	27.2580263	9.057248
744	553536	411830784	27.2763634	9.061309
745	555025	413493625	27.2946881	9.065367
746	556516	415160936	27.3130006	9.069422
747	558009	416832723	27.3313007	9.073472
748	559504	418508992	27.3495887	9.077519
749	561001	420189749	27.3678644	9.081563
750	562500	421875000	27.3861279	9.085603

Numb.	Square.	Cube.	Square Root.	Cube Root.
751	564001	423564751	27.4043792	9.089639
752	565504	425259008	27.4226184	9.093672
753	567009	426957777	27.4408455	9.097701
754	568516	428661064	27.4590604	9.101726
755	570025	430368875	27.4772633	9.105748
756	571536	432081216	27.4954542	9.109766
757	573049	433798093	27.5136330	9.113781
758	574564	435519512	27.5317998	9.117793
759	576081	437245479	27.5499546	9.121801
760	577600	438976000	27.5680975	9.125805
761	579121	440711081	27.5862284	9.129806
762	580644	442450728	27.6043475	9.133803
763	582169	444194947	27.6224546	9.137797
764	583696	445943744	27.6405499	9.141788
765	585225	447697125	27.6586334	9.145774
766	586756	449455096	27.6767050	9.149757
767	588289	451217663	27.6947648	9.153737
768	589824	452984832	27.7128129	9.157713
769	591361	454756609	27.7308492	9.161686
770	592900	456533000	27.7488739	9.165656
771	594441	458314011	27.7668868	9.169622
772	595984	460099648	27.7848880	9.173585
773	597529	461889917	27.8028775	9.177544
774	599076	463684824	27.8208555	9.181500
775	600625	465484375	27.8388218	9.185452
776	602176	467288576	27.8567766	9.189401
777	603729	469097433	27.8747197	9.193347
778	605284	470910952	27.8926514	9.197289
779	606841	472729139	27.9105715	9.201228
780	608400	474552000	27.9284801	9.205164
781	609961	476379541	27.9463772	9.209096
782	611524	478211768	27.9642629	9.213025
783	613089	480048687	27.9821372	9.216950
784	614656	481890304	28.0000000	9.220872
785	616225	483736025	28.0178515	9.224791
786	617796	485587656	28.0356915	9.228706
787	619369	487443403	28.0535203	9.232618
788	620944	489303872	28.0713377	9.237527
789	622521	491169069	28.0891438	9.240433
790	624100	493039000	28.1069386	9.244335
791	625681	494913671	28.1247222	9.248234
792	627264	496793088	28.1424946	9.252130
793	628849	498677257	28.1602557	9.256022
794	630436	500566184	28.1780056	9.259911
795	632025	502459875	28.1957444	9.263797
796	633616	504358336	28.2134720	9.267679
797	635209	506261573	28.2311884	9.271559
798	636804	508169592	28.2488938	9.275435
799	638401	510082399	28.2665881	9.279308
800	640000	512000000	28.2842712	9.283177

Numb.	Square.	Cube.	Square Root.	Cube Root.
801	641601	513922401	28.3019434	9.287044
802	643204	515849608	28.3196045	9.290907
803	644809	517781627	28.3372546	9.294767
804	646416	519718464	28.3548938	9.298623
805	648025	521660125	28.3725219	9.302477
806	649636	523606616	28.3901391	9.306327
807	651249	525557943	28.4077454	9.310175
808	652864	527514112	28.4253408	9.314019
809	654481	529475129	28.4429253	9.317859
810	656100	531441000	28.4604989	9.321697
811	657721	533411731	28.4780617	9.325532
812	659344	535387328	28.4956137	9.329363
813	660969	537366797	28.5131549	9.333191
814	662596	539353144	28.5306852	9.337016
815	664225	541343375	28.5482048	9.340838
816	665856	543338496	28.5657187	9.344657
817	667489	545338513	28.5832119	9.348473
818	669124	547343432	28.6006993	9.352285
819	670761	549353259	28.6181760	9.356095
820	672400	551368000	28.6356421	9.359901
821	674041	553387661	28.6530976	9.363704
822	675684	555412248	28.6705424	9.367505
823	677329	557441767	28.6879766	9.371302
824	678976	559476224	28.7054002	9.375096
825	680625	561515625	28.7228132	9.378887
826	682276	563559976	28.7402157	9.382675
827	683929	565609283	28.7576077	9.386460
828	685584	567663552	28.7749891	9.390241
829	687241	569722789	28.7923601	9.394020
830	688900	571787000	28.8097206	9.397796
831	690561	573856191	28.8270706	9.401569
832	692224	575930368	28.8444102	9.405338
833	693889	578009537	28.8617394	9.409105
834	695556	580093704	28.8790582	9.412869
835	697225	582182875	28.8963666	9.416630
836	698896	584277056	28.9136646	9.420387
837	700569	586376253	28.9309523	9.424141
838	702244	588480472	28.9482297	9.427893
839	703921	590589719	28.9654967	9.431642
840	705600	592704000	28.9827535	9.435388
841	707281	594823321	29.0000000	9.439130
842	708964	596947688	29.0172363	9.442870
843	710649	599077107	29.0344623	9.446607
844	712336	601211584	29.0516781	9.450341
845	714025	603351125	29.0688837	9.454071
846	715716	605495736	29.0860791	9.457799
847	717409	607645423	29.1032644	9.461524
848	719104	609800192	29.1204396	9.465247
849	720801	611960049	29.1376046	9.468966
850	722500	614125000	29.1547505	9.472682

Numb.	Square.	Cube.	Square Root.	Cube Root.
851	724201	616295051	29.1719043	9.476395
852	725904	618470208	29.1890390	9.480106
853	727609	620650477	29.2061637	9.483813
854	729316	622835864	29.2232784	9.487518
855	731025	625026375	29.2403830	9.491219
856	732736	627222016	29.2574777	9.494918
857	734449	629422793	29.2745623	9.498614
858	736164	631628712	29.2916370	9.502307
859	737881	633839779	29.3087018	9.505998
860	739600	636056000	29.3257566	9.509685
861	741321	638277381	29.3428015	9.513369
862	743044	640503928	29.3598365	9.517051
863	744769	642735647	29.3768616	9.520730
864	746496	644972544	29.3938769	9.524406
865	748225	647214625	29.4108823	9.528079
866	749956	649461896	29.4278779	9.531749
867	751689	651714363	29.4448637	9.535417
868	753424	653972032	29.4618397	9.539081
869	755161	656234909	29.4788059	9.542748
870	756900	658503000	29.4957624	9.546402
871	758641	660776311	29.5127091	9.550058
872	760384	663054848	29.5296461	9.553712
873	762129	665338617	29.5465734	9.557363
874	763876	667627624	29.5634910	9.561010
875	765625	669921875	29.5803989	9.564655
876	767376	672221376	29.5972972	9.568297
877	769129	674526133	29.6141858	9.571937
878	770884	676836152	29.6310648	9.575574
879	772641	679151439	29.6479325	9.579208
880	774400	681472000	29.6647939	9.582839
881	776161	683797841	29.6816442	9.586468
882	777924	686128968	29.6984848	9.590093
883	779689	688465387	29.7153159	9.593716
884	781456	690807104	29.7321375	9.597337
885	783225	693154125	29.7489496	9.600951
886	784996	695506456	29.7657521	9.604569
887	786769	697864103	29.7825452	9.608181
888	788544	700227072	29.7993289	9.611791
889	790321	702595369	29.8161030	9.615397
890	792100	704969000	29.8328678	9.619001
891	793881	707347971	29.8496231	9.622603
892	795664	709732288	29.8663690	9.626201
893	797449	712121957	29.8831056	9.629797
894	799236	714516984	29.8998328	9.633390
895	801025	716917375	29.9165506	9.636981
896	802816	719323136	29.9332591	9.640569
897	804609	721734273	29.9499583	9.644154
898	806404	724150792	29.9666481	9.647736
899	808201	726572699	29.9833287	9.651316
900	810000	729000000	30.0000000	9.654893

Numb.	Square.	Cube.	Square Root.	Cube Root.
901	811801	731432701	30.0166620	9.658468
902	813604	733870808	30.0333148	9.662040
903	815409	7363 4327	30.0499584	9.665609
904	817216	738763264	30.0665928	9.669176
905	819025	741217625	30.0832179	9.672740
906	820836	743677416	30.0998339	9.676301
907	822649	746142643	30.1164407	9.679860
908	824464	748613312	30.1330383	9.683416
909	826281	751089429	30.1496269	9.686970
910	828100	753571000	30.1632063	9.690521
911	829921	756058031	30.1827765	9.694069
912	831744	758550528	30.1993377	9.697615
913	833569	761048497	30.2158899	9.701158
914	835396	763551944	30.2324329	9.704698
915	837225	766060875	30.2489669	9.708236
916	839056	768575296	30.2654919	9.711772
917	840889	771095213	30.2820079	9.715305
918	842724	773620632	30.2985148	9.718835
919	844561	776151559	30.3150128	9.722363
920	846400	778688000	30.3315018	9.725888
921	848241	781229961	30.3479818	9.729410
922	850084	783777448	30.3644529	9.732930
923	851929	786330467	30.3809151	9.736448
924	853776	788889024	30.3973683	9.739963
925	855625	791453125	30.4138127	9.743475
926	857476	794022776	30.4302481	9.746985
927	859329	796597983	30.4466747	9.750493
928	861184	799178752	30.4630924	9.753998
929	863041	801765089	30.4795013	9.757500
930	864900	804357000	30.4959014	9.761000
931	866761	806954491	30.5122926	9.764497
932	868624	809557568	30.5286750	9.767992
933	870489	812166237	30.5450487	9.771484
934	872356	814780504	30.5614136	9.774974
935	874225	817400375	30.5777697	9.778461
936	876096	820025856	30.5941171	9.782946
937	877969	822656953	30.6104557	9.785428
938	879844	825293672	30.6267857	9.788908
939	881721	827936019	30.6431069	9.792386
940	883600	830584000	30.6594194	9.795861
941	885481	833237621	30.6757233	9.799333
942	887364	835896888	30.6920185	9.802803
943	889249	838561807	30.7083051	9.806271
944	891136	841232884	30.7245830	9.809736
945	893025	843908625	30.7408523	9.813198
946	894916	846590536	30.7571130	9.816659
947	896809	849278123	30.7733651	9.820117
948	898704	851971392	30.7896086	9.823572
949	900601	854670349	30.8058436	9.827025
950	902500	857375000	30.8220700	9.830475

SQUARES, CUBES, AND ROOTS.

109

Numb.	Square.	Cube.	Square Root.	Cube Root.
951	904401	860085351	30.8382879	9.833923
952	906304	862801408	30.8544972	9.837369
953	908209	865523177	30.8706981	9.840812
954	910116	868250664	30.8868904	9.844253
955	912025	870983875	30.9030743	9.847692
956	913936	873722816	30.9192497	9.851128
957	915849	876467493	30.9354166	9.854561
958	917764	879217912	30.9515751	9.857992
959	919681	881974079	30.9677251	9.861421
960	921600	884736000	30.9838668	9.861848
961	923521	887503681	31.0000000	9.868272
962	925444	890277128	31.0161248	9.871694
963	927369	893056347	31.0322413	9.875113
964	929296	895841344	31.0483494	9.878530
965	931225	893632125	31.0644491	9.881945
966	933156	901428696	31.0805405	9.885357
967	935089	904231063	31.0966236	9.888767
968	937024	907039232	31.1126984	9.892174
969	938961	909853209	31.1287648	9.895580
970	940900	912673000	31.1448230	9.898983
971	942841	915498611	31.1608729	9.902383
972	944784	918330048	31.1769145	9.905781
973	946729	921167317	31.1929479	9.909177
974	948676	924010424	31.2089731	9.912571
975	950625	926859375	31.2249900	9.915962
976	952576	929714176	31.2409987	9.919351
977	954529	932574833	31.2569992	9.922738
978	956484	935441352	31.2729915	9.926122
979	958441	938313739	31.2889757	9.929504
980	960400	941192001	31.3049517	9.932883
981	962361	944076141	31.3209195	9.936261
982	964324	946966168	31.3368792	9.939636
983	966289	949862087	31.3528308	9.943009
984	968256	952763904	31.3687743	9.946379
985	970225	955671625	31.3847097	9.949747
986	972196	958585256	31.4006369	9.953113
987	974169	961504803	31.4165561	9.956477
988	976144	964430272	31.4324673	9.959839
989	978121	967361669	31.4483704	9.963198
990	980100	970299000	31.4642654	9.966554
991	982081	973242271	31.4801525	9.969909
992	984064	976191488	31.4960315	9.973262
993	986049	979146657	31.5119025	9.976612
994	988036	982107784	31.5277655	9.979959
995	990025	985074875	31.5436206	9.983304
996	992016	988047936	31.5594677	9.986648
997	994009	991026973	31.5753068	9.989990
998	996004	994011992	31.5911380	9.993328
999	998001	997002999	31.6069613	9.996665

OF RATIOS, PROPORTIONS, AND PROGRESSIONS.

NUMBERS are compared to each other in two different ways: the one comparison considers the difference of the two numbers, and is named Arithmetical Relation; and the difference sometimes the Arithmetical Ratio: the other considers their quotient, which is called Geometrical Relation; and the quotient is the Geometrical Ratio. So, of these two numbers 6 and 3, the difference, or arithmetical ratio, is $6 - 3$ or 3, but the geometrical ratio is $\frac{6}{3}$ or 2.

There must be two numbers to form a comparison: the number which is compared, being placed first, is called the Antecedent; and that to which it is compared, the Consequent. So, in the two numbers above, 6 is the antecedent, and 3 the consequent.

If two or more couplets of numbers have equal ratios, or equal differences, the equality is named Proportion, and the terms of the ratios Proportionals. So, the two couplets, 4, 2 and 8, 6, are arithmetical proportionals, because $4 - 2 = 8 - 6 = 2$; and the two couplets 4, 2 and 6, 3, are geometrical proportionals, because $\frac{4}{2} = \frac{6}{3} = 2$, the same ratio.

To denote numbers as being geometrically proportional, a colon is set between the terms of each couplet, to denote their ratio; and a double colon, or else a mark of equality between the couplets or ratios. So, the four proportionals, 4, 2, 6, 3 are set thus, $4 : 2 :: 6 : 3$, which means, that 4 is to 2 as 6 is to 3; or thus, $4 : 2 = 6 : 3$, or thus, $\frac{4}{2} = \frac{6}{3}$, both which mean, that the ratio of 4 to 2, is equal to the ratio of 6 to 3.

Proportion is distinguished into Continued and Discontinued. When the difference or ratio of the consequent of one couplet, and the antecedent of the next couplet, is not the same as the common difference or ratio of the couplets, the proportion is discontinued. So, 4, 2, 8, 6 are in discontinued arithmetical proportion, because $4 - 2 = 8 - 6 = 2$, whereas $8 - 2 = 6$: and 4, 2, 6, 3 are in discontinued geometrical proportion, because $\frac{4}{2} = \frac{6}{3} = 2$, but $\frac{6}{2} = 3$, which is not the same.

But when the difference or ratio of every two succeeding terms is the same quantity, the proportion is said to be Continued, and the numbers themselves make a series of Continued Proportionals,

Proportionals, or a progression. So 2, 4, 6, 8 form an arithmetical progression, because $4 - 2 = 6 - 4 = 8 - 6 = 2$, all the same common difference; and 2, 4, 8, 16 a geometrical progression, because $\frac{4}{2} = \frac{8}{4} = \frac{16}{8} = 2$, all the same ratio.

When the following terms of a progression increase, or exceed each other it is called an Ascending Progression, or Series; but when the terms decrease, it is a descending one. So, 0, 1, 2, 3, 4, &c. is an ascending arithmetical progression, but 9, 7, 5, 3, 1, &c. is a descending arithmetical progression. Also 1, 2, 4, 8, 16, &c. is an ascending geometrical progression, and 16, 8, 4, 2, 1, &c. is a descending geometrical progression.

ARITHMETICAL PROPORTION *and* PROGRESSION.

IN Arithmetical Progression, the numbers or terms have all the same common difference. Also, the first and last terms of a Progression, are called the **Extremes**; and the other terms, lying between them, the **Means**. The most useful part of arithmetical proportions, is contained in the following theorems:

THEOREM 1. When four quantities are in arithmetical proportion, the sum of the two extremes is equal to the sum of the two means. Thus, of the four 2, 4, 6, 8, here $2 + 8 = 4 + 6 = 10$.

THEOREM 2. In any continued arithmetical progression, the sum of the two extremes is equal to the sum of any two means that are equally distant from them, or equal to double the middle term when there is an uneven number of terms.

Thus, in the terms 1, 3, 5, it is $1 + 5 = 3 + 3 = 6$.

And in the series 2, 4, 6, 8, 10, 12, 14, it is $2 + 14 = 4 + 12 = 6 + 10 = 8 + 8 = 16$.

THEOREM 3. The difference between the extreme terms of an arithmetical progression is equal to the common difference of the series multiplied by one less than the number of the terms. So, of the ten terms, 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, the common difference is 2, and one less than the number of terms 9; then the difference of the extremes is $20 - 2 = 18$, and $2 \times 9 = 18$ also.

Consequently,

Consequently, the greatest term is equal to the least term added to the product of the common difference multiplied by 1 less than the number of terms.

THEOREM 4. The sum of all the terms, of any arithmetical progression, is equal to the sum of the two extremes multiplied by the number of terms, and divided by 2; or the sum of the two extremes multiplied by the number of the terms, gives double the sum of all the terms in the series.

This is made evident by setting the terms of the series in an inverted order, under the same series in a direct order, and adding the corresponding terms together in that order. Thus, in the series 1, 3, 5, 7, 9, 11, 13, 15; ditto inverted 15, 13, 11, 9, 7, 5, 3, 1;

the sums are $16 + 16 + 16 + 16 + 16 + 16 + 16 + 16$, which must be double the sum of the single series, and is equal to the sum of the extremes repeated as often as are the number of the terms.

From these theorems may readily be found any one of these five parts; the two extremes, the number of terms, the common difference, and the sum of all the terms, when any three of them are given; as in the following problems:

PROBLEM. I.

Given the extremes, and the Number of Terms; to find the Sum of all the Terms.

ADD the extremes together, multiply the sum by the number of terms, and divide by 2.

EXAMPLES.

1. The extremes being 3 and 19, and the number of terms 9; required the sum of the terms?

$$\begin{array}{r} 19 \\ 3 \\ \hline \end{array}$$

$$\begin{array}{r} 22 \\ 9 \\ \hline \end{array}$$

$$2) 198$$

$$\hline \text{Ans. } 99$$

$$\text{Or, } \frac{19+3}{2} \times 9 = \frac{22}{2} \times 9 = 11 \times 9 = 99.$$

the same answer.

2. It is required to find the number of all the strokes a common clock strikes in one whole revolution of the index, or in 12 hours?

Ans. 78.

Ex.

Ex. 3. How many strokes do the clocks of Venice strike in the compass of a day, which go continually on from 1 to 24 o'clock? Ans. 300.

4. What debt can be discharged in a year, by weekly payments in arithmetical progression, the first payment being 1s, and the last or 52d payment 5l 3s? Ans. 135l 4s.

PROBLEM II.

Given the Extremes, and the Number of Terms; to find the Common Difference.

SUBTRACT the less extreme from the greater, and divide the remainder by 1 less than the number of terms, for the common difference.

EXAMPLES.

1. The extremes being 3 and 19, and the number of terms 9; required the common difference?

$$\begin{array}{r} 19 \\ 3 \\ \hline 8) 16 \\ \hline \text{Ans. } 2 \end{array} \quad \text{Or, } \frac{19-3}{9-1} = \frac{16}{8} = 2.$$

2. If the extremes be 10 and 70, and the number of terms 21; what is the common difference, and the sum of the series? Ans. the com. diff. is 3, and the sum is 840.

3. A certain debt can be discharged in one year, by weekly payments in arithmetical progression, the first payment being 1s, and the last 5l 3s; what is the common difference of the terms? Ans. 2.

PROBLEM III.

Given one of the Extremes, the Common Difference, and the Number of Terms; to find the other Extreme, and the sum of the Series.

MULTIPLY the common difference by 1 less than the number of terms, and the product will be the difference of the extremes: Therefore add the product to the less extreme, to give the greater; or subtract it from the greater, to give the less extreme.

EXAMPLES.

1. Given the least term 3, the common difference 2, of an arithmetical series of 9 terms; to find the greatest term, and the sum of the series.

$$\begin{array}{r}
 2 \\
 8 \\
 \hline
 16 \\
 3 \\
 \hline
 19 \text{ the greatest term} \\
 3 \text{ the least} \\
 \hline
 22 \text{ sum} \\
 9 \text{ number of terms.} \\
 \hline
 2) 198 \\
 \hline
 99 \text{ the sum of the series.} \\
 \hline
 \end{array}$$

2. If the greatest term be 70, the common difference 3, and the number of terms 21, what is the least term, and the sum of the series?

Ans. The least term is 10, and the sum is 840.

3. A debt can be discharged in a year, by paying 1 shilling the first week, 3 shillings the second, and so on, always 2 shillings more every week; what is the debt, and what will the last payment be?

Ans. The last payment will be 5*l* 3*s*, and the debt is 135*l* 4*s*.

PROBLEM IV.

To find an Arithmetical Mean between two given Terms.

ADD the two given extremes or terms together, and take half their sum for the arithmetical mean required

EXAMPLE.

To find an arithmetical mean between the two numbers 4 and 14.

$$\begin{array}{r}
 \text{Here} \\
 14 \\
 4 \\
 \hline
 2) 18 \\
 \hline
 \end{array}$$

Ans. 9 the mean required.

PROBLEM

PROBLEM V.

To find two Arithmetical Means between Two Given Extremes.

SUBTRACT the less extreme from the greater, and divide the difference by 3, so will the quotient be the common difference; which being continually added to the less extreme, or taken from the greater, gives the means.

EXAMPLE.

To find two arithmetical means between 2 and 8.

$$\begin{array}{r} \text{Here } 8 \\ \quad 2 \\ \hline 3 \overline{) 6} \\ \hline \text{com. dif. } 2 \\ \hline \end{array}$$

Then $2 + 2 = 4$ the one mean
and $4 + 2 = 6$ the other mean.

PROBLEM VI.

To find any Number of Arithmetical Means between Two Given Terms or Extremes.

SUBTRACT the less extreme from the greater, and divide the difference by 1 more than the number of means required to be found, which will give the common difference; then this being added continually to the least term, or subtracted from the greatest, will give the terms required.

EXAMPLE.

To find five arithmetical means between 2 and 14.

$$\begin{array}{r} \text{Here } 14 \\ \quad 2 \\ \hline 6 \overline{) 12} \\ \hline \text{com. dif. } 2 \\ \hline \end{array}$$

Then by adding this com. dif. continually,
the means are found 4, 6, 8, 10, 12.

See more of Arithmetical progression in the Algebra.

PROPORTION AND GEOMETRICAL PROGRESSION ;

Or Progression by equal Ratios.

In Geometrical Progression the numbers or terms have all the same multiplier or divisor. The most useful part of Proportion, is contained in the following theorems.

THEOREM 1. When four quantities are in proportion, the product of the two extremes is equal to the product of the two means.

Thus, in the four 2, 4, 3, 6, it is $2 \times 6 = 3 \times 4 = 12$.

And hence, if the product of the two means be divided by one of the extremes, the quotient will give the other extreme. So, of the above numbers, the product of the means $12 \div 2 = 6$ the one extreme, and $12 \div 6 = 2$ the other extreme ; and this is the foundation and reason of the practice in the Rule of Three.

THEOREM 2. In any continued geometrical progression, the product of the two extremes is equal to the product of any two means that are equally distant from them, or equal to the square of the middle term when there is an uneven number of terms.

Thus, in the terms 2, 4, 8, it is $2 \times 8 = 4 \times 4 = 16$.

And in the series 2, 4, 8, 16, 32, 64, 128,
it is $2 \times 128 = 4 \times 64 = 8 \times 32 = 16 \times 16 = 256$.

THEOREM 3. The quotient of the extreme terms of a geometrical progression, is equal to the common ratio of the series raised to the power denoted by 1 less than the number of the terms. Consequently the greatest term is equal to the least term multiplied by the said quotient.

So, of the ten terms, 2, 4, 8, 16, 32, 64, 128, 256, 512, 1024, the common ratio is 2, and one less than the number of term is 9 ; then the quotient of the extremes is $1024 \div 2 = 512$, and $2^9 = 512$ also.

THEOREM

THEOREM 4. The sum of all the terms, of any geometrical progression, is found by adding the greatest term to the difference of the extremes divided by 1 less than the ratio.

So, the sum of 2, 4, 8, 16, 32, 64, 128, 256, 512, 1024, $\frac{1024-2}{2-1}$
 (whose ratio is 2), is $1024 + \frac{1024-2}{2-1} = 1024 + 1022 = 2046$.

The foregoing, and several other properties of proportion, are demonstrated more at large in the Algebraic part of this work. A few examples may here be added of the theorems, just delivered, with some problems concerning mean proportionals.

EXAMPLES.

1. The least of ten terms, in geometrical progression, being 1, and the ratio 2; what is the greatest term, and the sum of all the terms?

Ans. The greatest term is 512, and the sum 1023.

2. What debt may be discharged in a year, or 12 months, by paying 1*l* the first month, 2*l* the second, 4*l* the third, and so on, each succeeding payment being double the last; and what will the last payment be?

Ans. The debt 4095*l*, and the last payment 2048*l*.

PROBLEM I.

To find One Geometrical Mean Proportional between any two Numbers.

MULTIPLY the two numbers together, and extract the square root of the product, which will give the mean proportional sought.

EXAMPLE.

To find a geometrical mean between the two numbers 3 and 12.

$$\begin{array}{r} 12 \\ 3 \\ \hline 36 \text{ (6 the mean.)} \\ \hline 36 \end{array}$$

PROBLEM

PROBLEM II.

To find Two Geometrical Mean Proportionals between any Two Numbers.

DIVIDE the greater number by the less, and extract the cube root of the quotient, which will give the common ratio of the terms. Then multiply the least given term by the ratio for the first mean, and this mean again by the ratio for the second mean : or, divide the greater of the two given terms by the ratio for the greater mean, and divide this again by the ratio for the less mean.

EXAMPLE.

To find two geometrical means between 3 and 24.

Here $3 \sqrt[3]{24}$ (8 ; its cube root 2 is the ratio.

Then $3 \times 2 = 6$, and $6 \times 2 = 12$, the two means.

Or $24 \div 2 = 12$, and $12 \div 2 = 6$, the same.

That is, the two means between 3 and 24, are 6 and 12.

PROBLEM III.

To find any Number of Geometrical Means between Two Numbers.

DIVIDE the greater number by the less, and extract such root of the quotient whose index is 1 more than the number of means required, that is, the 2d root for one mean, the 3d root for two means, the 4th root for three means, and so on ; and that root will be the common ratio of all the terms. Then, with the ratio, multiply continually from the first term, or divide continually from the last or greatest term.

EXAMPLE.

To find four geometrical means between 3 and 96.

Here $3 \sqrt[5]{96}$ (32 ; the 5th root of which is 2, the ratio.

Then $3 \times 2 = 6$, & $6 \times 2 = 12$, & $12 \times 2 = 24$, & $24 \times 2 = 48$.

Or $96 \div 2 = 48$, & $48 \div 2 = 24$, & $24 \div 2 = 12$, & $12 \div 2 = 6$.

That is, 6, 12, 24, 48, are the four means between 3 and 96.

OF

OF MUSICAL PROPORTION.

THERE is also a third kind of proportion, called Musical, which being but of little or no common use, a very short account of it may here suffice.

Musical Proportion is when, of three numbers, the first has the same proportion to the third, as the difference between the first and second, has to the difference between the second and third.

As in these three, 6, 8, 12 ;
 where $6 : 12 :: 8 - 6 : 12 - 8$,
 that is $6 : 12 :: 2 : 4$.

When four numbers are in musical proportion ; then the first has the same ratio to the fourth, as the difference between the first and second has to the difference between the third and fourth.

As in these, 6, 8, 12, 18 ;
 where $6 : 18 :: 8 - 6 : 18 - 12$.
 that is $6 : 18 :: 2 : 6$.

When numbers are in musical progression, their reciprocals are in arithmetical progression ; and the converse, that is, when numbers are in arithmetical progression, their reciprocals are in musical progression.

So in these musicals 6, 8, 12, their reciprocals $\frac{1}{6}$, $\frac{1}{8}$, $\frac{1}{12}$, are in arithmetical progression ; for $\frac{1}{6} + \frac{1}{12} = \frac{3}{12} = \frac{1}{4}$; and $\frac{1}{8} + \frac{1}{8} = \frac{2}{8} = \frac{1}{4}$; that is, the sum of the extremes is equal to double the mean, which is the property of arithmeticals.

The method of finding out numbers in musical proportion is best expressed by letters in Algebra.

 FELLOWSHIP, OR PARTNERSHIP.

FELLOWSHIP is a rule, by which any sum or quantity may be divided into any number of parts, which shall be in any given proportion to one another.

By this rule are adjusted the gains or loss or charges of
 partners

partners in company ; or the effects of bankrupts, or legacies in case of a deficiency of assets or effects ; or the shares of prizes ; or the numbers of men to form certain detachments ; or the division of waste lands among a number of proprietors.

Fellowship is either Single or Double. It is Single, when the shares or portions are to be proportional each to one single given number only ; as when the stocks of partners are all employed for the same time : And Double, when each portion is to be proportional to two or more numbers ; as when the stocks of partners are employed for different times.

SINGLE FELLOWSHIP.

GENERAL RULE.

ADD together the numbers that denote the proportion of the shares. Then say,

As the sum of the said proportional numbers,
Is to the whole sum to be parted or divided,
So is each several proportional number,
To the corresponding share or part.

Or, as the whole stock, is to the whole gain or loss,
So is each man's particular stock,
To his particular share of the gain or loss.

TO PROVE THE WORK. Add all the shares or parts together, and the sum will be equal to the whole number to be shared when the work is right.

EXAMPLES.

1. To divide the number 240 into three such parts, as shall be in proportion to each other as the three numbers 1, 2 and 3.

Here $1+2+3=6$, the sum of the numbers.

Then, as $6 : 240 :: 1 : 40$ the 1st part,

and as $6 : 240 :: 2 : 80$ the 2d part,

also as $6 : 240 :: 3 : 120$ the 3d part,

Sum of all 240, the proof.

Ex. 2.

Ex. 2. Three persons, A, B, C, freighted a ship with 340 tuns of wine; of which, A loaded 110 tuns, B 97, and C the rest: in a storm the seamen were obliged to throw overboard 85 tuns; how much must each person sustain of the loss?

Here $110 + 97 = 207$ tuns, loaded by A and B;

theref. $340 - 207 = 133$ tuns, loaded by C.

Hence, as $340 : 85 :: 110$

or as $4 : 1 :: 110 : 27\frac{1}{2}$ tuns = A's loss;

and as $4 : 1 :: 97 : 24\frac{1}{4}$ tuns = B's loss;

also as $4 : 1 :: 133 : 33\frac{1}{4}$ tuns = C's loss;

Sum 85 tuns, the proof.

3. Two merchants, C and D, made a stock of 120*l.* of which C contributed 75*l.*, and D the rest: by trading they gained 30*l.*; what must each have of it?

Ans. C 18*l.* 15*s.*, and D 11*l.* 5*s.*

4. Three merchants, E, F, G, made a stock of 700*l.*, of which E contributed 123*l.*, F 358*l.*, and G the rest: by trading they gain 125*l.* 10*s.*; what must each have of it?

Ans. E must have 22*l.* 1*s.* 0*d.* $2\frac{2}{3}\frac{7}{5}$.

F - - - 64 3 8 $0\frac{2}{3}\frac{2}{5}$.

G - - - 39 5 3 $1\frac{1}{3}\frac{1}{5}$.

5. A General imposing a contribution* of 700*l.* on four villages, to be paid in proportion to the number of inhabitants contained in each; the 1st containing 250, the 2d 350, the 3d 400, and the 4th 500 persons, what part must each village pay?

Ans. the 1st to pay 116*l.* 13*s.* 4*d.*

the 2d - - - 163 6 8

the 3d - - - 186 13 4

the 4th - - - 233 6 8

6. A piece of ground, consisting of 37 ac 2 ro 14 ps. is to be divided among three persons, L, M, and N, in proportion to their estates: now if L's estate be worth 500*l.* a year, M's 320*l.*, and N's 75*l.*; what quantity of land must each one have?

Ans. L must have 20 ac 3 ro $39\frac{1}{17}\frac{3}{5}$ ps.

M - - - 13 1 $30\frac{4}{17}\frac{6}{5}$.

N - - - 3 0 $23\frac{1}{17}\frac{3}{5}$.

7. A person is indebted to O 57*l.* 15*s.*, to P 108*l.* 3*s.* 8*d.*, to Q 22*l.* 10*d.*, and to R 73*l.*; but at his decease, his effects are found

* Contribution is a tax paid by provinces, towns, villages, &c. to excuse them from being plundered. It is paid in provisions or in money, and sometimes in both.

to be worth no more than 170*l* 14*s*; how must it be divided among his creditors?

Ans. o must have 37*l* 15*s* 5*d* $2\frac{5302}{10439}q$.
 P - - - 70 15 2 $2\frac{7498}{10439}$.
 Q - - - 14 8 4 $0\frac{4720}{10439}$.
 R - - - 47 14 11 $2\frac{3358}{10439}$.

Ex. 8. A ship worth 900*l*, being entirely lost, of which $\frac{1}{8}$ belonged to s, $\frac{1}{4}$ to r, and the rest to v; what loss will each sustain, supposing 540*l* of her were insured?

Ans. s will lose 45*l*, r 90*l*, and v 225*l*.

9. Four persons, w, x, y, and z, spent among them 25*s*, and agree that their shares are to be in proportion as $\frac{1}{3}$, $\frac{1}{4}$, and $\frac{1}{5}$: what are their shares?

Ans. w must pay 9*s* 8*d* $3\frac{1}{7}q$.
 x - - - 6 5 $3\frac{5}{7}q$.
 y - - - 4 10 $1\frac{5}{7}q$.
 z - - - 3 10 $3\frac{1}{7}q$.

10. A detachment, consisting of 5 companies, being sent into a garrison, in which the duty required 76 men a day; what number of men must be furnished by each company, in proportion to their strength; the first consisting of 54 men, the 2d of 51 men, and the 3d of 48 men, the 4th of 39, and the 5th of 36 men?

Ans. The 1st must furnish 18, the 2d 17, the 3d 16, the 4th 13, and the 5th 12 men.*

DOUBLE FELLOWSHIP.

DOUBLE FELLOWSHIP, as has been said, is concerned in cases in which the stocks of partners are employed or continued for different times.

* Questions of this nature frequently occurring in military service, General Haviland, an officer of great merit, contrived an ingenious instrument, for more expeditiously resolving them; which is distinguished by the name of the inventor, being called a Haviland.

RULE.*—Multiply each person's stock by the time of its continuance ; then divide the quantity, as in Single Fellowship, into shares, in proportion to these products, by saying,
 As the total sum of all the said products,
 Is to the whole gain or loss, or quantity to be parted,
 So is each particular product,
 To the correspondent share of the gain or loss.

EXAMPLES.

1. A had in company 50*l* for 4 months, and B had 60*l* for 5 months ; at the end of which time they find 24*l* gained : how must it be divided between them ?

$$\begin{array}{r}
 \text{Here } 50 \quad 60 \\
 \quad 4 \quad 5 \\
 \hline
 200 + 300 = 500
 \end{array}$$

Then, as 500 : 24 :: 200 : $9\frac{3}{5}$ = 9*l* 12*s* = A's share.
 and as 500 : 24 :: 300 : $14\frac{2}{5}$ = 14 8 = B's share.

2. c and d hold a piece of ground in common, for which they are to pay 54*l*. c put in 23 horses for 27 days, and d 21 horses for 39 days ; how much ought each man to pay of the rent ?

Ans. c must pay 23*l* 5*s* 9*d*.
 d must pay 30 14 3

4. Three persons, E, F, G, hold a pasture in common, for which they are to pay 39*l* per annum ; into which E put 7 oxen for 3 months, F put 9 oxen for 5 months, and G put in 4 oxen for 12 months ; how much must each person pay of the rent ?

Ans. E must pay 5*l* 10*s* 6*d* $1\frac{5}{9}q$.
 F - - - 11 16 10 $0\frac{8}{9}$.
 G - - - 12 12 7 $2\frac{6}{9}$.

4. A ship's company take a prize of 1000*l*, which they agree to divide among them according to their pay and the time they have been on board : now the officers and midshipmen have been on board 6 months, and the sailors 3 months ;

* The proof of this rule is as follows : When the times are equal, the shares of the gain or loss are evidently as the stocks, as in Single Fellowship ; and when the stocks are equal, the shares as the times ; therefore, when neither are equal, the shares must be as their products.

the officers have 40s a month, the midshipmen 30s, and the sailors 22s a month; moreover there are 4 officers, 12 midshipmen, and 110 sailors; what will each man's share be?

Ans. each officer must have 23*l* 2*s* 5*d* $0\frac{2}{17\frac{2}{3}}q$.
 each midshipman - 17 6 9 $3\frac{6}{17\frac{2}{3}}$.
 each seamen - - 6 7 2 $0\frac{8}{17\frac{2}{3}}$.

Ex. 5. H, with a capital of 1000*l*, began trade the first of January, and, meeting with success in business, took in I as a partner, with a capital of 1500*l*, on the first of March following. Three months after that they admit K as a third partner, who brought into stock 2800*l*. After trading together till the end of the year, they find there has been gained 1776*l* 10*s*; how must this be divided among the partners?

Ans. H must have 457*l* 9*s* 4 $\frac{1}{4}$ *d*.
 I - - - 571 16 8 $\frac{1}{4}$.
 K - - - 747 3 11 $\frac{1}{4}$.

6. x, y, and z made a joint-stock for 12 months; x at first put in 20*l*, and 4 months after 20*l* more; y put in at first 30*l*, at the end of 3 months he put in 20*l* more, and 2 months after he put in 40*l* more; z put in at first 60*l*, and 5 months after he put in 10*l* more, 1 month after which he took out 30*l*; during the 12 months they gained 50*l*; how much of it must each have?

Ans. x must have 10*l* 18*s* 6*d* $3\frac{4}{11}q$.
 y - - - 22 8 1 $0\frac{1}{6}q$.
 z - - - 16 13 4 0.

SIMPLE INTEREST.

INTEREST is the premium or sum allowed for the loan, or forbearance of money. The money lent, or forborn, is called the Principal. And the sum of the principal and its interest, added together, is called the Amount. Interest is allowed at so much per cent. per annum; which premium per cent. per annum, or interest of 100*l* for a year, is called the rate of interest:—So,

When

When interest is at 3 per cent. the rate is 3 ;
 - - - - 4 per cent. - - 4 ;
 - - - - 5 per cent. - - 5 ;
 - - - - 6 per cent. - - 6 ;

But, by law in England, interest ought not to be taken higher than at the rate of 5 per cent

Interest is of two sorts ; Simple and Compound.

Simple Interest is that which is allowed for the principal lent or forborn only, for the whole time of forbearance. As the interest of any sum, for any time, is directly proportional to the principal sum, and also to the time of continuance ; hence arises the following general rule of calculation.

As 100*l* is to the rate of interest, so is any given principal to its interest for one year. And again,

As 1 year is to any given time, so is the interest for a year, just found, to the interest of the given sum for that time.

OTHERWISE. Take the interest of 1 pound for a year, which multiply by the given principal, and this product again by the time of loan or forbearance, in years and parts, for the interest of the proposed sum for that time.

Note, When there are certain parts of years in the time, as quarters or months, or days : they may be worked for, either by taking the aliquot or like parts of the interest of a year, or by the Rule of Three, in the usual way. Also to divide by 100, is done by only pointing off two figures for decimals.

EXAMPLES.

1. To find the interest of 230*l* 10*s*, for 1 year, at the rate of 4 per cent. per annum.

Here, As 100 : 4 :: 230*l* 10*s* : 9*l* 4*s* 4 $\frac{3}{4}$ *d*.

$$\begin{array}{r}
 100 \overline{) 9,22 \ 0} \\
 \underline{20} \\
 4 \ 40 \\
 \underline{12} \\
 4 \ 80 \\
 \underline{4} \\
 3 \ 20
 \end{array}$$

Ans. 9*l* 4*s* 4 $\frac{3}{4}$ *d*.

Ex. 2. To find the interest of 547*l* 15*s*, for 3 years, at 5 per cent. per annum.

As 100 : 5 :: 547·75 :

Or 20 : 1 :: 547·75 : 27·3875 interest for 1 year.

$$\begin{array}{r}
 \\
 \\
 \\
 \hline
 27\cdot3875 \text{ ditto for 3 years.} \\
 20 \\
 \hline
 s\ 3\cdot2500 \\
 12 \\
 \hline
 d\ 3\cdot00 \text{ Ans. } 82l\ 3s\ 3d.
 \end{array}$$

3. To find the interest of 200 guineas, for 4 years 7 months and 25 days, at 4½ per cent. per annum.

	ds	l	ds
210 <i>l</i>	As 365 ::	9·45 :	25 : l
4½	or 73 ::	9·45 :	5 : ·6472
<hr style="width: 50px; margin-left: 0;"/>		5	
840		<hr style="width: 50px; margin-left: 0;"/>	
105	73) 47·25 (·6472		
<hr style="width: 50px; margin-left: 0;"/>	345		
9·45 interest for 1 yr,	530		
4	19		
<hr style="width: 50px; margin-left: 0;"/>			
37·80 ditto 4 years.			
6 mo = ½ 4·725 ditto 6 months.			
1 mo = ⅙ ·7875 ditto 1 month.			
·6472 ditto 25 days.			

$$\begin{array}{r}
 l\ 43\cdot9597 \\
 20 \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 s\ 19\cdot1940 \\
 12 \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 d\ 2\cdot3280 \\
 4 \\
 \hline
 \end{array}$$

Ans. 43*l* 19*s* 2¼*d*.

$$\begin{array}{r}
 q\ 1\cdot3120 \\
 \hline
 \end{array}$$

4. To find the interest of 450*l*, for a year at 5 per cent. per annum. Ans. 22*l* 10*s*.

5. To find the interest of 715*l* 12*s* 6*d*, for a year, at 4½ per cent. per annum. Ans. 32*l* 4*s* 0*d*.

6. To

6. To find the interest of 720*l*, for 3 years, at 5 per cent. per annum. Ans. 108*l*.

7. To find the interest of 355*l* 15*s*, for 4 years, at 4 per cent. per annum. Ans. 56*l* 18*s* 4 $\frac{3}{4}$ *d*.

8. To find the interest of 32*l* 5*s* 8*d*, for 7 years, at 4 $\frac{1}{4}$ per cent. per annum. Ans. 9*l* 12*s* 1*d*.

9. To find the interest of 170*l*, for 1 $\frac{1}{2}$ year, at 5 per cent. per annum. Ans. 12*l* 5*s*.

10. To find the insurance on 205*l* 15*s*, for $\frac{1}{4}$ of a year, at 4 per cent. per annum. Ans. 2*l* 1*s* 1 $\frac{3}{4}$ *d*.

11. To find the interest of 319*l* 6*d*, for 5 $\frac{3}{4}$ years, at 3 $\frac{3}{4}$ per cent. per annum. Ans. 68*l* 15*s* 9 $\frac{1}{2}$ *d*.

12. To find the insurance on 207*l*, for 117 days, at 4 $\frac{3}{4}$ per cent. per annum. Ans. 1*l* 12*s* 7*d*.

13. To find the interest of 17*l* 5*s*, for 117 days, at 4 $\frac{3}{4}$ per cent. per annum. Ans. 5*s* 3*d*.

14. To find the insurance on 712*l* 6*s*, for 8 months, at 7 $\frac{1}{2}$ per cent. per annum. Ans. 35*l* 12*s* 3 $\frac{1}{2}$ *d*.

Note. The Rules for Simple Interest, serve also to calculate Insurances, or the Purchase of Stocks, or any thing else that is rated at so much per cent.

See also more on the subject of Interest, with the algebraical expression and investigation of the rules at the end of the Algebra, next following.

COMPOUND INTEREST.

COMPOUND INTEREST, called also Interest upon Interest, is that which arises from the principal and interest, taken together, as it becomes due, at the end of each stated time of payment. Though it be not lawful to lend money at Compound Interest, yet in purchasing annuities, pensions, or leases in reversion, it is usual to allow Compound Interest to the purchaser for his ready money.

RULES.—1. Find the amount of the given principal, for the time of the first payment, by Simple Interest. Then consider this amount as a new principal for the second payment, whose amount calculate as before. And so on through all the payments to the last, always accounting the last amount as a new principal for the next payment. The reason of which is evident from the definition of Compound Interest.
Or else,

2. Find the amount of 1 pound for the time of the first payment, and raise or involve it to the power whose index is denoted by the number of payments. Then that power multiplied

multiplied by the given principal, will produce the whole amount. From which the said principal being subtracted, leaves the Compound Interest of the same. As is evident from the first Rule.

EXAMPLES.

1. To find the amount of 720*l*, for 4 years, at 5 per cent. per annum,

Here 5 is the 20th part of 100, and the interest of 1*l* for a year is $\frac{1}{20}$ or .05, and its amount 1.05. Therefore,

	<i>l</i>	<i>s</i>	<i>d</i>		
20)	720	0	0	1st yr's princip.	1.05 amount of 1 <i>l</i> .
	36	0	0	1st yr's interest.	1.05
<hr/>					
20)	756	0	0	2d yr's princip.	1.1025 2d power of it.
	37	16	0	2d yr's interest.	1.1025
<hr/>					
20)	793	16	0	3d yr's princip.	1.21550625 4th pow. of it.
	39	13	9 $\frac{1}{2}$	3d yr's interest.	720
<hr/>					
20)	833	9	9 $\frac{1}{2}$	4th yr's princip.	1 875.1645
	41	13	5 $\frac{3}{4}$	4th yr's interest.	20
<hr/>					
	£ 875	3	3 $\frac{1}{4}$	the whole amount.	3.2900
<hr/>					
				or ans. required.	12
<hr/>					
					3.4800

2. To find the amount of 50*l*, in 5 years, at 5 per cent. per annum, compound interest. Ans. 63*l* 16*s* 3 $\frac{1}{4}$ *d*.

3. To find the amount of 50*l* in 5 years, or 10 half-years, at 5 per cent. per annum,* compound interest, the interest payable half-yearly. Ans. 64*l* 0*s* 1*d*.

4. To find the amount of 50*l*, in 5 years, or 20 quarters, at 5 per cent. per annum,† compound interest, the interest payable quarterly. Ans. 64*l* 2*s* 0 $\frac{1}{4}$ *d*.

5. To find the compound interest of 370*l* forborn for 6 years, at 4 per cent. per annum. Ans. 98*l* 3*s* 4 $\frac{1}{4}$ *d*.

6. To find the compound interest of 410*l* forborn for 2 $\frac{1}{2}$ years, at 4 $\frac{1}{2}$ per cent. per annum, the interest payable half-yearly. Ans. 48*l* 4*s* 11 $\frac{1}{4}$ *d*.

7. To find the amount, at compound interest, of 217*l*, forborn for 2 $\frac{1}{4}$ years, at 5 per cent. per annum, the interest payable quarterly. Ans. 242*l* 13*s* 4 $\frac{1}{2}$ *d*.

Note. See the Rules for Compound Interest algebraically investigated, at the end of the Algebra.

* That is, at 2 $\frac{1}{2}$ per cent. per half-year.

† That is, at 1 $\frac{1}{4}$ per cent. per quarter of a year.

ALLIGATION.

ALLIGATION teaches how to compound or mix together several simples of different qualities, so that the composition may be of some intermediate quality or rate. It is commonly distinguished into two cases, Alligation Medial, and Alligation Alternate.

ALLIGATION MEDIAL.

ALLIGATION MEDIAL is the method of finding the rate or quality of the composition, from having the quantities and rates or qualities of the several simples given. And it is thus performed :

* MULTIPLY the quantity of each ingredient by its rate or quality ; then add all the products together, and add also all

* *Demonstration.* The rule is thus proved by Algebra.

Let $a, b, c,$ be the quantities of the ingredients,
and $m, n, p,$ their rates or qualities, or prices ;
then $am, bn, cp,$ are their several values,
and $am, + bn + cp$ the sum of their values,
also $a + b + c$ is the sum of the quantities,
and if r denote the rate of the whole composition,

then $a + b + c \times r$ will be the value of the whole,

conseq. $a + b + c \times r = am + bn + cp,$

and $r = am + bn + cp \div a + b + c,$ which is the rule.

Note, If an ounce or any other quantity of pure gold be reduced into 24 equal parts, these parts are called Caracts ; but gold is often mixed with some base metal, which is called the Alloy, and the mixture is said to be of so many caracts fine, according to the proportion of pure gold contained in it ; thus, if 22 caracts of pure gold, and 2 of alloy be mixed together, it is said to be 22 caracts fine.

If any one of the simples be of little or no value with respect to the rest, its rate is supposed to be nothing ; as water mixed with wine, and alloy with gold and silver.

the quantities together into another sum; then divide the former sum by the latter, that is, the sum of the products by the sum of the quantities, and the quotient will be the rate or quality of the composition required.

EXAMPLES.

If three sorts of gunpowder be mixed together, viz. 50lb at $12d$ a pound, 44lb at $9d$, and 26lb at $8d$ a pound; how much a pound is the composition worth?

Here 50, 44, 26 are the quantities,
and 12, 9, 8 the rates or qualities;
then $50 \times 12 = 600$
 $44 \times 9 = 396$
 $26 \times 8 = 208$

$\begin{array}{r} \hline 120) \\ \hline \end{array}$
 $\begin{array}{r} \hline 1204 \\ \hline \end{array}$
 $(10 \frac{4}{120} = 10 \frac{1}{30}$

Ans. The rate or price is $10 \frac{1}{30}d$ the pound.

2. A composition being made of 5lb of tea at $7s$ per lb, 9lb at $8s 6d$ per lb, and $14\frac{1}{2}lb$ at $5s 10d$ per lb; what is a lb of it worth? Ans. $6s 10\frac{1}{2}d$.

3. Mixed 4 gallons of wine at $4s 10d$ per gallon, with 7 gallons at $5s 3d$ per gallon, and $9\frac{3}{4}$ gallons at $5s 8d$ per gallon; what is a gallon of this composition worth? Ans. $5s 4\frac{1}{4}d$.

4. A mealman would mix 3 bushels of flour at $3s 5d$ per bushel, 4 bushels at $5s 6d$ per bushel, and 5 bushels at $4s 8d$ per bushel; what is the worth of a bushel of this mixture? Ans. $4s 7\frac{1}{2}d$.

5. A farmer mixes 10 bushels of wheat at $5s$ the bushel, with 18 bushels of rye at $3s$ the bushel, and 20 bushels of barley at $2s$ per bushel: how much is a bushel of the mixture worth? Ans. $3s$.

6. Having melted together 7 oz of gold of 22 caracts fine, $12\frac{1}{2}$ oz of 21 caracts fine, and 17 oz of 19 caracts fine: I would know the fineness of the composition? Ans. $20\frac{1}{3}$ caracts fine.

7. Of what fineness is that composition, which is made by mixing 3lb of silver of 9 oz fine, with 5lb 8 oz of 10 oz fine, and 1lb 10 oz of alloy? Ans. $7\frac{2}{3}$ oz fine.

ALLIGATION

ALLIGATION ALTERNATE.

ALLIGATION ALTERNATE is the method of finding what quantity of any number of simples, whose rates are given, will compose a mixture of a given rate. So that it is the reverse of Alligation Medial, and may be proved by it.

RULE I.*

1. SET the rates of the simples in a column under each other. —2. Connect, or link with a continued line, the rate of each simple, which is less than that of the compound, with one, or any number, of those that are greater than the compound; and each greater rate with one or any number of the less. —3. Write the difference between the mixture rate, and that of each of the simples, opposite the rate with which they are linked. —4. Then if only one difference stand against any rate, it will be the quantity belonging to that rate; but if there be several, their sum will be the quantity.

The examples may be proved by the rule for Alligation Medial.

* *Demonst.*: By connecting the less rate to the greater, and placing the difference between them and the rate alternately, the quantities resulting are such that there is precisely as much gained by one quantity as is lost by the other, and therefore the gain and loss upon the whole is equal, and is exactly the proposed rate: and the same will be true of any other two simples managed according to the Rule.

In like manner, whatever the number of simples may be, and with how many soever every one is linked, since it is always a less with a greater than the mean price, there will be an equal balance of loss and gain between every two, and consequently an equal balance on the whole. Q. E. D.

It is obvious, from this Rule, that questions of this sort admit of a great variety of answers; for, having found one answer, we may find as many more as we please, by only multiplying or dividing each of the quantities found, by 2, or 3, or 4, &c: the reason of which is evident; for, if two quantities, of two simples, make a balance of loss and gain, with respect to the mean price, so must also the double or treble, the $\frac{1}{2}$ or $\frac{1}{3}$ part, or any other ratio of these quantities, and so on *ad infinitum*.

These kinds of questions are called by algebraists *indeterminate* or *unlimited* problems; and by an analytical process, theorems may be raised that will give all the *possible* answers.

EXAMPLES.

EXAMPLES.

1. A merchant would mix wines at 16s, at 18s, and at 22s per gallon, so as that the mixture may be worth 20s the gallon : what quantity of each must be taken ?

$$\text{Here } 20 \left\{ \begin{array}{l} 16 \\ 18 \\ 22 \end{array} \right. \begin{array}{l} 2 \text{ at } 16s \\ 2 \text{ at } 18s \\ 4+2=6 \text{ at } 22s. \end{array}$$

Ans. 2 gallons at 16s, 2 gallons at 18s, and 6 at 22s.

2. How much wine at 6s per gallon, and at 4s per gallon must be mixed together, that the composition may be worth 5s per gallon ?

Ans. 1 qt or 1 gall, &c.

3. How much sugar at 4d, at 6d, and at 11d per lb, must be mixed together, so that the composition formed by them may be worth 7d per lb ?

Ans. 1 lb, or 1 stone, or 1 cwt, or any other equal quantity of each sort.

4. How much corn at 2s 6d, 3s 8d, 4s, and 4s 8d per bushel, must be mixed together, that the compound may be worth 3s 10d per bushel ?

Ans. 2 at 2s 6d, 2 at 3s 8d, 3 at 4s, and 3 at 4s 8d.

5. A goldsmith has gold of 16, of 18, of 23, and of 24 caracts fine : how much must he take of each, to make it 21 caracts fine ?

Ans. 3 of 16, 2 of 18, 3 of 23, and 5 of 24.

6. It is required to mix brandy at 12s, wine at 10s, cyder at 1s, and water at 0 per gallon together, so that the mixture may be worth 8s per gallon ?

Ans. 8 gals of brandy, 7 of wine, 2 of cyder, and 4 of water.


RULE II.

WHEN the whole composition is limited to a certain quantity : Find an answer as before by linking ; then say, as the sum of the quantities, or differences thus determined, is to the given quantity ; so is each ingredient, found by linking, to the required quantity of each.

EXAMPLES.

1. How much gold of 15, 17, 18, and 22 caracts fine, must be mixed together, to form a composition of 40 oz of 20 caracts fine ?

Here



16

Then, as 16 : 40 :: 2 : 5
 and 16 : 40 :: 10 : 25

Ans. 5 oz of 15, of 17, and of 18 caracts fine, and 25 oz of 22 caracts fine.*

Ex. 2. A vintner has wine at 4s, at 5s, at 5s 6d, and at 6s a gallon ; and he would make a mixture of 18 gallons, so that it might be afforded at 5s 4d per gallon ; how much of each sort must he take ?

Ans. 3 gal. at 4s, 3 at 5s, 6 at 5s 6d, and 6 at 6s.

* A great number of questions might be here given relating to the specific gravities of metals, &c. but one of the most curious may here suffice.

Hiero, king of Syracuse, gave orders for a crown to be made entirely of pure gold ; but suspecting the workman had debased it by mixing it with silver or copper, he recommended the discovery of the fraud to the famous Archimedes, and desired to know the exact quantity of alloy in the crown.

Archimedes, in order to detect the imposition, procured two other masses, the one of pure gold, the other of silver or copper, and each of the same weight with the former ; and by putting each separately into a vessel full of water the quantity of water expelled by them determined their specific gravities ; from which, and their given weights, the exact quantities of gold and alloy in the crown may be determined.

Suppose the weight of each crown to be 10lb, and that the water expelled by the copper or silver was 92lb, by the gold 52lb, and by the compound crown 64lb ; what will be the quantities of gold and alloy in the crown ?

The rates of the simples are 92 and 52, and of the compound 64 ; therefore.

$$64 \begin{cases} 92 - 12 \text{ of copper} \\ 52 - 28 \text{ of gold} \end{cases}$$

And the sum of these is 12 + 28 = 40, which should have been but 10 ; therefore by the Rule,

$$\begin{cases} 40 : 10 :: 12 : 3\text{lb of copper} \\ 40 : 10 :: 28 : 7\text{lb of gold} \end{cases} \text{ the answer.}$$

RULE III.*

WHEN one of the ingredients is limited to a certain quantity ; Take the difference between each price, and the mean rate as before ; then say, As the difference of that simple, whose quantity is given, is to the rest of the differences severally ; so is the quantity given, to the several quantities required.

EXAMPLES.

1. How much wine at 5s, at 5s 6d, and 6s the gallon, must be mixed with 3 gallons at 4s per gallon, so that the mixture may be worth 5s 4d per gallon ?

$$\begin{array}{r} \text{Here } 64 \left\{ \begin{array}{l} 48 \\ 60 \\ 66 \\ 72 \end{array} \right. \begin{array}{l} 8+2=10 \\ 8+2=10 \\ 16+4=20 \\ 16+4=20 \end{array} \end{array}$$

$$\text{Then } 10 : 10 :: 3 : 3$$

$$10 : 20 :: 3 : 6$$

$$10 : 20 :: 3 : 6$$

Ans. 3 gallons at 5s, 6 at 5s 6d, and 6 at 6s.

2. A grocer would mix teas at 12s, 10s, and 6s per lb, with 20lb at 4s per lb. how much of each sort must he take to make the composition worth 8s per lb ?

Ans. 20lb at 4s, 10lb at 6s, 10lb at 10s, and 20lb at 12s.

3. How much gold of 15, of 17, and of 22 caracts fine, must be mixed with 5 oz of 18 caracts fine, so that the composition may be 20 caracts fine ?

Ans. 5 oz of 15 caracts fine, 5 oz of 17, and 25 of 22.

* In the very same manner questions may be wrought when several of the ingredients are limited to certain quantities, by finding first for one limit, and then for another. The two last Rules can need no demonstration, as they evidently result from the first, the reason of which has been already explained.

POSITION.

POSITION is a method of performing certain questions, which cannot be resolved by the common direct rules. It is sometimes called False Position, or False Supposition, because it makes a supposition of false numbers, to work with the same as if they were the true ones, and by their means discovers the true numbers sought. It is sometimes also called Trial-and-Error, because it proceeds by *trials* of false numbers, and thence finds out the true ones by a comparison of the *errors*.—Position is either Single or Double.

SINGLE POSITION.

SINGLE POSITION is that by which a question is resolved by means of one supposition only. Questions which have their result proportional to their suppositions, belong to Single Position: such as those which require the multiplication or division of the number sought by any proposed number; or when it is to be increased or diminished by itself, or any parts of itself, a certain proposed number of times. The rule is as follows:

TAKE or assume any number for that which is required, and perform the same operations with it, as are described or performed in the question. Then say, As the result of the said operation, is to the position, or number assumed; so is the result in the question, to a fourth term, which will be the number sought.*

* The reason of this Rule is evident, because it is supposed that the results are proportional to the suppositions.

Thus, $na : a :: nz : z,$

or $\frac{a}{n} : a :: \frac{z}{n} : z,$

or $\frac{a}{n} \pm \frac{a}{m} \&c. : a :: \frac{z}{n} \pm \frac{z}{m} \&c. : z,$

and so on.

EXAMPLES.

EXAMPLES.

1. A person after spending $\frac{1}{3}$ and $\frac{1}{4}$ of his money, has yet remaining 60*l*; what had he at first?

Suppose he had at first 120*l*.

Now $\frac{1}{3}$ of 120 is 40

$\frac{1}{4}$ of it is 30

their sum is 70
which taken from 120

leaves 50

Then, 50 : 120 :: 60 : 144, the Answer.

Proof.

$\frac{1}{3}$ of 144 is 48

$\frac{1}{4}$ of 144 is 36

their sum 84
taken from 144

leaves 60 as
per question.

2. What number is that which being multiplied by 7, and the product divided by 6, the quotient may be 21? Ans. 18.

3. What number is that, which being increased by $\frac{1}{2}$, $\frac{1}{3}$, and $\frac{1}{4}$ of itself, the sum shall be 75? Ans. 36.

4. A general, after sending out a foraging $\frac{1}{2}$ and $\frac{1}{3}$ of his men. had yet remaining 1000; what number had he in command? Ans. 6000.

5. A gentleman distributed 52 pence among a number of poor people, consisting of men, women, and children; to each man he gave 6*d*, to each woman 4*d*, and to each child 2*d*: moreover there were twice as many women as men, and thrice as many children as women. How many were there of each? Ans. 2 men, 4 women, and 12 children.

6. One being asked his age, said, if $\frac{3}{5}$ of the years I have lived, be multiplied by 7, and $\frac{2}{3}$ of them be added to the product, the sum will be 219. What was his age?

Ans. 45 years.

DOUBLE POSITION.

DOUBLE POSITION is the method of resolving certain questions by means of two suppositions of false numbers.

To the Double Rule of Position belong such questions as have their results not proportional to their positions : such are those, in which the numbers sought, or their parts, or their multiples, are increased or diminished by some given absolute number, which is no known part of the number sought.

RULE I.*

TAKE OR assume any two convenient numbers, and proceed with each of them separately, according to the conditions of the question, as in Single Position ; and find how much each result is different from the result mentioned in the question, calling these differences the *errors*, noting also whether the results are too great or too little.

* *Demonstr.* The Rule is founded on this supposition, namely, that the first error is to the second, as the difference between the true and first supposed number, is to the difference between the true and second supposed number ; when that is not the case, the exact answer to the question cannot be found by this Rule.—That the Rule is true, according to that supposition, may be thus proved.

Let a and b be the two suppositions, and A and B their results, produced by similar operation ; also r and s their errors, or the differences between the results A and B from the true result N ; and let x denote the number sought, answering to the true result N of the question.

Then is $N - A = r$, and $N - B = s$. And, according to the supposition on which the Rule is founded, $r : s :: x - a : x - b$: hence, by multiplying extremes and means, $rx - rb = sx - sa$; then, by transposition, $rx - sx = rb - sa$; and, by division, $x = \frac{rb - sa}{r - s}$ the number sought, which is the rule when the results are both too little.

If the results be both too great, so that A and B are both greater than N ; then $N - A = -r$, and $N - B = -s$, or r and s are both negative ; hence $-r : -s :: x - a : x - b$, but $-r : -s :: +r : +s$, therefore $r : s :: x - a : x - b$; and the rest will be exactly as in the former case.

But if one result A only be too little, and the other B too great, or one error r positive, and the other s negative, then the theorem becomes $x = \frac{rb + sa}{r + s}$, which

is the Rule in this case, or when the errors are unlike

Then multiply each of the said errors by the contrary supposition, namely, the first position by the second error, and the second position by the first error. Then,

If the errors are alike, divide the difference of the products by the difference of the errors, and the quotient will be the answer.

But if the errors are unlike, divide the sum of the products by the sum of the errors, for the answers.

Note, The errors are said to be alike, when they are either both too great or both too little ; and unlike, when one is too great and the other too little.

EXAMPLES.

1. What number is that, which being multiplied by 6, the product increased by 18, and the sum divided by 9, the quotient shall be 20 ?

Suppose the two numbers 18 and 30. Then,

	First Position.		Second Position.	Proof.
	18	Suppose	30	27
	6	mult.	6	6
	<hr style="width: 50px; margin: 0 auto;"/>		<hr style="width: 50px; margin: 0 auto;"/>	<hr style="width: 50px; margin: 0 auto;"/>
	108		108	162
	18	add	18	18
	<hr style="width: 50px; margin: 0 auto;"/>		<hr style="width: 50px; margin: 0 auto;"/>	<hr style="width: 50px; margin: 0 auto;"/>
	9) 126	div.	9) 198	9) 180
	<hr style="width: 50px; margin: 0 auto;"/>		<hr style="width: 50px; margin: 0 auto;"/>	<hr style="width: 50px; margin: 0 auto;"/>
	14	results	22	20
	20	true res.	20	<hr style="width: 50px; margin: 0 auto;"/>
	<hr style="width: 50px; margin: 0 auto;"/>		<hr style="width: 50px; margin: 0 auto;"/>	
	+6	errors unlike	-2	
2nd pos.	30	mult.	18	1st pos.
	<hr style="width: 50px; margin: 0 auto;"/>		<hr style="width: 50px; margin: 0 auto;"/>	
Er- } 2	180		36	
rors } 6	36		<hr style="width: 50px; margin: 0 auto;"/>	
sum 8)	216	sum of products.		
	<hr style="width: 50px; margin: 0 auto;"/>			
	27	Answer sought.		
	<hr style="width: 50px; margin: 0 auto;"/>			

RULE II.

FIND, by trial, two numbers, as near the true number as convenient, and work with them as in the question ; marking the errors which arise from each of them.

Multiply the difference of the two numbers assumed, or found by trial, by one of the errors, and divide the product by the difference of the errors, when they are alike, but by their sum when they are unlike.

Add

Add the quotient, last found, to the number belonging to the said error, when that number is too little, but subtract it when too great, and the result will give the true quantity sought*.

EXAMPLES.

1. So, the foregoing example, worked by this 2d rule will be as follows :

30 positions 18 ;	their dif. 12
—2 errors + 6 ;	least error 2

sum of errors 8) 24 (3 subtr.
from the position 30

leaves the answer 27

Ex. 2. A son asking his father how old he was, received this answer : Your age is now one-third of mine ; but 5 years ago, your age was only one-fourth of mine. What then are their two ages ? Ans. 15 and 45.

3. A workman was hired for 20 days, at 3s per day, for every day he worked ; but with this condition, that for every day he played, he should forfeit 1s. Now it so happened, that upon the whole he had 2l 4s to receive. How many of the days did he work ? Ans. 16.

4. A and B began to play together with equal sums of money : A first won 20 guineas, but afterwards lost back $\frac{2}{3}$ of what he then had ; after which, B had 4 times as much as A. What sum did each begin with ? Ans. 100 guineas.

5. Two persons, A and B, have both the same income, A saves $\frac{1}{5}$ of his ; but B, by spending 50l per annum more than A, at the end of 4 years finds himself 100l in debt. What does each receive and spend per annum ?

Ans. They receive 125l per annum ; also A spends 100l, and B spends 150l per annum.

* For since, by the supposition, $r : s :: x - a : x - b$, therefore by division, $r - s : s :: b - a : x - b$, which is the 2d Rule.

PERMUTATIONS AND COMBINATIONS.

PERMUTATION is the altering, changing or varying the position or order of things ; or the showing how many different ways they may be placed.—This is otherwise called Alternation, Changes, or Variation ; and the only thing to be regarded here, is the order they stand in ; for no two parcels are to have all their quantities placed in the same situation ; as, how many changes may be rung on a number of bells, or how many different ways any number of persons may be placed, or how many several variations may be made of any number of letters, or any other things proposed to be varied.

COMBINATION is the showing how often a less number of things can be taken out of a greater, and combined together, without considering their places, or the order they stand in. This is sometimes called Election or Choice ; and here every parcel must be different from all the rest, and no two are to have precisely the same quantities or things.

Combinations of the same Form, are those in which there are the same number of quantities, and the same repetitions : thus, *abc, bcda, cdba*, are of the same form ; *abc, abbb, aabb*, are of different forms.

Composition of Quantities, is the taking a given number of quantities out of as many equal rows of different quantities, one out of every row, and combining them together.

Illustrations of these definitions are in the following Problems :

PROBLEM I.

To assign the Number of Permutations, or Changes, that can be made of any Given Number of Things, all different from each other.

RULE*.

MULTIPLY all the terms of the natural series of numbers, from 1 up to the given number, continually together, and the last product will be the answer required.

EXAMPLES.

* The reason of the Rule may be shown thus ; any one thing *a* is capable only of one position, as *a*.

Any two things *a* and *b*, are only capable of two variations ; as *ab, ba* ; whose number is expressed by 1×2 .

EXAMPLES.

1. How many changes may be rung on 6 bells.

$$\begin{array}{r}
 1 \\
 2 \\
 \hline
 2 \\
 3 \\
 \hline
 6 \\
 4 \\
 \hline
 24 \\
 5 \\
 \hline
 120 \\
 6 \\
 \hline
 720 \text{ the Answer.}
 \end{array}$$

Or $1 \times 2 \times 3 \times 4 \times 5 \times 6 = 720$ the Answer.

2. How many days can 7 persons be placed in a different position at dinner? Ans. 5040 days.

3. How many changes may be rung on 12 bells, and what time would it require, supposing 10 changes to be rung in 1 minute, and the year to consist of 365 days, 5 hours, and 49 minutes?

Ans. 479001600 changes, and 91 years, 26 days, 22 hours, 41 minutes.

4. How many changes may be made of the words in the following verse: *Tot tibi sunt dotes, virgo, quot sidera cælo?*

Ans. 40320 changes.

If there be three things, *a*, *b*, and *c*; then any two of them, leaving out the 3d, will have 1×2 variations; and consequently when the 3d is taken in, there will be $1 \times 2 \times 3$ variations.

In the same manner, when there are 4 things, every three, leaving out the 4th, will have $1 \times 2 \times 3$ variations; consequently by taking in successively the 4 left out, there will be $1 \times 2 \times 3 \times 4$ variations. And so on as far as we please.

PROBLEM II.

Any Number of different things being given; to find how many Changes can be made out of them, by taking a Given Number of Quantities at a Time.

RULE.*

TAKE a series of numbers, beginning at the number of things given, and decreasing by 1 to the number of quantities to be taken at a time, and the product of all the terms will be the answer required.

EXAMPLES.

1. How many changes may be rung with 3 bells out of 8?

$$\begin{array}{r} 8 \\ 7 \\ \hline 56 \\ 6 \\ \hline \end{array}$$

336 the answer.

Or, $8 \times 7 \times 6 (= 3 \text{ terms}) = 336$ the Answer.

2. How many words can be made with 5 letters of the alphabet, supposing 24 letters in all, and that a number of consonants alone will make a word. Ans. 5100480.

3. How many words can be made with 5 letters of the alphabet in each word, there being 26 letters in all, and 6 vowels, admitting that a number of consonants alone will not make a word? Ans. 137858400.

PROB-

* This Rule, expressed in algebraic terms, is as follows;

$m \times m-1 \times m-2 \times m-3$ &c. to n terms: where $m =$ the number of things given, and $n =$ the quantities to be taken at a time.

In order to demonstrate the Rule, it will be proper to premise the following Lemma;

LEMMA. The number of changes of m things, taken n at a time, is equal to m changes of $m-1$ things, taken $n-1$ at a time.

Demonstr. Let any five quantities $a b c d e$ be given.

First, leave out the a , and let $v =$ the number of all the variations of every two, bc, bd , &c. that can be taken out of the four remaining quantities $b c d e$.

Now, let a be put in the first place of each of them, a, b, c, a, b, d , &c. and the number of changes which still remain the same; that is, $v =$ the number of variations of every 3 out of the 5, a, b, c, d, e , when a is first.

fin

PROBLEM III.

Any Number of Things being given; of which there are several given Things of one Sort, and several of another, &c.; to Find how many Changes can be made out of them all.

RULE.*

TAKE the series $1 \times 2 \times 3 \times 4$, &c. up to the number of things given, and find the product of all the terms.

Take the series $1 \times 2 \times 3 \times 4$, &c. up to the number of given things of the first sort, and the series $1 \times 2 \times 3 \times 4$, &c. up to the number of given things of the second sort, &c.

Divide

In like manner, if b, c, d, e be successively left out, the number of variations of all the two's will also be $=v$; and putting b, c, d, e respectively in the first place, to make 3 quantities out of 5, there will still be v variations as before.

But these are all the variations that can happen of 3 things out of 5, when a, b, c, d, e , are successively put first; and therefore the sum of all these is the sum of all the changes of 3 things out of 5.

But the sum of these is so many times v as is the number of things; that is $5v$, or mv , = all the changes of 3 times out of 5.

And the same way of reasoning may be applied to any numbers whatever.

Demon. of the Rule. Let any 7 things, $a b c d e f g$, be given, and let 3 be the number of quantities to be taken.

Then $m = 7$, and $n = 3$.

Now, it is evident, that the number of changes that can be made by taking 1 by 1 out of 5 things, will be 5, which let $= v$.

Then, by the Lemma, when $m = 6$, and $n = 2$, the number of changes will be $= mv = 6 \times 5$; which let be $= v$ a second time.

Again, by the Lemma, when $m = 7$ and $n = 3$, the number of changes is $mv = 7 \times 6 \times 5$; that is $mv = m \times (m-1) \times (m-2)$, continued to 3, or n terms.

And the same may be shown for any other numbers.

* This Rule is expressed in terms thus:

$$1 \times 2 \times 3 \times 4 \times 5, \text{ \&c. to } m$$

$$1 \times 2 \times 3, \text{ \&c. to } p \times 1 \times 2 \times 3, \text{ \&c. to } q, \text{ \&c.}$$

where m = the number of things given, p = the number of things of the first sort, q = the number of things of the second sort, &c.

The Demonstration may be shown as follows;

Any two quantities, a, b , both different, admit of 2 changes; but if the quantities are the same, or $a b$ becomes $a a$, there will be only one position; which

may be expressed by $\frac{1 \times 2}{1 \times 2} = 1$.

Ans

Divide the product of all the terms of the first series by the joint product of all the terms of the remaining ones, and the quotient will be the answer required.

EXAMPLES.

1. How many variations can be made of the letters in the word Bacchanalia?

$$\begin{aligned}
 & 1 \times 2 \quad (\text{= number of } c\text{'s}) = 2 \\
 & 1 \times 2 \times 3 \times 4 \quad (\text{= number of } a\text{'s}) = 24 \\
 & 1 \times 2 \times 3 \times 4 \times 5 \times 6 \times 7 \times 8 \times 9 \times 10 \times 11 \\
 & \quad (\text{= number of letters in the word}) = 39916800 \\
 & 2 \times 24 = 48 \quad 39916800 \quad (831600 \text{ the Answer.}
 \end{aligned}$$

151

76

288

2. How many different numbers can be made of the following figures, 1220005555? Ans. 12600.

3. How many varieties will take place in the succession of the following musical notes, fa, fa, fa, fa, sol, sol, la, mi? Ans. 840.

Any 3 quantities, a, b, c , all different from each other, afford 6 variations; but if the quantities be all alike, or $a b c$ becomes $a a a$, then the 6 variations will be

reduced to 1; which may be expressed by $\frac{1 \times 2 \times 3}{1 \times 2 \times 3} = 1$. Again, if two of the quantities only are alike, or $a b c$ becomes $a a c$; then the 6 variations will be reduced to these 3, $a a c, c a a$, and $a c a$; which may be expressed by $\frac{1 \times 2 \times 3}{1 \times 2} = 3$.

Any 4 quantities, $a b c d$, all different from each other, will admit of 24 variations. But if the quantities be the same, or $a b c d$ becomes $a a a a$, the number of variations will be reduced to one; which is $= \frac{1 \times 2 \times 3 \times 4}{1 \times 2 \times 3 \times 4} = 1$.

Again, if three of the quantities only be the same, or $a b c d$ becomes $a a a b$, the number of variations will be reduced to these 4, $a a a b, a a b a, a b a a$, and $b a a a$; which is $= \frac{1 \times 2 \times 3 \times 4}{1 \times 2 \times 3} = 4$.

And thus it may be shown, that if two of the quantities be alike, or the 4 quantities be $a a b c$, the number of variations will be reduced to 12; which may be expressed by $\frac{1 \times 2 \times 3 \times 4}{1 \times 2} = 12$.

And by reasoning in the same manner, it will appear, that the number of changes which can be made of the quantities $a b b c c$, is equal to 60; which may be expressed by $\frac{1 \times 2 \times 3 \times 4 \times 5 \times 6}{1 \times 2 \times 1 \times 2 \times 3} = 60$. And so on for any other quantities whatever

PROBLEM IV.

To find the Changes of any Given Number of Things, taking a Given Number at a Time: in which there are several Given Things of one Sort, several of another, &c.

RULE.*

FIND all the different forms of combination of all the given things, taken as many at a time as in the question.

Find the number of changes in any form, and multiply it by the number of combinations in that form.

Do the same for every distinct form, and the sum of all the products will give the whole number of changes required.

EXAMPLES.

1. How many alterations, or changes, can be made of every four letters out of these 8, *aaabbbcc*?

No. of forms.	No. of changes.
<i>a³b, a²c, b³a, b³c</i> - - - - -	4
<i>a²b², a²c², b²c²</i> - - - - -	6
<i>a²bc, b²ac, c²ab</i> - - - - -	12

$$\text{Therefore } \left\{ \begin{array}{l} 4 \times 4 = 16 \\ 3 \times 6 = 18 \\ 3 \times 12 = 36 \end{array} \right.$$

70 = number of changes
— required.

2. How many changes can be made of every 8 letters out of these 10; *aaaabbbccde*? Ans. 22260.

3. How many different numbers can be made out of 1 unit,

* The reason of this Rule is plain from what has been shown before, and the nature of the problem.

A Rule for finding the Number of Forms.

1. PLACE the things so, that the greatest indices may be first, and the rest in order.

2. Begin with the first letter, and join it to the second, third, fourth, &c. to the last.

3. Then take the second letter, and join it to the third, fourth, &c. to the last. And so on, till they are entirely exhausted, always remembering to reject such combinations as have occurred before; and this will give the combinations of all the two's.

4. Join the first letter to every one of the twos, and the second, third, &c. as before; and it will give the combinations of all the threes.

5. Proceed in the same manner to get the combinations of all the fours, &c. and you will at last get all the several forms of combinations, and the number in each form.

2 twos, 3 threes, 4 fours, and 5 fives; taken 5 at a time?
 Ans. 2111.

PROBLEM V.

To find the Number of Combinations of any Given Number of things all different from each other, taken any Given Number at a time.

RULE.*

TAKE the series 1, 2, 3, 4, &c. up to the number to be taken at a time, and find the product of all the terms.

Take a series of as many terms, decreasing by 1, from the given number, out of which the election is to be made, and find the product of all the terms.

Divide the last product by the former, and the quotient will be the number sought.

EXAMPLES.

1. How many combinations can be made of 6 letters out of ten?

* This Rule, expressed algebraically, is,

$$\frac{m}{1} \times \frac{m-1}{2} \times \frac{m-2}{3} \times \frac{m-3}{4} \text{ \&c. to } n \text{ terms; where } m \text{ is the number of given quantities, and } n \text{ those to be taken at a time.}$$

Demonstr. of the Rule. 1. Let the number of things to be taken at a time be 2, and the things to be combined = m .

Now, when m , or the number of things to be combined, is only two, as a and b , it is evident that there can be but one combination, as ab ; but if m be increased by one, or the letters to be combined be 3, as a, b, c ; then it is plain that the number of combinations will be increased by 2, since with each of the former letters a and b , the new letter c may be joined. In this case therefore, it is evident that the whole number of combinations will be truly expressed by $1 + 2$.

Again, if m be increased by one letter more, or the whole number of letters be four, as a, b, c, d ; then it will appear that the whole number of combinations must be increased by 3, since with each of the preceding letters the new letter d may be combined. The combinations, therefore, in this case will be truly expressed by $1 + 2 + 3$.

And in the same manner it may be shown that the whole number of combinations of 2, in 5 things, will be $1 + 2 + 3 + 4$; of 2 in 6 things, $1 + 2 + 3 + 4 + 5$; and of 2, in 7 things, $1 + 2 + 3 + 4 + 5 + 6$, &c.; whence, universally, the number of combinations of m things, taken 2 by 2, is = $1 + 2 + 3 + 4 + 5 + 6$, &c. to $(m-1)$ terms.

But the sum of this series is = $\frac{m}{1} \times \frac{m-1}{2}$; which is the same as the rule.

2. Let now the number of quantities in each combination be supposed to be three.

Then

$1 \times 2 \times 3 \times 4 \times 5 \times 6$ (= the number to be taken at a time) = 720.

$10 \times 9 \times 8 \times 7 \times 6 \times 5$ (= same number from 10) = 151200.

Then 720) 151200 (210 the Answer.

1440

720

720

2. How many combinations can be made of 2 letters out of the 24 letters of the alphabet? Ans. 276.

3. A general, who had often been successful in war, was asked by his king what reward he should confer upon him for his services; the general only desired a farthing for every file, of 10 men in a file, which he could make with a body of 100 men; what is the amount in pounds sterling?

Ans. 18031572350l 9s 2d.

Then it is plain, that when $m = 3$, or the things to be combined are a, b, c , there can be only one combination. But if m be increased by 1, or the things to be combined are 4, as a, b, c, d , then will the number of combinations be increased by 3: since 3 is the number of combinations of 2 in all the preceding letters, a, b, c , and with each two of these the new letter d may be combined.

The number of combinations, therefore in this case, is $1 + 3$.

Again, if m be increased by one more, or the number of letters be supposed 5; then the former number of combinations will be increased by 6, that is, by all the combinations of 2 in the 4 preceding letters, a, b, c, d ; since, as before, with each two of these the new letter e may be combined.

The number of combinations, therefore, in this case, is $1 + 3 + 6$.

Whence, universally, the number of combinations of m things, taken 3 by 3, is $1 + 3 + 6 + 10$, &c. to $m - 2$ terms.

But the sum of this series is $\frac{m}{1} \times \frac{m-1}{2} \times \frac{m-2}{3}$; which is the same as the

rule.

And the same thing will hold, let the number of things to be taken at a time be what it will; therefore the number of combinations of m things, taken n at a time, will be =

$\frac{m}{1} \times \frac{m-1}{2} \times \frac{m-2}{3} \times \frac{m-3}{4}$, &c. to n terms. Q. E. D.

PROBLEM VI.

To find the Number of Combinations of any Given Number of Things, by taking any Given Number at a time; in which there are several Things of one Sort, several of another, &c.

RULE.

FIND, by trial, the number of different forms which the things to be taken at a time will admit of, and the number of combinations there are in each.

Add all the combinations, thus found together, and the sum will be the number required.

EXAMPLES.

1. Let the things proposed be $a a a b b c$; it is required to find the number of combinations made of every 3 of these quantities?

Forms.	Combinations.
a^3 - - - - - -	1
a^2b, a^2c, b^2a, b^2c - - - -	4
$a b c$ - - - - - -	1

Number of combinations required = 6

2. Let $a a a b b b c c$ be proposed; it is required to find the number of combinations of these quantities, taken 4 at a time? Ans. 10.

3. How many combinations are there in $a a a a b b c c d e$, taking 8 at a time? Ans. 13.

4. How many combinations are there in $a a a a a b b b b b c c c c d d d e e e e f f f g$, taking 10 at a time? Ans. 2319.

PROBLEM VII.

To find the Compositions of any Number, in an equal Number of Sets, the Things themselves being all different.

RULE*.

MULTIPLY the number of things in every set continually together, and the product will be the answer required.

* *Demonstr.* Suppose there are only two sets; then, it is plain, that every quantity of the one set being combined with every quantity of the other, will make all the compositions, of two things in these two sets; and the number of these compositions

EXAMPLES.

1. Suppose there are four companies, in each of which there are 9 men ; it is required to find how many ways 4 men may be chosen, one out of each company ?

$$\begin{array}{r}
 9 \\
 9 \\
 \hline
 81 \\
 9 \\
 \hline
 729 \\
 9 \\
 \hline
 6561 \text{ the Answer.} \\
 \hline
 \hline
 \end{array}$$

Or, $9 \times 9 \times 9 \times 9 = 6561$ the Answer.

2. Suppose there are 4 companies ; in one of which there are 6 men, in another 8, and in each of the other two 9 ; what are the choices, by a composition of 4 men, one out of each company ?

Ans. 3888.

3. How many changes are there in throwing 5 dice ?

Ans. 7776.

compositions is evidently the product of the number of quantities in one set by that in the other.

Again, suppose there are three sets ; then the composition of two, in any two of the sets, being combined with every quantity of the third, will make all the compositions of three in the three sets. That is, the compositions of two in any two of the sets, being multiplied by the number of quantities in the remaining set, will produce the compositions of three in the three sets ; which is evidently the continual product of all the three numbers in the three sets.

And the same manner of reasoning will hold, let the number of sets be what it will. Q. E. D.

The doctrine of permutations, combinations, &c. is of very extensive use in different parts of the Mathematics ; particularly in the calculation of annuities and chances. The subject might have been pursued to a much greater length ; but what is here done, will be found sufficient for most of the purposes to which things of this nature are applicable.

PRACTICAL QUESTIONS IN ARITHMETIC.

QUEST. 1. The swiftest velocity of a cannon-ball, is about 2000 feet in a second of time. Then in what time, at that rate, would such a ball be in moving from the earth to the sun, admitting the distance to be 100 millions of miles, and the year to contain 365 days 6 hours.

Ans. $8\frac{4}{7}\frac{8}{31}\frac{0}{49}$ years.

QUEST. 2. What is the ratio of the velocity of light to that of a cannon-ball, which issues from the gun with a velocity of 1500 feet per second; light passing from the sun to the earth in $7\frac{1}{2}$ minutes?

Ans. the ratio of $782222\frac{2}{5}$ to 1.

QUEST. 3. The slow or parade-step being 70 paces per minute, at 28 inches each pace, it is required to determine at what rate per hour that movement is?

Ans. $1\frac{1}{3}\frac{2}{3}$ miles.

QUEST. 4. The quick-time or step, in marching, being 2 paces per second, or 120 per minute, at 28 inches each; then at what rate per hour does a troop march on a route, and how long will they be in arriving at a garrison 20 miles distant, allowing a halt of one hour by the way to refresh?

Ans. $\left\{ \begin{array}{l} \text{the rate is } 3\frac{2}{3} \text{ miles an hour.} \\ \text{and the time } 7\frac{2}{3} \text{ hr. or } 7 \text{ h. } 17\frac{1}{2} \text{ min.} \end{array} \right.$

QUEST. 5. A wall was to be built 700 yards long in 29 days. Now, after 12 men had been employed on it for 11 days, it was found that they had completed only 220 yards of the wall. It is required then to determine how many men must be added to the former, that the whole number of them may just finish the wall in the time proposed, at the same rate of working?

Ans. 4 men to be added.

QUEST. 6. To determine how far 500 millions of guineas will reach, when laid down in a straight line touching one another; supposing each guinea to be an inch in diameter, as it is very nearly.

Ans. 7891 miles, 728 yards, 2 ft. 8 in.

QUEST. 7. Two persons, A and B, being on opposite sides of a wood, which is 536 yards about, they being to go round it, both the same way, at the same instant of time; A goes at the rate of 11 yards per minute, and B 34 yards in 3 minutes; and the question is, how many times will the wood be gone round before the quicker overtake the slower?

Ans. 17 times.

QUEST-

QUEST. 8. A can do a piece of work alone in 12 days, and B alone in 14; in what time will they both together perform a like quantity of work? Ans. $6\frac{6}{13}$ days.

QUEST. 9. A person who was possessed of a $\frac{3}{8}$ share of a copper mine, sold $\frac{3}{4}$ of his interest in it for 1800*l*; what was the reputed value of the whole at the same rate? Ans. 4000*l*.

QUEST. 10. A person after spending 20*l* more than $\frac{1}{4}$ of his yearly income, had then remaining 30*l* more than the half of it; what was his income? Ans. 200*l*.

QUEST. 11. The hour and minute hand of a clock are exactly together at 12 o'clock; when are they next together? Ans. $1\frac{1}{11}$ hr, or $1\text{ hr } \frac{5}{11}$ min.

QUEST. 12. If a gentleman whose annual income is 1500*l*, spends 20 guineas a week; whether will he save or run in debt, and how much in the year? Ans. save 408*l*.

QUEST. 13. A person bought 180 oranges at 2 a penny, and 180 more at 3 a penny; after which, selling them out again at 5 for 2 pence, whether did he gain or lose by the bargain? Ans. he lost 6 pence.

QUEST. 14. If a quantity of provisions serves 1500 men 12 weeks, at the rate of 20 ounces a day for each man; how many men will the same provisions maintain for 20 weeks, at the rate of 8 ounces a day for each man? Ans. 2250 men.

QUEST. 15. In the latitude of London, the distance round the earth measured on the parallel of latitude, is about 15550 miles; now as the earth turns round in 23 hours 56 minutes, at what rate per hour is the city of London carried by this motion from west to east? Ans. $649\frac{2}{3}\frac{9}{10}$ miles an hour.

QUEST. 16. A father left his son a fortune, $\frac{1}{4}$ of which he ran through in 8 months; $\frac{3}{7}$ of the remainder lasted him 12 months longer; after which he had bare 820*l* left. What sum did the father bequeath his son? Ans. 1913*l* 6*s* 8*d*.

QUEST. 17. If 1000 men, besieged in a town with provisions for 5 weeks, allowing each man 16 ounces a day, be reinforced with 500 men more; and supposing that they cannot be relieved till the end of 8 weeks, how many ounces a day must each man have, that the provision may last that time? Ans. $6\frac{2}{3}$ ounces.

QUEST. 18. A younger brother received 8400*l*, which was just $\frac{7}{9}$ of his elder brother's fortune: What was the father worth at his death? Ans. 19200*l*.

QUEST.

QUEST. 19. A person, looking on his watch, was asked what was the time of the day, who answered, It is between 5 and 6 ; but a more particular answer being required, he said that the hour and minute hands were then exactly together ; What was the time ?

Ans. $27 \frac{3}{11}$ min. past 5.

QUEST. 20. If 20 men can perform a piece of work in 12 days, how many men will accomplish another thrice as large in one-fifth of the time ?

Ans. 300.

QUEST. 21. A father devised $\frac{7}{8}$ of his estate to one of his sons, and $\frac{7}{8}$ of the residue to another, and the surplus to his relict for life. The children's legacies were found to be 514*l* 6*s* 8*d* different : Then what money did he leave the widow the use of ?

Ans. 1270*l* 1*s* 9 $\frac{1}{8}$ *d*.

QUEST. 22. A person, making his will, gave to one child $\frac{1}{2}$ of his estate, and the rest to another. When these legacies came to be paid the one turned out 1200*l* more than the other : What did the testator die worth ?

Ans. 4000*l*.

QUEST. 23. Two persons, A and B, travel between London and Lincoln, distant 100 miles, A from London, and B from Lincoln, at the same instant. After 7 hours they meet on the road when it appeared that A had rode $1\frac{1}{2}$ miles an hour more than B. At what rate per hour then did each of the travellers ride ?

Ans. A $7\frac{2}{3}$, and B $6\frac{1}{3}$ miles.

QUEST. 24. Two persons, A and B, travel between London and Exeter. A leaves Exeter at 8 o'clock in the morning, and walks at the rate of 3 miles an hour, without intermission : and B sets out from London at 4 o'clock the same evening, and walks for Exeter at the rate of 4 miles an hour constantly. Now, supposing the distance between the two cities to be 130 miles, whereabouts on the road will they meet ?

Ans. $69\frac{3}{4}$ miles from Exeter.

QUEST. 25. One hundred eggs being placed on the ground in a straight line, at the distance of a yard from each other : How far will a person travel who shall bring them one by one to a basket, which is placed at one yard from the first egg ?

Ans. 10100 yards, or 5 miles and 1300 yds.

QUEST. 26. The clocks of Italy go on to 24 hours ; Then how many strokes do they strike in one complete revolution of the index ?

Ans. 300.

QUEST. 27. One Sessa, an Indian, having invented the game of chess, showed it to his prince, who was so delighted

with

with it, that he promised him any reward he should ask ; on which Sessa requested that he might be allowed one grain of wheat for the first square on the chess board, 2 for the second, 4 for the third, and so on, doubling continually, to 64 the whole number of squares. Now, supposing a pint to contain 7680 of these grains, and one quarter or 8 bushels to be worth 27s 6d, it is required to compute the value of all the corn ?

Ans. 6450468216285l 17s 3d $\frac{32757}{2768}q$.

QUEST. 28. A person increased his estate annually by 100l more than the $\frac{1}{4}$ part of it ; and at the end of 4 years found that his estate amounted to 10342l 3s 9d. What had he at first ?

Ans. 4000l.

QUEST. 29. Paid 1012l 10s for a principal of 750l, taken in 7 years before ; at what rate per cent. per annum did I pay interest ?

Ans. 5 per cent.

QUEST. 30. Divide 1000l among A, B, C ; so as to give A 120 more, and B 95 less than C.

Ans. A 445, B 230, C 325.

QUEST. 31. A person being asked the hour of the day, said, the time past noon is equal to $\frac{4}{5}$ ths of the time till midnight. What was the time ?

Ans. 20 min. past 5.

QUEST. 32. Suppose that I have $\frac{3}{16}$ of a ship worth 1200l ; what part of her have I left after selling $\frac{2}{5}$ of $\frac{4}{9}$ of my share, and what is it worth ?

Ans. $\frac{37}{246}$, worth 185l.

QUEST. 33. Part 1200 acres of land among A, B, C ; so that B may have 100 more than A, and C 64 more than B.

Ans. A 312, B 412, C 476.

QUEST. 34. What number is that, from which if there be taken $\frac{2}{7}$ of $\frac{3}{8}$, and to the remainder be added $\frac{3}{15}$ of $\frac{5}{16}$, the sum will be 10 ?

Ans. $9\frac{7}{24}$.

QUEST. 35. There is a number which if multiplied by $\frac{2}{3}$ of $\frac{4}{5}$ of $1\frac{1}{2}$, will produce 1 : what is the square of that number ?

Ans. $1\frac{9}{16}$.

QUEST. 36. What length must be cut off a board, $8\frac{1}{2}$ inches broad to contain a square foot, or as much as 12 inches in length and 12 in breadth ?

Ans. $16\frac{16}{17}$ inches.

QUEST. 37. What sum of money will amount to 138l 2s 6d, in 15 months, at 5 per cent. per annum simple interest ?

Ans. 130l.

QUEST. 38. A father divided his fortune among his three sons, A, B, C, giving A 4 as often as B 3, and C 5 as often as

B 6 ;

B 6; what was the whole legacy, supposing A's share was 4000*l*. Ans. 9500*l*.

QUEST. 39. A young hare starts 40 yards before a greyhound, and is not perceived by him till she has been up 40 seconds; she scuds away at the rate of 10 miles an hour, and the dog, on view, makes after her at the rate of 18: how long will the course hold, and what ground will be run over, counting from the outseting of the dog?

Ans. $60\frac{5}{2}$ sec. and 530 yards run.

QUEST. 40. Two young gentlemen, without private fortune, obtain commissions at the same time, and at the age of 18. One thoughtlessly spends 10*l* a year more than his pay; but shocked at the idea of not paying his debts, gives his creditor a bond for the money, at the end of every year, and also insures his life for the amount; each bond costs him 30 shillings, besides the lawful interest of 5 per cent. and to insure his life costs him 6 per cent.

The other, having a proper pride, is determined never to run in debt; and, that he may assist a friend in need, perseveres in saving 10*l* every year, for which he obtains an interest of 5 per cent. which interest is every year added to his savings, and laid out, so as to answer the effect of compound interest.

Suppose these two officers to meet at the age of 50, when each receives from Government 400*l* per annum; that the one, seeing his past errors, is resolved in future to spend no more than he actually has, after paying the interest for what he owes, and the insurance on his life.

The other, having now something before hand, means in future, to spend his full income, without increasing his stock.

It is desirable to know how much each has to spend per annum, and what money the latter has by him to assist the distressed, or leave to those who deserve it?

Ans. The reformed officer has to spend 66*l* 19*s* $1\frac{3}{4}$ ·5389*d* per annum.

The prudent officer has to spend 437*l* 12*s* $11\frac{3}{4}$ ·4379*d* per annum.

And the latter has saved, to dispose of, 752*l* 19*s* 9·1396*d*.

OF LOGARITHMS.*

LOGARITHMS are made to facilitate troublesome calculations in numbers. This they do, because they perform multiplication by only addition, and division by only subtraction, and raising of powers by multiplying the logarithm by the index of the power, and extracting of roots by dividing the logarithm of the number by the index of the root. For, logarithms are numbers so contrived, and adapted to other numbers, that the sums and differences of the former shall correspond to, and show, the products and quotients of the latter, &c.

Or, more generally, logarithms are the numerical exponents of ratios ; or they are a series of numbers in arithmetical

* The invention of Logarithms is due to Lord Napier, Baron of Merchiston, in Scotland, and is properly considered as one of the most useful inventions of modern times. A table of these numbers was first published by the inventor at Edinburgh, in the year 1614, in a treatise entitled *Canon Mirificum Logarithmorum* ; which was eagerly received by all the learned throughout Europe. Mr. Henry Briggs, then professor of geometry at Gresham College, soon after the discovery, went to visit the noble inventor ; after which, they jointly undertook the arduous task of computing new tables on this subject, and reducing them to a more convenient form than that which was at first thought of. But Lord Napier dying soon after, the whole burden fell upon Mr. Briggs, who, with prodigious labour and great skill, made an entire Canon, according to the new form, for all numbers from 1 to 20000, and from 90000 to 10100, to 14 places of figures, and published it at London, in the year 1624, in a treatise entitled *Arithmetica Logarithmica*, with directions for supplying the intermediate parts.

tical progression, answering to another series of numbers in geometrical progression.

Thus, $\left\{ \begin{array}{l} 0, 1, 2, 3, 4, 5, 6, \text{Indices, or logarithms.} \\ 1, 2, 4, 8, 16, 32, 64, \text{Geometric progression.} \end{array} \right.$

Or $\left\{ \begin{array}{l} 0, 1, 2, 3, 4, 5, 6, \text{Indices, or logarithms.} \\ 1, 3, 9, 27, 81, 243, 729, \text{Geometric progression.} \end{array} \right.$

Or $\left\{ \begin{array}{l} 0, 1, 2, 3, 4, 5, \text{Indices, or logs.} \\ 1, 10, 100, 1000, 10000, 100000, \text{Geom. progress.} \end{array} \right.$

Where it is evident, that the same indices serve equally for any geometric series ; and consequently there may be an

This Canon was again published in Holland by Adrain Vlacq, in the year 1628, together with the Logarithms of all the numbers which Mr. Briggs had omitted ; but he contracted them down to 10 places of decimals. Mr. Briggs also computed the Logarithms of the sines, tangents, and secants, to every degree, and centesim, or 100th part of a degree, of the whole quadrant ; and annexed them to the natural sines, tangents, and secants, which he had before computed, to fifteen places of figures. These Tables, with their construction and use, were first published in the year 1633, after Mr. Briggs's death, by Mr. Henry Gellibrand, under the title of *Trigonometria Britannica*.

Benjamin Ursinus also gave a Table of Napier's Logs. and of sines, to every 10 seconds. And Chr. Wolf, in his *Mathematical Lexicon*, says that one Van Loser had computed them to every single second, but his untimely death prevented their publication. Many other authors have treated on this subject ; but as their numbers are frequently inaccurate and incommodiously disposed, they are now generally neglected. The Tables in most repute at present, are those of Gardiner in 4to, first published in the year 1742 ; and my own Tables in 8vo, first printed in the year 1785, where the Logarithms of all numbers may be easily found from 1 to 10000000 ; and those of the sines, tangents, and secants, to any degree of accuracy required.

Also Mr. Michael Taylor's Tables in large 4to, containing the common logarithms, and the logarithmic sines and tangents to every second of the quadrant. And, in France, the new book of logarithms by Callet ; the 2d edition of which, in 1795, has the tables still farther extended, and are printed with what are called stereotypes, the types in each page being soldered together into a solid mass or block.

Dodson's Antilogarithmic Canon is likewise a very elaborate work, and used for finding the numbers answering to any given logarithm.

endless

endless variety of systems of logarithms, to the same common numbers, by only changing the second term, 2, 3, or 10, &c. of the geometrical series of whole numbers; and by interpolation the whole system of numbers may be made to enter the geometric series, and receive their proportional logarithms, whether integers or decimals.

It is also apparent, from the nature of these series, that if any two indices be added together, their sum will be the index of that number which is equal to the product of the two terms, in the geometric progression, to which those indices belong. Thus, the indices 2 and 3 being added together, make 5; and the numbers 4 and 8, or the terms corresponding to those indices, being multiplied together, make 32, which is the number answering to the index 5.

In like manner, if any one index be subtracted from another, the difference will be the index of that number which is equal to the quotient of the two terms to which those indices belong. Thus, the index 6, minus the index 4, is = 2; and the terms corresponding to those indices are 64 and 16, whose quotient is = 4, which is the number answering to the index 2.

For the same reason, if the logarithm of any number be multiplied by the index of its power, the product will be equal to the logarithm of that power. Thus, the index or logarithm of 4 in the above series, is 2; and if this number be multiplied by 3, the product will be = 6; which is the logarithm of 64, or the third power of 4.

And, if the logarithm of any number be divided by the index of its root, the quotient will be equal to the logarithm of that root. Thus, the index or logarithm of 64 is 6; and if this number be divided by 2, the quotient will be = 3; which is the logarithm of 8, or the square root of 64.

The logarithms most convenient for practice, are such as are adapted to a geometric series increasing in a tenfold proportion, as in the last of the above forms; and are those which are to be found, at present, in most of the common tables on this subject. The distinguishing mark of this system of logarithms is, that the index or logarithm of 10 is 1; that of 100 is 2; that of 1000 is 3; &c. And, in decimals, the

the logarithm of $\cdot 1$ is -1 ; that of $\cdot 01$ is -2 ; that of $\cdot 001$ is -3 ; &c. The log. of 1 being 0 in every system. Whence it follows, that the logarithm of any number between 1 and 10, must be 0 and some fractional parts; and that of a number between 10 and 100, will be 1 and some fractional parts; and so on, for any other number whatever. And since the integral part of a logarithm, usually called the Index, or Characteristic, is always thus readily found, it is commonly omitted in the tables; being left to be supplied by the operator himself, as occasion requires.

Another Definition of Logarithms is, that the logarithm of any number is the index of that power of some other number, which is equal to the given number. So, if there be $N = r^n$, then n is the log. of N ; where n may be either positive or negative, or nothing, and the root r any number whatever, according to the different systems of logarithms. When n is $= 0$, then N is $= 1$, whatever the value of r is; which shows that the log. of 1 is always 0, in every system of logarithms. When n is $= 1$, then N is $= r$; so that the radix r is always that number whose log is 1, in every system. When the radix r is $= 2.718281828459$, &c. the indices n are the hyperbolic or Napier's log. of the numbers N ; so that n is always the hyp. log. of the number N or $(2.718 \text{ \&c.})^n$.

But when the radix r is $= 10$, then the index n becomes the common or Brigg's log. of the number N : so that the common log. of any number 10^n or N , is n the index of that power of 10 which is equal to the said number. Thus 100, being the second power of 10, will have 2 for its logarithm; and 1000, being the third power of 10, will have 3 for its logarithm: hence also, if 50 be $= 10^{1.69897}$, then is 1.69897 the common log. of 50. And, in general, the following decuple series of terms, viz. $10^4, 10^3, 10^2, 10^1, 10^0, 10^{-1}, 10^{-2}, 10^{-3}, 10^{-4}$, or 10000, 1000, 100, 10, 1, $\cdot 1, \cdot 01, \cdot 001, \cdot 0001$, have 4, 3, 2, 1, 0, $-1, -2, -3, -4$, for their logarithms, respectively. And from this scale of numbers and logarithms, the same properties easily follow, as above mentioned.

PROBLEM.

To compute the Logarithm to any of the Natural Numbers
1, 2, 3, 4, 5, &c.

RULE I.*

TAKE the geometric series, 1, 10, 100, 1000, 10000, &c. and apply to it the arithmetic series, 0, 1, 2, 3, 4, &c. as logarithms.—Find a geometric mean between 1 and 10, or between 10 and 100, or any other two adjacent terms of the series, between which the number proposed lies.—In like manner, between the mean, thus found, and the nearest extreme, find another geometrical mean; and so on, till you arrive within the proposed limit of the number whose logarithm is sought.—Find also as many arithmetical means, in the same order as you found the geometrical ones, and these will be the logarithms answering to the said geometrical means.

EXAMPLE.

Let it be required to find the logarithm of 9.

Here the proposed number lies between 1 and 10.

First, then, the log of 10 is 1, and the log of 1 is 0;

theref. $\frac{1+0}{2} \div 2 = \frac{1}{2} = \cdot 5$ is the arithmetical mean.

and $\sqrt{10 \times 1} = \sqrt{10} = 3\cdot 1622777$ the geom. mean.

hence the log. of $3\cdot 1622777$ is $\cdot 5$.

Secondly, the log. of 10 is 1, and the log. of $3\cdot 1622777$ is $\cdot 5$;

theref. $\frac{1+5}{2} \div 2 = \cdot 75$ is the arithmetical mean,

and $\sqrt{10 \times 3\cdot 1622777} = 5\cdot 6234132$ is the geom. mean;

hence the log. of $5\cdot 6234132$ is $\cdot 75$.

Thirdly, the log. of 10 is 1, and the log. of $5\cdot 6234132$ is $\cdot 75$;

theref. $\frac{1+\cdot 75}{2} \div 2 = \cdot 875$ is the arithmetical mean,

and $\sqrt{10 \times 5\cdot 6235132} = 7\cdot 4989422$ the geom. mean;

hence the log. of $7\cdot 4989422$ is $\cdot 875$.

Fourthly, the log. of 10 is 1, and the log. of $7\cdot 4989422$ is $\cdot 875$;

theref. $\frac{1+\cdot 875}{2} \div 2 = \cdot 9375$ is the arithmetical mean,

and $\sqrt{10 \times 7\cdot 4989422} = 8\cdot 6596431$ the geom. mean;

hence the log. of $8\cdot 6596431$ is $\cdot 9375$.

* The reader who wishes to inform himself more particularly concerning the history, nature, and construction of Logarithms, may consult the Introduction to my mathematical Tables, lately published, where he will find his curiosity amply gratified.

Fifthly,

Fifthly, the $\log.$ of 10 is 1 and the $\log.$ of 8.6596431 is .9875 ;
 theref. $1 + .9375 \div 2 = .96875$ is the arithmetical mean,
 and $\sqrt{10 \times 8.6596431} = 9.3057204$ the geom. mean ;
 hence the $\log.$ of 9.3057204 is .96875.

Sixthly, the $\log.$ of 8.6596431 is .9375, and the $\log.$ of
 9.3057204 is .96875 ;

theref. $.9375 + .96875 \div 2 = .953125$ is the arith. mean,
 and $\sqrt{8.6596431 \times 9.3057204} = 8.9768713$ the geome-
 tric mean ;
 hence the $\log.$ of 8.9768713 is .953125.

And proceeding in this manner, after 25 extractions, it will
 be found that the logarithm of 8.9999998 is 9542425 ; which
 may be taken for the logarithm of 9, as it differs so little from
 it, that it is sufficiently exact for all practical purposes. And
 in this manner were the logarithms of almost all the prime
 numbers at first computed.

RULE II.*

LET b be the number whose logarithm is required to be
 found ; and a the number next less than b , so that $b - a = 1$,
 the logarithm of a being known ; and let s denote the sum of
 the two numbers $a + b$. Then

1. Divide the constant decimal .8685889638 &c. by s ,
 and reserve the quotient : divide the reserved quotient by
 the square of s , and reserve this quotient : divide this last
 quotient also by the square of s , and again reserve the
 quotient : and thus proceed, continually dividing the last
 quotient by the square of s , as long as division can be
 made.

2. Then write these quotients orderly under one another,
 the first uppermost, and divide them respectively by the odd
 numbers, 1, 3, 5, 7, 9, &c. as long as division can be made ;
 that is, divide the first reserved quotient by 1, the second by
 3, the third by 5, the fourth by 7, and so on.

3. Add all these last quotients together. and the sum will
 be the logarithm of $b \div a$; therefore to this logarithm add
 also the given logarithm of the said next less number a , so
 will the last sum be the logarithm of the number b proposed.

* For the demonstration of this rule, see my Mathematical Tables, p. 109, &c.

That is,

Log. of b is $\log. a + \frac{n}{s} \times (1 + \frac{1}{3s^2} + \frac{1}{5s^4} + \frac{1}{7s^6} + \&c.)$ where n denotes the constant given decimal $\cdot 8685889638$, &c.

EXAMPLES.

Ex. 1. Let it be required to find the log. of number 2. Here the given number b is 2, and the next less number a is 1, whose log. is 0; also the sum $2+1=3=s$, and its square $s^2=9$. Then the operation will be as follows :

3)	·868588964	1)	·289529654	(·289529654
9)	·289529654	3)	32169962	(10723321
9)	32169962	5)	3574440	(714888
9)	3574440	7)	397160	(56737
9)	397160	9)	44129	(4903
9)	44129	11)	4903	(446
9)	4903	13)	545	(42
9)	545	15)	61	(4
9)	61				

log. of $\frac{2}{1}$ - ·301029995
 add log. 1 - ·000000000

 log. of 2 - ·301029995

Ex. 2. To compute the logarithm of the number 3.

Here $b=3$, the next less number $a=2$, and the sum $a+b=5=s$, whose square s^2 is 25, to divide by which, always multiply by $\cdot 04$. Then the operation is as follows :

5)	·868588964	1)	·173717793	(·173717793
25)	·173717793	3)	6948712	(2316237
25)	6948712	5)	277948	(55590
25)	277948	7)	11118	(1588
25)	11118	9)	445	(50
25)	445	11)	18	(2
	18				

log. of $\frac{3}{2}$ - ·176091260
 log. of 2 add ·301029995

log. of 3 sought ·477121255

Then, because the sum of the logarithms of numbers, gives the logarithm of their product; and the difference of the logarithms, gives the logarithm of the quotient of the numbers;

numbers ; from the above two logarithms, and the logarithm of 10, which is 1, we may raise a great many logarithms, as in the following examples :

<p style="text-align: center;">EXAMPLE 3.</p> <p>Because $2 \times 2 = 4$, therefore</p> <p>to log. 2 - $\cdot 301029995\frac{2}{3}$</p> <p>add log. 2 - $\cdot 301029995\frac{2}{3}$</p> <hr style="width: 50%; margin-left: 0;"/> <p>sum is log. 4 $\cdot 602059991\frac{1}{3}$</p> <hr style="width: 50%; margin-left: 0;"/>	<p style="text-align: center;">EXAMPLE 6.</p> <p>Because $3^2 = 9$, therefore</p> <p>log. 3 - $\cdot 477121254\frac{7}{10}$</p> <p>mult. by 2 2</p> <hr style="width: 50%; margin-left: 0;"/> <p>gives log. 9 $\cdot 954242509$</p> <hr style="width: 50%; margin-left: 0;"/>
<p style="text-align: center;">EXAMPLE 4.</p> <p>Because $2 \times 3 = 6$, therefore</p> <p>to log. 2 - $\cdot 301029995$</p> <p>add log. 3 - $\cdot 477121255$</p> <hr style="width: 50%; margin-left: 0;"/> <p>sum is log. 6 $\cdot 778151250$</p> <hr style="width: 50%; margin-left: 0;"/>	<p style="text-align: center;">EXAMPLE 7.</p> <p>Because $\frac{1}{2} = 5$, therefore</p> <p>from log. 10 $1\cdot 000000000$</p> <p>take log. 2 $\cdot 301029995\frac{2}{3}$</p> <hr style="width: 50%; margin-left: 0;"/> <p>leaves log. 5 $\cdot 698970004\frac{1}{3}$</p> <hr style="width: 50%; margin-left: 0;"/>
<p style="text-align: center;">EXAMPLE 5.</p> <p>Because $2 = 8$, therefore</p> <p>log. 2 - $\cdot 301029995\frac{2}{3}$</p> <p>mult. by 3 3</p> <hr style="width: 50%; margin-left: 0;"/> <p>gives log. 8 $\cdot 903089987$</p> <hr style="width: 50%; margin-left: 0;"/>	<p style="text-align: center;">EXAMPLE 8.</p> <p>Because $3 \times 4 = 12$, therefore</p> <p>to log. 3 - $\cdot 477121255$</p> <p>add log. 4 - $\cdot 602059991$</p> <hr style="width: 50%; margin-left: 0;"/> <p>gives log. 12 $1\cdot 079181246$</p> <hr style="width: 50%; margin-left: 0;"/>

And thus, computing, by this general rule, the logarithms to the other prime numbers, 7, 11, 13, 17, 19, 23, &c. and then using composition and division, we may easily find as many logarithms as we please, or may speedily examine any logarithm in the table*.

* There are, besides these, many other ingenious methods, which later writers have discovered for finding the logarithms of numbers, in a much easier way than by the original inventor; but, as they cannot be understood without a knowledge of some of the higher branches of the mathematics, it is thought proper to omit them, and to refer the reader to those works which are written expressly on the subject. It would likewise much exceed the limits of this compendium, to point out all the particular artifices that are made use of for constructing an entire table of these numbers; but any information of this kind, which the learner may wish to obtain, may be found in my Tables, before mentioned.

Description and Use of the TABLE of LOGARITHMS.

HAVING explained the manner of forming a table of the logarithms of numbers, greater than unity; the next thing to be done is, to show how the logarithms of fractional quantities may be found. In order to this, it may be observed, that as in the former case a geometric series is supposed to increase towards the left, from unity, so in the latter case it is supposed to decrease towards the right hand, still beginning with unit; as exhibited in the general description, page 148, where the indices being made negative, still show the logarithms to which they belong. Whence it appears, that as $+1$ is the log. of 10, so -1 is the log. of $\frac{1}{10}$ or $\cdot 1$; and as $+2$ is the log. of 100, so, -2 is the log. of $\frac{1}{100}$ or $\cdot 01$: and so on.

Hence it appears in general, that all numbers which consist of the same figures, whether they be integral, or fractional, or mixed, will have the decimal parts of their logarithms the same, but differing only in the index, which will be more or less, and positive or negative, according to the place of the first figure of the number.

Thus, the logarithm of 2651 being 3.423410, the log. of $\frac{1}{10}$, or $\frac{1}{100}$, or $\frac{1}{1000}$, &c. part of it; will be as follows:

Numbers.		Logarithms.
2 6 5 1		3.423410
2 6 5.1		2.423410
2 6.5 1		1.423410
2.6 5 1		0.423410
.2 6 5 1		-1.423410
.0 2 6 5 1		-2.423410
.0 0 2 6 5 1		-3.423410

Hence it also appears, that the index of any logarithm, is always less by 1 than the number of integer figures which the natural number consists of; or it is equal to the distance of the first figure from the place of units, or first place of integers, whether on the left or on the right, of it: and this index is constantly to be placed on the left hand side of the decimal part of the logarithm.

When there are integers in the given number, the index is always affirmative; but when there are no integers, the index is negative, and is to be marked by a short line drawn before it, or else above it. Thus.

A number having 1, 2, 3, 4, 5, &c. integer places,
the index of its log. is 0, 1, 2, 3, 4, &c. or 1 less than those places.

And

And a decimal fraction having its first figure in the 1st, 2d, 3d, 4th, &c. place of the decimals, has always — 1, — 2, — 3, — 4, &c. for the index of its logarithm.

It may also be observed, that though the indices of fractional quantities are negative, yet the decimal parts of their logarithms are always affirmative. And the negative mark (—) may be set either before the index or over it.

I. TO FIND, IN THE TABLE, THE LOGARITHM TO ANY NUMBER.*

1. *If the given Number be less than 100, or consist of only two figures; its log. is immediately found by inspection in the first page of the table, which contains all numbers from 1 to 100, with their logs. and the index immediately annexed in the next column.*

So the log. of 5 is 0.698970. The log. of 23 is 1.361728. The log. of 50 is 1.698970. And so on.

2. *If the Number be more than 100 but less than 10000; that is, consisting of either three or four figures; the decimal part of the logarithm is found by inspection in the other pages of the table, standing against the given number, in this manner; viz. the first three figures of the given number in the first column of the page, and the fourth figure one of those along the top line of it; then in the angle of meeting are the last four figures of the logarithm, and the first two figures of the same at the beginning of the same line in the second column of the page: to which is to be prefixed the proper index, which is always 1 less than the number of integer figures.*

So the logarithm of 251 is 2.399674, that is, the decimal .399674 found in the table, with the index 2 prefixed, because the given number contains three integers. And the log. of 34.09 is 1.532627, that is, the decimal .532627 found in the table, with the index 1 prefixed, because the given number contains two integers.

3. *But if the given Number contain more than four figures; take out the logarithm of the first four figures by inspection in the table, as before, as also the next greater logarithm, subtracting the one logarithm from the other, as also their corresponding numbers the one from the other. Then say,*

As the difference between the two numbers,
Is to the difference of their logarithms,
So is the remaining part of the given number,
To the proportional part of the logarithm.

* See the table of Logarithms at the end of the 2d volume.

Which part being added to the less logarithm, before taken out, gives the whole logarithm sought very nearly.

EXAMPLE.

To find the logarithm of the number 34·0926.
 The log. of 340900, as before is 532627.
 And log. of 341000 - - is 532754.
 The diffs. are 100 and 127

Then as 100 : 127 :: 26 : 33, the proportional part.
 This added to - - 532627, the first log.
 Gives, with the index, 1·532660, for the log. of 34·0926.

4. If the number consist both of integers and fractions, or is entirely fractional; find the decimal part of the logarithm the same as if all its figures were integral; then this, having prefixed to it the proper index, will give the logarithm required.

5. And if the given number be a proper vulgar fraction: subtract the logarithm of the denominator from the logarithm of the numerator, and the remainder will be the logarithm sought; which, being that of a decimal fraction, must always have a negative index.

6. But if it be a mixed number; reduce it to an improper fraction, and find the difference of the logarithms of the numerator and denominator, in the same manner as before.

EXAMPLES.

<p>1. To find the log. of $\frac{37}{94}$.</p> <p>Log. of 37 - 1·568202</p> <p>Log. of 94 - 1·973128</p> <hr style="width: 100px; margin-left: 100px;"/> <p>Dif. log. of $\frac{37}{94}$ -1·595074</p> <hr style="width: 100px; margin-left: 100px;"/> <p>Where the index 1 is negative.</p>	<p>2. To find the log. of $17\frac{1}{3}$</p> <p>First, $17\frac{1}{3} = \frac{52}{3}$. Then</p> <p>Log. of 405 - 2·607455</p> <p>Log. of 23 - 1·361728</p> <hr style="width: 100px; margin-left: 100px;"/> <p>Dif. log. of $17\frac{1}{3}$ 1·245727</p> <hr style="width: 100px; margin-left: 100px;"/>
--	---

II. TO FIND THE NATURAL NUMBER TO ANY GIVEN LOGARITHM.

THIS is to be found in the tables by the reverse method to the former, namely, by searching for the proposed logarithm among those in the table, and taking out the corresponding number by inspection, in which the proper number of integers are to be pointed off, viz. 1 more than the index. For, in finding the number answering to any given logarithm, the index always shows how far the first figure must

must be removed from the place of units, viz. to the left hand, or integers, when the index is affirmative ? but to the right hand, or decimals, when it is negative.

EXAMPLES.

So, the number to the log: 1.532882 is 34.11.

And the number of the log. 1.532882 is .3411.

But if the logarithm cannot be exactly found in the table ; take out the next greater and the next less, subtracting the one of these logarithms from the other, as also their natural numbers the one from the other, and the less logarithm from the logarithm proposed. Then say,

As the difference of the first or tabular logarithms,
Is to the difference of their natural numbers,
So is the differ. of the given log. and the least tabular log.
To their corresponding numeral difference.

Which being annexed to the least natural number above taken, gives the natural number sought, corresponding to the proposed logarithm.

EXAMPLE.

So, to find the natural number answering to the given logarithm 1.532708.

Here the next greater and next less tabular logarithms, with their corresponding numbers, are as below :

Next greater 532744 its num. 341000 ; given log. 532708
Next less 532627 its num. 340900 ; next less 532627

Differences	<u>127</u>	<u>100</u>	<u>81</u>
-------------	------------	------------	-----------

Then, as 127 : 100 :: 81 : 64 nearly, the numeral differ. Therefore 34.0964 is the number sought, marking off two integers, because the index of the given logarithm is 1.

Had the index been negative, thus 1.532708, its corresponding number would have been .340965, wholly decimal.

MULTIPLI-

MULTIPLICATION BY LOGARITHMS.

RULE.

TAKE out the logarithms of the factors from the table, then add them together, and their sum will be the logarithm of the product required. Then, by means of the table, take out the natural number, answering to the sum, for the product sought.

Observing to add what is to be carried from the decimal part of the logarithm to the affirmative index or indices, or else subtract it from the negative.

Also, adding the indices together when they are of the same kind, both affirmative or both negative; but subtracting the less from the greater, when the one is affirmative and the other negative, and prefixing the sign of the greater to the remainder,

EXAMPLES.

1. To Multiply 23·14 by
5·062.

Numbers.	Logs.
23·14	- 1·364363
5·062	- 0·704322

Product 117·1347 2·068685

2. To multiply 2·581926
by 3·457291.

Numbers.	Logs.
2·581926	- 0·411944
3·457291	- 0·538736

Prod. 8·92648 0·950680

3. To mult. 3·902 and 597·16
and ·0314728 all together.

Numbers.	Logs.
3·902	- 0·591287
597·16	- 2·776091
·0314728	- 2·497935

Prod. 73·3333 1·865313

Here the—2 cancels the 2,
and the 1 to carry from the
decimals is set down.

4. To mult. 3·586, and 2·1046,
and 0·8372, and 0·0294 all
together.

Numbers.	Logs.
3·586	- 0·554610
2·1046	- 0·323170
0·8372	- 1·922829
0·0294	- 2·468347

Prod. 0·1057618—1·268956

Here the 2 to carry can-
cels the—2, and there re-
mains the—1 to set down.

DIVISION

DIVISION BY LOGARITHMS.

RULE.

From the logarithm of the dividend subtract the logarithm of the divisor, and the number answering to the remainder will be the quotient required.

Observing to change the sign of the index of the divisor, from affirmative to negative, or from negative to affirmative; then take the sum of the indices if they be of the same name, or their difference when of different signs, with the sign of the greater, for the index to the logarithm of the quotient.

And also, when 1 is borrowed, in the left-hand place of the decimal part of the logarithm, add it to the index of the divisor when that index is affirmative, but subtract it when negative; then let the sign of the index arising from hence be changed, and worked with as before.

EXAMPLES.

<p>1. To divide 24163 by .4567.</p> <table border="0"> <tr> <td></td> <td style="text-align: center;">Numbers.</td> <td style="text-align: center;">Logs.</td> </tr> <tr> <td>Dividend</td> <td>24163</td> <td>- 4.383151</td> </tr> <tr> <td>Divisor</td> <td>- 4567</td> <td>- 3.659631</td> </tr> <tr> <td>Quot.</td> <td>5.29078</td> <td>0.723520</td> </tr> </table>		Numbers.	Logs.	Dividend	24163	- 4.383151	Divisor	- 4567	- 3.659631	Quot.	5.29078	0.723520	<p>2. To divide 37.149 by 523.76</p> <table border="0"> <tr> <td></td> <td style="text-align: center;">Numbers.</td> <td style="text-align: center;">Logs.</td> </tr> <tr> <td>Dividend</td> <td>37.149</td> <td>- 1.569947</td> </tr> <tr> <td>Divisor</td> <td>523.76</td> <td>- 2.719132</td> </tr> <tr> <td>Quot.</td> <td>.0709275</td> <td>- 2.850815</td> </tr> </table>		Numbers.	Logs.	Dividend	37.149	- 1.569947	Divisor	523.76	- 2.719132	Quot.	.0709275	- 2.850815
	Numbers.	Logs.																							
Dividend	24163	- 4.383151																							
Divisor	- 4567	- 3.659631																							
Quot.	5.29078	0.723520																							
	Numbers.	Logs.																							
Dividend	37.149	- 1.569947																							
Divisor	523.76	- 2.719132																							
Quot.	.0709275	- 2.850815																							
<p>3. Divide .06314 by .007241</p> <table border="0"> <tr> <td></td> <td style="text-align: center;">Numbers.</td> <td style="text-align: center;">Logs.</td> </tr> <tr> <td>Divid.</td> <td>.06314</td> <td>- 2.800305</td> </tr> <tr> <td>Divisor</td> <td>.007241</td> <td>- 3.859799</td> </tr> <tr> <td>Quot.</td> <td>8.71979</td> <td>0.940506</td> </tr> </table>		Numbers.	Logs.	Divid.	.06314	- 2.800305	Divisor	.007241	- 3.859799	Quot.	8.71979	0.940506	<p>4. To divide .7438 by 12.9476</p> <table border="0"> <tr> <td></td> <td style="text-align: center;">Numbers.</td> <td style="text-align: center;">Logs.</td> </tr> <tr> <td>Divid.</td> <td>.7438</td> <td>- 1.871456</td> </tr> <tr> <td>Divisor</td> <td>12.9476</td> <td>1.112189</td> </tr> <tr> <td>Quot.</td> <td>.057447</td> <td>- 2.759267</td> </tr> </table>		Numbers.	Logs.	Divid.	.7438	- 1.871456	Divisor	12.9476	1.112189	Quot.	.057447	- 2.759267
	Numbers.	Logs.																							
Divid.	.06314	- 2.800305																							
Divisor	.007241	- 3.859799																							
Quot.	8.71979	0.940506																							
	Numbers.	Logs.																							
Divid.	.7438	- 1.871456																							
Divisor	12.9476	1.112189																							
Quot.	.057447	- 2.759267																							

Here 1 carried from the decimals to the — 3, makes it become — 2, which taken from the other — 2, leaves 0 remaining.

Here the 1 taken from the — 1, makes it become — 2, to set down.

Note. As to the Rule-of-Three, or Rule of Proportion, it is performed by adding the logarithms of the 2d and 3d terms, and subtracting that of the first term from their sum.

INVOLUTION BY LOGARITHMS.

RULE.

TAKE out the logarithm of the given number from the table. Multiply the log thus found, by the index of the power proposed. Find the number answering to the product, and it will be the power required.

Note. In multiplying a logarithm with a negative index, by an affirmative number, the product will be negative. But what is to be carried from the decimal part of the logarithm, will always be affirmative. And therefore their difference will be the index of the product, and is always to be made of the same kind with the greater.

EXAMPLES.

1. To square the number
2·5791.

Numb.	Log.
Root 2·5791 - -	0·411468
The index - -	2
<hr/>	
Power 6·65174	0·822936
<hr/>	

2. To find the cube of
3·07146.

Numb.	Log.
Root 3·07146 - - -	0·487345
The index - - -	3
<hr/>	
Power 28·9758	1·462035
<hr/>	

3. To raise ·09163 to the 4th
power.

Numb.	Log.
Root ·09163	—2·962038
The index - -	4
<hr/>	
Pow. ·000070494	—5·848152
<hr/>	

4. To raise 1·0045 to the
365th power.

Numb.	Log.
Root 1·0045 - -	0·001950
The index - -	365
<hr/>	
	9750
	11700
	5850
<hr/>	
Power 5·14932*	0·711750
<hr/>	

Here 4 times the negative index being —8, and 3 to carry, the difference —5 is the index of the product.

* This answer 5·14932 though found strictly according to the general rule, is not correct in the last two figures 32; nor can the answers to such questions relating to very high powers be generally found true to 6 places of figures by the table of logarithms in this work: if any power above the hundred thousandth were required, not one figure of the answer found by the table of logarithms here given could be depended on.

The logarithm of 1·0045 is 00194994103 true to eleven places, which multiplied by 365 gives ·7117285 true to 7 places, and the corresponding number true to 7 places is 5·149067. *Ed.*

EVOLUTION BY LOGARITHMS.

TAKE the log. of the given number out of the table.

Divide the log. thus found by the index of the root. Then the number answering to the quotient, will be the root.

Note. When the index of the logarithm, to be divided, is negative, and does not exactly contain the divisor, without some remainder, increase the index by such a number as will make it exactly divisible by the index, carrying the units borrowed, as so many tens, to the left-hand place of the decimal, and then divide as in whole numbers.

Ex. 1. To find the square root of 365.

	Numb.	Log.
Power	365 2)	2·562293
Root	19·10496	1·281146½

Ex. 2. To find the 3d root of 12345.

	Numb.	Log.
Power	12345 3)	4·091491
Root	23·1116	1·363830½

Ex. 3. To find the 10th root of 2.

	Numb.	Log.
Power	2 - - 10)	0·301030
Root	1·071773	0·030103

Ex. 4. To find the 365th root of 1·045.

	Numb.	Log.
Power	1·045 365)	0·019116
Root	1·000121	0·000052½

Ex. 5. To find $\sqrt[2]{093}$.

	Numb.	Log.
Power	·093 2)	-2·968483
Root	·304959	-1·484241½

Ex. 6. To find the $\sqrt[3]{\cdot00048}$.

	Numb.	Log.
Power	·00048 3)	-4·681241
Root	·0782973	-2·893747

Here the divisor 2 is contained exactly once in the negative index - 2, and therefore the index of the quotient is - 1.

Here the divisor 3 not being exactly contained in -4, it is augmented by 2, to make up 6, in which the divisor is contained just 2 times; then the 2, thus borrowed, being carried to the decimal figure 6, makes 26, which divided by 3, gives 8, &c.

Ex. 7. To find $3\cdot1416 \times 82 \times \frac{13}{41}$.

Ex. 8. To find $\cdot02916 \times 751\cdot3 \times \frac{9}{341}$.

Ex. 9. As 7241 : 3·58 :: 20·46 : ?

Ex. 10. As $\sqrt[2]{724} : \sqrt[3]{\frac{51}{13}} :: 6\cdot927 : ?$

ALGEBRA.

DEFINITIONS AND NOTATION.

1. **ALGEBRA** is the science of computing by symbols. It is sometimes also called Analysis ; and is a general kind of arithmetic, or universal way of computation.

2. In this science, quantities of all kinds are represented by the letters of the alphabet. And the operation to be performed with them, as addition or subtraction, &c. are denoted by certain simple characters, instead of being expressed by words at length.

3. In algebraical questions, some quantities are known or given, viz. those whose values are known : and others unknown, or are to be found out, viz. those whose values are not known. The former of these are represented by the leading letters of the alphabet, $a, b, c, d,$ &c. ; and the latter, or unknown quantities, by the final letters, $z, y, x, u,$ &c.

4. The characters used to denote the operations, are chiefly the following :

$+$ signifies addition, and is named *plus*.

$-$ signifies subtraction, and is named *minus*.

\times or $.$ signifies multiplication, and is named *into*.

\div signifies division, and is named *by*.

$\sqrt{\quad}$ signifies the square root ; $\sqrt[3]{\quad}$ the cube root ; $\sqrt[4]{\quad}$ the 4th root, &c. ; and $\sqrt[n]{\quad}$ the n th root.

$:::$ signifies proportion.

$=$ signifies equality, and is named *equal to*.

And so on for other operations.

Thus $a+b$ denotes that the number represented by b is to be added to that represented by a .

$a-b$ denotes, that the number represented by b is to be subtracted from that represented by a .

$a \oslash b$ denotes the difference of a and b , when it is not known which is the greater.

ab , or $a \times b$, or $a.b$, expresses the product, by multiplication, of the numbers represented by a and b .

$a \div b$,

$a \div b$, or $\frac{a}{b}$, denotes, that the number represented by a is

to be divided by that which is expressed by b .

$a : b :: c : d$, signifies that a is in the same proportion to b , as c is to d .

$x = a - b + c$ is an equation, expressing that x is equal to the difference of a and b , added to the quantity c .

$a > b$ signifies that a is greater than b .

$a < b$ signifies that a is less than b .

\sqrt{a} , or $a^{\frac{1}{2}}$, denotes the square root of a ; $\sqrt[3]{a}$, or $a^{\frac{1}{3}}$, the cube root of a ; and $\sqrt[3]{a^2}$ or $a^{\frac{2}{3}}$ the cube root of the square of a ; also $\sqrt[m]{a}$, or $a^{\frac{1}{m}}$, is the m th root, of a ; and $\sqrt[m]{a^n}$ or a_m^n is the n th power of the m th root of a , or it is a to the $\frac{n}{m}$ power.

a^2 denotes the square of a ; a^3 the cube of a ; a^4 the fourth power of a ; and a^n the n th power of a .

$\overline{a+b} \times c$, or $(a+b)c$, denotes the product of the compound quantity $a + b$ multiply by the simple quantity c . Using the bar $\overline{\quad}$, or the parenthesis (\quad) as a vinculum, to connect several simple quantities into one compound.

$\overline{a+b} \div \overline{a-b}$ or $\frac{a+b}{a-b}$, expressed like a fraction, means the

quotient of $a+b$ divided by $a-b$.

$\sqrt{ab+cd}$, or $(ab+cd)^{\frac{1}{2}}$, is the square root of the compound quantity $ab+cd$. And $c \sqrt{ab+cd}$, or $c(ab+cd)^{\frac{1}{2}}$, denotes the product of c into the square root of the compound quantity $ab+cd$.

$\overline{a+b-c}^3$, or $(a+b-c)^3$, denotes the cube, or third power, of the compound quantity $a+b-c$.

$3a$ denotes that the quantity a is to be taken 3 times, and $4(a+b)$ is 4 times $a+b$. And these numbers, 3 or 4, showing how often the quantities are to be taken, or multiplied, are called Co-efficients.

Also $\frac{3}{4}x$ denotes that x is multiplied by $\frac{3}{4}$; thus $\frac{3}{4} \times x$ or $\frac{3}{4}x$.

5. Like Quantities, are those which consist of the same letters, and powers. As a and $3a$; or $2ab$ and $4ab$; or $3a^2bc$ and $-5a^2bc$.

6. Unlike Quantities, are those which consist of different letters, or different powers. As a and b ; or $2a$ and a^2 ; or $3ab^2$ and $3abc$.

7. Simple

7. Simple Quantities, are those which consist of one term only. As $3a$, or $5ab$, or $6abc^2$.

8. Compound Quantities are those which consist of two or more terms. As $a+b$, or $2a-3c$, or $a+2b-3c$.

9. And when the compound quantity consists of two terms it is called a Binomial, as $a+b$; when of three terms, it is a Trinomial, as $a+2b-3c$; when of four terms, a Quadrinomial, as $2a-3b+c-4d$; and so on. Also, a Multinomial or Polynomial, consists of many terms.

10. A Residual Quantity, is a binomial having one of the terms negative. As $a-2b$.

11. Positive or Affirmative Quantities, are those which are to be added, or have the sign $+$. As a or $+a$, or ab : for when a quantity is found without a sign, it is understood to be positive, or have the sign $+$ prefixed.

12. Negative Quantities, are those which are to be subtracted. As $-a$, or $2ab$, or $3ab^2$.

13. Like Signs, are either all positive ($+$), or all negative ($-$).

14. Unlike Signs, are when some are positive ($+$), and others negative ($-$).

15. The Co-efficient of any quantity, as shown above, is the number prefixed to it. As 3, in the quantity $3ab$.

16. The Power of a quantity (a), is its square (a^2), or cube (a^3), or biquadrate (a^4), &c; called also, the 2d power, or 3d power, or 4th power, &c.

17. The Index or Exponent, is the number which denotes the power or root of a quantity. So 2 is the exponent of the square or second power a^2 ; and 3 is the index of the cube or 3d power; and $\frac{1}{2}$ is the index of the square root, $a^{\frac{1}{2}}$ or \sqrt{a} ; and $\frac{1}{3}$ is the index of the cube root, $a^{\frac{1}{3}}$, or $\sqrt[3]{a}$.

18. A Rational Quantity, is that which has no radical sign ($\sqrt{\quad}$) or index annexed to it. As a , or $3ab$.

19. An Irrational Quantity, or Surd, is that of which the value cannot be accurately expressed in numbers, as the square roots of 2, 3, 5. Surds are commonly expressed by means of the radical sign $\sqrt{\quad}$, as $\sqrt{2}$, \sqrt{a} , $\sqrt[3]{a^2}$, or $ab^{\frac{1}{2}}$.

20. The Reciprocal of any quantity, is that quantity inverted, or unity divided by it. So, the reciprocal of a , or

$\frac{a}{1}$, is $\frac{1}{a}$, and the reciprocal of $\frac{a}{b}$ is $\frac{b}{a}$.

21. The letters by which any simple quantity is expressed, may be ranged according to any order at pleasure. So the product of a and b , may be either expressed by ab , or ba ; and the product of a , b , and c , by either abc , or acb , or bac , or bca , or cab , or cba ; as it matters not which quantities are placed or multiplied first. But it will be sometimes found convenient in long operations, to place the several letters according to their order in the alphabet, as abc , which order also occurs most easily or naturally to the mind.

22. Likewise, the several members, or terms, of which a compound quantity is composed, may be disposed in any order at pleasure, without altering the value of the signification of the whole. Thus, $3a - 2ab + 4abc$ may also be written $3a + 4abc - 2ab$, or $4abc + 3a - 2ab$, or $-2ab + 3a + 4abc$, &c; for all these represent the same thing, namely, the quantity which remains, when the quantity or term $2ab$ is subtracted from the sum of the terms or quantities $3a$ and $4abc$. But it is most usual and natural, to begin with a positive term, and with the first letters of the alphabet.

SOME EXAMPLES FOR PRACTICE.

In finding the numeral values of various expressions, or combinations, of quantities.

Supposing $a=6$, and $b=5$, and $c=4$, and $d=1$, and $e=0$.
Then

1. Will $a^2 + 3ab - c^2 = 36 + 90 - 16 = 110$.

2. And $2a^3 - 3a^2b + c^3 = 432 - 540 + 64 = -44$.

3. And $a^2 \times a + b - 2abc = 36 \times 11 - 240 = 156$.

4. And $\frac{a^3}{a+3c} + c^2 = \frac{216}{18} + 16 = 12 + 16 = 28$.

5. And $\sqrt{2ac+c^2}$ or $2ac+c^2 \} \frac{1}{2} = \sqrt{64} = 8$.

6. And $\sqrt{c + \frac{2bc}{\sqrt{2ac+c^2}}} = 2 + \frac{40}{8} = 7$.

7. And $\frac{a^2 - \sqrt{b^2 - ac}}{2a - \sqrt{b^2 + ac}} = \frac{36 - 1}{12 - 7} = \frac{35}{5} = 7$.

8. And $\sqrt{b^2 - ac + \sqrt{2ac + c^2}} = 1 + 8 = 9$.

9. And $\sqrt{b^2 - ac + \sqrt{2ac + c^2}} = \sqrt{25 - 24 + 8} = 3$.

10. And $a^2b + c - d = 183$.

11. And $9ab - 10b^2 + c = 24$.

12. And

12. And $\frac{a^2b}{c} \times d = 45.$

13. And $\frac{a+b}{c} \times \frac{b}{d} = 13\frac{3}{4}.$

14. And $\frac{a+b}{c} - \frac{a-b}{d} = 1\frac{3}{4}.$

15. And $\frac{a^2b}{c} + e = 45.$

16. And $\frac{a^2b}{c} \times e = 0.$

17. And $\overline{b-c} \times \overline{d-e} = 1.$

18. And $\overline{a+b} - \overline{c-d} = 8.$

19. And $\overline{a+b} - c - d = 6.$

20. And $a^2c \times d^3 = 144.$

21. And $acd - d = 23.$

22. And $a^2e + b^2e + d = 1.$

23. And $\frac{b-e}{d-e} \times \frac{a+b}{c-d} = 18\frac{1}{3}.$

24. And $\sqrt{a^2+b^2} - \sqrt{a^2-b^2} = 4.4936249.$

25. And $3ac^2 + \sqrt[3]{a^3-b^3} = 292.497942.$

26. And $4a^2 - 3a\sqrt{a^2} - \frac{2}{3}ab = 72.$

ADDITION.

ADDITION, in Algebra, is the connecting the quantities together by their proper signs, and incorporating or uniting into one term or sum, such as are similar, and can be united. As, $3a+2b-2a=a+2b$, the sum.

The rule of addition in algebra, may be divided into three cases: one when the quantities are like, and their signs like also; a second, when the quantities are like, but their signs unlike; and the third, when the quantities are unlike. Which are performed as follows.*

CASE

* The reasons on which these operations are founded, will readily appear, by a little reflection on the nature of the quantities to be added or collected together.
For

CASE I.

When the Quantities are Like, and have Like Signs.

ADD the co-efficients together, and set down the sum ; after which set the common letter or letters of the like quantities, and prefix the common sign $+$ or $-$.

For, with regard to the first example, where the quantities are $3a$ and $5a$ whatever a represents in the one term, it will represent the same thing in the other ; so that three times any thing and 5 times the same thing, collected together, must needs make 8 times that thing. As if a denote a shilling ; then $3a$ is 3 shillings ; and $5a$ is 5 shillings, and their sum 8 shillings. In like manner, $-2ab$ and $-7ab$, or -2 times any thing, and -7 times the same thing, make -9 times that thing.

As to the second case, in which the quantities are like, but the signs unlike ; the reason of its operation will easily appear, by reflecting, that addition means only the uniting of quantities together by means of the arithmetical operations denoted by their signs $+$ and $-$, or of addition and subtraction ; which being of contrary or opposite natures, the one co-efficient must be subtracted from the other, to obtain the incorporated or united mass.

As to the third case, where the quantities are unlike, it is plain that such quantities cannot be united into one, or otherwise added, than by means of their signs ; thus, for example, if a be supposed to represent a crown, and b a shilling ; then the sum of a and b can be neither $2a$ nor $2b$, that is neither 2 crowns nor 2 shillings, but only 1 crown plus 1 shilling, that is $a + b$.

In this rule, the word *addition* is not very properly used ; being much too limited to express the operation here performed. The business of this operation is to incorporate into one mass, or algebraic expression, different algebraic quantities, as far as an actual incorporation or union is possible ; and to retain the algebraic marks for doing it, in cases where the former is not possible. When we have several quantities, some affirmative and some negative ; and the relation of these quantities can in the whole or in part be discovered ; such incorporation of two or more quantities into one, is plainly effected by the foregoing rules.

It may seem a paradox, that what is called addition in algebra, should sometimes mean addition, and sometimes subtraction. But the paradox wholly arises from the scantiness of the name given to the algebraic process ; from employing an old term in a new and more enlarged sense. Instead of addition, call it incorporation, or union, or striking a balance, or any name to which a more extensive idea may be annexed, than that which is usually implied by the word addition ; and the paradox vanishes.

Thus,

ADDITION.

177

Thus, $3a$ added to $5a$, makes $8a$.

And $-2ab$ added to $-7ab$, makes $-9ab$.

And $5a+7b$ added to $7a+3b$, makes $12a+10b$.

OTHER EXAMPLES FOR PRACTICE.

$3a$	$-3bx$	bxy
$9a$	$-5bx$	$2bxy$
$5a$	$-4bx$	$5bxy$
$12a$	$-2bx$	bxy
a	$-7bx$	$3bxy$
$2a$	$-bx$	$6bxy$
$32a$	$-22bx$	$18bxy$

$3z$	$3x^2+5xy$	$2ax-4y$
$2z$	x^2+xy	$4ax-y$
$4z$	$2x^2+4xy$	$ax-3y$
z	$5x^2+2xy$	$5ax-5y$
$5z$	$4x^2+3xy$	$7ax-2y$
$15z$	$15x^3+15xy$	$19ax-15y$

$5xy$	$-12y^2$	$4a-4b$
$14xy$	$-7y^2$	$5a-5b$
$22xy$	$-2y^2$	$6a-b$
$17xy$	$-4y^2$	$3a-2b$
$1\frac{1}{2}xy$	$-y^2$	$2a-7b$
$\frac{1}{3}xy$	$-3y^2$	$8a-b$

$30-13x^{\frac{1}{2}}-3xy$	$5xy-3x+4ab$
$23-10x^{\frac{1}{2}}-4xy$	$8xy-4x+3ab$
$14-14x^{\frac{1}{2}}-7xy$	$3xy-5x+5ab$
$10-16x^{\frac{1}{2}}-5xy$	$xy-2x+ab$
$16-20x^{\frac{1}{2}}-xy$	$4xy-x+7ab$

CASE II.

When the Quantities are Like, but have Unlike Signs ;

ADD the affirmative co-efficients into one sum, and all the negative ones into another, when there are several of a kind. Then subtract the less sum, or the less co-efficient, from the greater, and to the remainder prefix the sign of the greater, and subjoin the common quantity or letters.

So $+ 5a$ and $- 3a$, united make $+ 2a$.

And $- 5a$ and $+ 3a$, united, make $- 2a$.

OTHER EXAMPLES FOR PRACTICE.

$- 5a$	$+ 3ax^2$	$+ 8x^3 + 3y$
$+ 4a$	$+ 4ax^2$	$- 5x^3 + 4y$
$+ 6a$	$- 8ax^2$	$- 16x^3 + 5y$
$- 3a$	$- 6ax^2$	$+ 3x^3 - 7y$
$+ a$	$+ 5ax^2$	$+ 2x^3 - 2y$
$+ 3a$	$- 2ax^2$	$- 8x^3 + 3y$

$- 3a^2$	$+ 3b^2y^2$	$+ 4ab + 4$
$- 5a^2$	$+ 9b^2y^2$	$- 4ab + 12$
$- 10a^2$	$- 10b^2y^2$	$+ 7ab - 14$
$+ 10a^2$	$- 19b^2y^2$	$+ ab + 3$
$+ 14a^2$	$- 2b^2y^2$	$- 5ab - 10$

$- 3ax^{\frac{1}{2}}$	$+ 10\sqrt{ax}$	$+ 3y + 4ax^{\frac{1}{2}}$
$+ ax^{\frac{1}{2}}$	$- 3\sqrt{ax}$	$- y - 5ax^{\frac{1}{2}}$
$+ 5ax^{\frac{1}{2}}$	$+ 4\sqrt{ax}$	$+ 4y + 2ax^{\frac{1}{2}}$
$- 6ax^{\frac{1}{2}}$	$- 12\sqrt{ax}$	$- 2y + 6ax^{\frac{1}{2}}$

CASE III.

When the Quantities are Unlike.

HAVING collected together all the like quantities, as in the two foregoing cases, set down those that are unlike, one after another, with their proper signs.

EXAMPLES.

$3xy$	$6xy - 12x^2$	$4ax - 130 + 3x^{\frac{1}{2}}$
$2ax$	$-4x^2 + 3xy$	$5x^2 + 3ax + 9x^2$
$-5xy$	$+4x^2 - 2xy$	$7xy - 4x^{\frac{1}{2}} + 90$
$6ax$	$-3xy + 4x^2$	$\sqrt{x} + 40 - 6x^2$
$-2xy + 8ax$	$4xy - 8x^2$	$7ax + 8x^2 + 7xy$

$9x^2y$	$14ax - 2x^2$	$9 + 10\sqrt{ax} - 5y$
$-7x^2y$	$5ax + 3xy$	$2x + 7\sqrt{xy} + 5y$
$+3axy$	$8y^2 - 4ax$	$5y + 3\sqrt{ax} - 4y$
$-4x^2y$	$3x^2 + 26$	$10 - 4\sqrt{ax} + 4y$

$4x^2y$	$4\sqrt{x} - 3y$	$3a^2 + 9 + x^{\frac{1}{2}} - 4$
$-6xy^2$	$2\sqrt{xy} + 14x$	$2a - 8 + 2a^2 - 3x$
$+3y^2x$	$3x + 2y$	$4x^2 - 2a^2 + 18 - 7$
$-7x^2y$	$-9 + 2\sqrt{xy}$	$-12 + a - 3x^2 - 2y$

Add $a + b$ and $3a - 5b$ together.

Add $5a - 8x$ and $3a - 4x$ together.

Add $6x - 5b + a + 8$ to $-5a - 4x + 4b - 3$.

Add $a + 2b - 3c - 10$ to $3b - 4a + 5c + 10$ and $5b - c$.

Add $a + b$ and $a - b$ together.

Add $3a + b - 10$ to $c - d - a$ and $-4c + 2a - 3b - 7$.

Add $3a^2 + b^2 - c$ to $2ab - 3a^2 + bc - b$.

Add $a^3 + b^2c - b^2$ to $ab^2 - abc + b^2$.

Add $9a - 8b + 10x - 6d - 7c + 50$ to $2x - 3a - 5c + 4b + 6d - 10$.

SUBTRAC-

SUBTRACTION.

SET down in one line the first quantities from which the subtraction is to be made; and underneath them place all the other quantities composing the subtrahend: ranging the like quantities under each other as in Addition.

Then change all the signs (+ and -) of the lower line, or conceive them to be changed; after which, collect all the terms together as in the cases of Addition.*

EXAMPLES.

From $7a^2 - 3b$ Take $3a^2 - 8b$ <hr style="width: 80%; margin: 0 auto;"/> Rem. $4a^2 + 5b$ <hr style="width: 80%; margin: 0 auto;"/>	$9x^2 - 4y + 8$ $6x^2 + 5y - 4$ <hr style="width: 80%; margin: 0 auto;"/> $3x^2 - 9y + 12$ <hr style="width: 80%; margin: 0 auto;"/>	$8xy - 3 + 6x - y$ $4xy - 7 - 6x - 4y$ <hr style="width: 80%; margin: 0 auto;"/> $4xy + 4 + 12x + 5y$ <hr style="width: 80%; margin: 0 auto;"/>
From $5xy - 6$ Take $-2xy + 6$ <hr style="width: 80%; margin: 0 auto;"/> Rem. $7xy - 12$ <hr style="width: 80%; margin: 0 auto;"/>	$4y^2 - 3y - 4$ $2y^2 + 2y + 4$ <hr style="width: 80%; margin: 0 auto;"/> $2y^2 - 5y - 8$ <hr style="width: 80%; margin: 0 auto;"/>	$-20 - 6x - 5xy$ $3xy - 9x + 8 - 2ay$ <hr style="width: 80%; margin: 0 auto;"/> $-28 + 3x - 8xy + 2ay$ <hr style="width: 80%; margin: 0 auto;"/>
From $8x^2y - 6$ Take $-2x^2y + 2$ <hr style="width: 80%; margin: 0 auto;"/> Rem. _____ <hr style="width: 80%; margin: 0 auto;"/>	$5\sqrt{xy} + 2x\sqrt{xy}$ $7\sqrt{xy} + 3 - 2xy$ <hr style="width: 80%; margin: 0 auto;"/> _____ <hr style="width: 80%; margin: 0 auto;"/>	$7x^2 + 2\sqrt{x} - 18 + 3b$ $9x^2 - 12 + 5b + x^{\frac{1}{2}}$ <hr style="width: 80%; margin: 0 auto;"/> _____ <hr style="width: 80%; margin: 0 auto;"/>

* This rule is founded on the consideration, that addition and subtraction are opposite to each other in their nature and operation, as are the signs + and -, by which they are expressed and represented. So that, since to unite a negative quantity with a positive one of the same kind, has the effect of diminishing it, or subtracting an equal positive one from it, therefore to subtract a positive (which is the opposite of uniting or adding) is to add the equal negative quantity. In like manner, to subtract a negative quantity, is the same in effect as to add or unite an equal positive one. So that, by changing the sign of a quantity from + to -, or from - to +, changes its nature from a subtractive quantity to an additive one; and any quantity is in effect subtracted, by barely changing its sign.

$$\begin{array}{r} 5xy-30 \\ 7xy-50 \\ \hline \end{array}$$

$$\begin{array}{r} 7x^3-2(a+b) \\ 2x^2-4(a+b) \\ \hline \end{array}$$

$$\begin{array}{r} 3xy^3+20a\sqrt{(xy+10)} \\ 4x^2y^2+12a\sqrt{(xy+10)} \\ \hline \end{array}$$

From $a+b$, take $a-b$.

From $4a+4b$, take $b+a$.

From $4a-4b$, take $3a+5b$.

From $8a-12x$, take $4a-3x$.

From $2x-4a-2b+5$, take $8-5b+a+6x$.

From $3a+b+c-d-10$, take $c+2a-d$.

From $3a+b+c-d-10$, take $b-19+3a$.

From $2ab+b^2-4c+bc-b$, take $3a^2-c+b^2$.

From $a^3+3b^2c+ab^2-abc$, take b^2+ab^2-abc .

From $12x+6a-4b+40$, take $4b-3a+4x+6d-10$.

From $2x-3a+4b+6c-50$, take $9a+x+6b-6c-40$

From $6a-4b-12c+12x$, take $2x-8a+4b-5c$.

MULTIPLICATION.

This consists of several cases, according as the factors are simple or compound quantities.

CASE I. *When both the Factors are Simple Quantities;*

FIRST multiply the co-efficients of the two terms together, then to the product annex all the letters in those terms, which will give the whole product required.

*Note.** Like signs, in the factors, produce + and unlike signs -, in the products.

EXAMPLES.

* That this rule for the signs is true, may be thus shown.

1. When + a is to be multiplied by + c; the meaning is, that + a is to be taken as many times as there are units in c; and since the sum of any number of positive terms is positive, it follows that + a X + c makes + ac.

2. When

EXAMPLES.

$10a$	$-3a$	$7a$	$-6x$
$2b$	$+2b$	$-4c$	$-4a$
$20ab$	$-6ab$	$-28ac$	$+24ax$
$4ac$	$9a^2x$	$-2x^2y$	$-4xy$
$-3ab$	$4x$	$3xy^2$	$-xy$
$-12a^2bc$	$36a^2x^2$	$-6x^3y^3$	$+4x^2y^2$
$-3ax$	$-ax$	$+3xy$	$-5xyz$
$4x$	$-6c$	-4	$-5ax$

CASE II.

When one of the Factors is a Compound Quantity.

MULTIPLY every term of the multiplicand, or compound quantity, separately, by the multiplier, as in the former case; placing the products one after another, with the proper signs; and the result will be the whole product required.

2. When two quantities are to be multiplied together, the result will be exactly the same, in whatever order they are placed; for a times c is the same as c times a , and therefore, when $-a$ is to be multiplied by $+c$, or $+c$ by $-a$, this is the same thing as taking $-a$ as many times as there are units in $+c$; and as the sum of any number of negative terms is negative, it follows that $-a \times +c$, or $+a \times -c$ make or produce $-ac$.

3. When $-a$ is to be multiplied by $-c$: here $-a$ is to be subtracted as often as there are units in c : but subtracting negatives is the same thing as adding affirmatives, by the demonstration of the rule for subtraction; consequently the product is c times a , or $+ac$.

Otherwise. Since $a - a = 0$, therefore $(a - a) \times -c$ is also $= 0$, because 0 multiplied by any quantity, is still but 0 ; and since the first term of the product, or $a \times -c$ is $= -ac$, by the second case; therefore the last term of the product, or $-a \times -c$, must be $+ac$, to make the sum $= 0$, or $-ac + ac = 0$; that is, $-a \times -c = +ac$.

EXAMPLES.

EXAMPLES.

$5a-3c$	$3ac-4b$	$2a^3-3c+5$
$2a$	$3a$	bc
<hr/>	<hr/>	<hr/>
$10a^2-6ac$	$9a^2c-12ab$	$2a^2bc-3bc^2+5bc$
<hr/>	<hr/>	<hr/>
$12x-2ac$	$25c-7b$	$4x-b+3ab$
$4a$	$-2a$	$2ab$
<hr/>	<hr/>	<hr/>
$3c^2+x$	$10x^2-3y^2$	$3a^2-2x^2-6b$
$4xy$	$-4x^2$	$2ax^2$
<hr/>	<hr/>	<hr/>
<hr/>	<hr/>	<hr/>

CASE III.

When both the Factors are Compound Quantities ;

MULTIPLY every term of the multiplier by every term of the multiplicand, separately ; setting down the products one after or under another, with their proper signs ; and add the several lines of products all together for the whole product required.

$a+b$	$3x+2y$	$2x^2+xy-2y^2$
$a+b$	$4x-5y$	$3x-3y$
<hr/>	<hr/>	<hr/>
a^2+ab	$12x^2+8xy$	$6x^3+3x^2y-6xy^2$
$+ab+b^2$	$-15xy-10y^2$	$-6x^2y-3xy^2+6y^3$
<hr/>	<hr/>	<hr/>
$a^2+2ab+b^2$	$12x^2-7xy-10y^2$	$6x^3-3x^2y-9xy^2+6y^3$
<hr/>	<hr/>	<hr/>
$a+b$	x^2+y	a^2+ab+b^2
$a-b$	x^2+y	$a-b$
<hr/>	<hr/>	<hr/>
a^2+ab	x^4+yx^2	$a^3+a^2b+ab^2$
$-ab-b^2$	$+yx^2+y^2$	$-a^2b-ab^2-b^3$
<hr/>	<hr/>	<hr/>
$a^2 * -b^2$	$x^4+2yx+y^2$	$a^3 * * -b^3$
<hr/>	<hr/>	<hr/>

Note.

Note. In the multiplication of compound quantities, it is the best way to set them down in order, according to the powers and the letters of the alphabet. And in multiplying them, begin at the left hand side, and multiply from the left hand towards the right, in the manner that we write, which is contrary to the way of multiplying numbers. But in setting down the several products, as they arise, in the second and following lines, range them under the like terms in the lines above, when there are such like quantities; which is the easiest way for adding them up together.

In many cases, the multiplication of compound quantities is only to be performed by setting them down one after another, each within or under a vinculum with a sign of multiplication between them. As $(a+b) \times (a-b) \times 3ab$, or $\overline{a+b} \cdot \overline{a-b} \cdot 3ab$.

EXAMPLES FOR PRACTICE.

1. Multiply $10ac$ by $2a$. Ans. $20a^2c$.
2. Multiply $3a^2 - 2b$ by $3b$. Ans. $9a^2b - 6b^2$.
3. Multiply $3a + 2b$ by $3a - 2b$. Ans. $9a^2 - 4b^2$.
4. Multiply $x^2 - xy + y^2$ by $x + y$. Ans. $x^3 + y^3$.
5. Multiply $a^3 + a^2b + ab^2 + b^3$ by $a - b$. Ans. $a^4 - b^4$.
6. Multiply $a^2 + ab + b^2$ by $a^2 - ab + b^2$.
7. Multiply $3x^2 - 2xy + 5$ by $x^2 + 2xy - 6$.
8. Multiply $3a^2 - 2ax + 5x^2$ by $3a^2 - 4ax - 7x^2$.
9. Multiply $3x^3 + 2x^2y^2 + 3y^3$ by $2x^3 - 3y^2 + 3y^3$.
10. Multiply $a^2 + ab + b^2$ by $a - 2b$.

DIVISION.

DIVISION in Algebra, like that in numbers, is the converse of multiplication; and it is performed like that of numbers also, by beginning at the left hand side, and dividing all the parts of the dividend by the divisor, when they can be so divided; or else by setting them down like a fraction, the dividend over the divisor, and then abbreviating the fraction as much as can be done. This will naturally divide into the following particular cases.

CASE

CASE I.

When the Divisor and Dividend are both Simple Quantities ;

SET the terms both down as in division of numbers, either the divisor before the dividend, or below it, like the denominator of a fraction. Then abbreviate these terms as much as can be done, by cancelling or striking out all the letters that are common to them both, and also dividing the one co-efficient by the other, or abbreviating them after the manner of a fraction, by dividing them by their common measure.

Note. Like signs in the two factors make + in the quotient; and unlike signs make —; the same as in multiplication*.

EXAMPLES.

1. To divide $6ab$ by $3a$,

Here $6ab \div 3a$ or $3a) 6ab$ (or $\frac{6ab}{3a} = 2b$.

2. Also $c \div c = \frac{c}{c} = 1$; and $abx \div bxy = \frac{abx}{bxy} = \frac{a}{y}$.

3. Divide $16x^2$ by $8x$.

Ans. $2x$.

4. Divide $12a^2x^2$ by $-3a^2x$.

Ans. $-4x$.

5. Divide $-15ay^2$ by $3ay$.

Ans. $-5y$.

6. Divide $-18ax^2y$ by $-8axz$.

Ans. $\frac{9xy}{4z}$.

* Because the divisor multiplied by the quotient, must produce the dividend. Therefore,

1. When both the terms are +, the quotient must be +; because + in the divisor \times + in the quotient, produces + in the dividend.

2. When the terms are both—, the quotient is also +; because — in the divisor \times + in the quotient, produces — in the dividend.

3. When one term is + and the other —, the quotient must be —; because + in the divisor \times — in the quotient produces — in the dividend, or— in the divisor, \times + in the quotient gives—in the dividend.

So that the rule is general; viz, that like signs give +, and unlike signs give —, in the quotient.

CASE II.

When the Dividend is a Compound Quantity, and the Divisor a Simple one :

DIVIDE every term of the dividend by the divisor, as in the former case.

EXAMPLES.

$$1. (ab + b^2) \div 2b, \text{ or } \frac{ab + b^2}{2b} = \frac{a + b}{2} = \frac{1}{2}a + \frac{1}{2}b.$$

$$2. (10ab + 15ax) \div 5a, \text{ or } \frac{10ab + 15ax}{5a} = 2b + 3x.$$

$$3. (30az - 48z) \div z, \text{ or } \frac{30az - 48z}{z} = 30a - 48.$$

$$4. \text{ Divide } 6ab - 8ax + a \text{ by } 2a.$$

$$5. \text{ Divide } 3x^2 - 15 + 6x + 6a \text{ by } 3x.$$

$$6. \text{ Divide } 6abc + 12abx - 9a^2b \text{ by } 3ab.$$

$$7. \text{ Divide } 10a^2x - 15x^2 - 25x \text{ by } 5x.$$

$$8. \text{ Divide } 15a^2bc - 15acx^2 + 5ad^2 \text{ by } -5ac.$$

$$9. \text{ Divide } 15a + 3ay - 18y^2 \text{ by } 21a.$$

$$10. \text{ Divide } -20ab + 60ab^3 \text{ by } -6ab.$$

CASE III.

When the Divisor and Dividend are both Compound Quantities ;

1. SET them down as in common Division of numbers, the divisor before the dividend, with a small curved line between them, and ranging the terms according to the powers of some one of the letters in both, the higher powers before the lower.

2. Divide the first term of the dividend by the first term of the divisor, as in the first case, and set the result in the quotient,

3. Multiply the whole divisor by the term thus found, and subtract the result from the dividend.

4. To this remainder bring down as many terms of the dividend as are requisite for the next operation, dividing as before ; and so on to the end, as in common arithmetic.

Note.

Note. If the divisor be not exactly contained in the dividend, the quantity which remains after the operation is finished, may be placed over the divisor, like a vulgar fraction, and set down at the end of the quotient as in common arithmetic.

EXAMPLES.

$$\begin{array}{r}
 a-b) a^2 - 2ab + b^2 \quad (a-b \\
 \underline{a^2 - ab} \\
 -ab + b^2 \\
 \underline{-ab + b^2} \\
 + b^2
 \end{array}$$

$$\begin{array}{r}
 a-c) a^3 - 4a^2c + 4ac^2 - c^3 \quad (a^2 - 3ac + c^2 \\
 \underline{a^3 - a^2c} \\
 -3a^2c + 4ac^2 \\
 \underline{-3a^2c + 3ac^2} \\
 + ac^2 - c^3 \\
 - c^3 \\
 \underline{ - c^3} \\

 \end{array}$$

$$\begin{array}{r}
 a-2) a^3 - 6a^2 + 12a - 8 \quad (a^2 - 4a + 4 \\
 \underline{a^3 - 2a^2} \\
 -4a^2 + 12a \\
 \underline{-4a^2 + 8a} \\
 + 4a - 8 \\
 \underline{4a - 8} \\
 - 8 \\
 \underline{4a - 8} \\

 \end{array}$$

$$\begin{array}{r}
 a+z) a^2 + z^3 \quad (a^2 - az + z^2 \\
 \underline{a^2 + a^2z} \\
 -a^2z + z^3 \\
 \underline{-a^2z - az^2} \\
 + z^3 \\
 - az^2
 \end{array}$$

$a+x)$

$$\begin{array}{r}
 a+x) a^4 - 3x^4 \left(a^3 - a^2x + ax^2 - x^3 - \frac{2x^4}{a+x} \right) \\
 \underline{a^4 + a^3x} \\
 -a^3x - 3x^4 \\
 \underline{-a^3x - a^2x^2} \\
 a^2x^2 - 3x^4 \\
 \underline{a^2x^2 + ax^3} \\
 -ax^3 - 3x^4 \\
 \underline{-ax^3 - x^4} \\
 -2x^4 \\
 \underline{\hspace{1.5cm}}
 \end{array}$$

EXAMPLES FOR PRACTICE.

1. Divide $a^2 + 4ax + 4x^2$ by $a + 2x$. Ans. $a + 2x$.
2. Divide $a^3 - 3a^2z + 3az^2 - z^3$ by $a - z$. Ans. $a^2 - 2az + z^2$.
3. Divide 1 by $1 + a$, Ans. $1 - a + a^2 - a^3 + \&c$.
4. Divide $12x^4 - 192$ by $3x - 6$. Ans. $4x^3 + 8x^2 + 16x + 32$.
5. Divide $a^5 - 5a^4b + 10a^3b^2 - 10a^2b^3 + 5ab^4 - b^5$ by $a^2 - 2ab + b^2$. Ans. $a^3 - 3a^2b + 3ab^2 - b^3$.
6. Divide $48z^3 - 96az^2 - 64a^2z + 150a^3$ by $2z - 3a$.
7. Divide $b^6 - 3b^4x^2 + 3b^2x^4 - x^6$ by $b^3 - 3b^2x + 3bx^2 - x^3$.
8. Divide $a^7 - x^7$ by $a - x$.
9. Divide $a^3 + 5a^2x + 5ax^2 + x^3$ by $a + x$.
10. Divide $a^4 + 4a^2b^2 - 32b^4$ by $a + 2b$.
11. Divide $24a^4 - b^4$ by $3a - 2b$.

—◆—

ALGEBRAIC FRACTIONS.

ALGEBRAIC FRACTIONS have the same names and rules of operation, as numeral fractions in common arithmetic ; as appears in the following Rules and Cases.

CASE

CASE I.

To reduce a Mixed Quantity to an Improper Fraction.

MULTIPLY the integer by the denominator of the fraction, and to the product add the numerator, or connect it with its proper sign, + or — ; then the denominator being set under this sum, will give the improper fraction required.

EXAMPLES.

1. Reduce $3\frac{4}{5}$, and $a - \frac{b}{x}$ to improper fractions

First, $3\frac{4}{5} = \frac{3 \times 5 + 4}{5} = \frac{15 + 4}{5} = \frac{19}{5}$ the Answer.

And, $a - \frac{b}{x} = \frac{a \times x - b}{x} = \frac{ax - b}{x}$ the Answer.

2. Reduce $a + \frac{a^2}{b}$ and $a - \frac{z^2 + a^2}{a}$ to improper fractions.

First, $a + \frac{a^2}{b} = \frac{a \times b + a^2}{b} = \frac{ab + a^2}{b}$ the Answer.

And, $a - \frac{z^2 + a^2}{a} = \frac{a^2 - z^2 + a^2}{a} = \frac{2a^2 - z^2}{a}$ the Answer.

3. Reduce $5\frac{3}{7}$ to an improper fraction. Ans. $\frac{38}{7}$.

4. Reduce $1 - \frac{3a}{x}$ to an improper fraction. Ans. $\frac{x - 3a}{x}$.

5. Reduce $2a - \frac{3ax + a^2}{4x}$ to an improper fraction.

6. Reduce $12 + \frac{4x - 18}{5x}$ to an improper fraction.

7. Reduce $x + \frac{1 - 3a - c}{c}$ to an improper fraction.

8. Reduce $4 + 2x - \frac{2x^3 - 3a}{5a}$ to an improper fraction.

CASE II.

To Reduce an Improper Fraction to a Whole or Mixed Quantity.

DIVIDE the numerator by the denominator, for the integral part; and set the remainder, if any over the denominator, for the fractional part; the two joined together will be the mixed quantity required.

EXAMPLES.

EXAMPLES.

1. To reduce $\frac{16}{3}$ and $\frac{ab+a^2}{b}$ to mixed quantities.

First, $\frac{16}{3} = 16 \div 3 = 5\frac{1}{3}$, the Answer required.

And, $\frac{ab+a^2}{b} = ab+a^2 \div b = a + \frac{a^2}{b}$. Answer.

2. To reduce $\frac{2ac-3a^2}{c}$ and $\frac{3ax+4x^2}{a+x}$ to mixed quantities.

First, $\frac{2ac-3a^2}{c} = 2ac-3a^2 \div c = 2a - \frac{3a^2}{c}$. Answer.

And, $\frac{3ax+4x^2}{a+x} = 3ax+4x^2 \div a+x = 3x + \frac{x^2}{a+x}$. Ans.

3. Reduce $\frac{33}{5}$ and $\frac{2ax-3x^2}{a}$ to mixed quantities.

Ans. $6\frac{3}{5}$, and $2x - \frac{3x^2}{a}$.

4. Reduce $\frac{4a^2x}{2a}$ and $\frac{2a^2+2b^2}{a-b}$ to whole or mixed quantities.

5. Reduce $\frac{3x^2-3y^2}{x+y}$, and $\frac{2x^3-2y^3}{x-y}$ to whole or mixed quantities.

6. Reduce $\frac{10a^2-4a+6}{5a}$ to a mixed quantity.

7. Reduce $\frac{15a^3+5a^2}{3a^3+2a^2-2a-4}$ to a mixed quantity.

CASE III.

To Reduce Fractions to a Common Denominator.

MULTIPLY every numerator, separately, by all the denominators except its own, for the new numerators; and all the denominators together, for the common denominator.

When the denominators have a common divisor, it will be better, instead of multiplying by the whole denominators, to multiply only by those parts which arise from dividing by the common divisor. And observing also the several rules and directions as in Fractions in the Arithmetic.

EXAMPLES.

EXAMPLES.

1. Reduce $\frac{a}{x}$ and $\frac{b}{z}$ to a common denominator.

Here $\frac{a}{x}$ and $\frac{b}{z} = \frac{az}{xz}$ and $\frac{bx}{xz}$, by multiplying the terms of the first fraction by z , and the terms of the 2d by x .

2. Reduce $\frac{a}{x}$, $\frac{x}{b}$ and $\frac{b}{c}$ to a common denominator.

Here $\frac{a}{x}$, $\frac{x}{b}$, and $\frac{b}{c} = \frac{abc}{bcx}$, $\frac{cx^2}{bcx}$ and $\frac{b^2x}{bcx}$, by multiplying the terms of the 1st fraction by bc , of the second by cx , and of the 3d by bx .

3. Reduce $\frac{2a}{x}$ and $\frac{3b}{2c}$ to a common denominator.

$$\text{Ans. } \frac{4ac}{2cx} \text{ and } \frac{3bx}{2cx}$$

4. Reduce $\frac{2a}{b}$ and $\frac{3a+2b}{2c}$ to a common denominator.

$$\text{Ans. } \frac{4ac}{2bc} \text{ and } \frac{3ab+2b^2}{2bc}$$

5. Reduce $\frac{5a}{3x}$ and $\frac{8b}{2c}$, and $4d$ to a common denominator.

$$\text{Ans. } \frac{10ac}{6cx} \text{ and } \frac{9bx}{6cx} \text{ and } \frac{24cdx}{6cx}$$

6. Reduce $\frac{5}{6}$ and $\frac{3a}{4}$ and $2b + \frac{3a}{b}$, to fractions having a common denominator. Ans. $\frac{20b}{24b}$ and $\frac{18ab}{24b}$, and $\frac{48b^2+72a}{24b}$.

7. Reduce $\frac{1}{3}$ and $\frac{2a^2}{4}$ and $\frac{2a^2+b^2}{a+b}$ to a common denominator.

8. Reduce $\frac{3b}{4a^2}$ and $\frac{2c}{3a}$ and $\frac{d}{2a}$ to a common denominator.

3H - 7 - 11

CASE IV.

To find the Greatest Common Measure of the Terms of a Fraction.

DIVIDE the greater term by the less, and the last divisor by the last remainder, and so on till nothing remains; then the divisor last used will be the common measure required; just the same as in common numbers.

But note, that it is proper to range the quantities according to the dimensions of some letters, as is shown in division. And note also, that all the letters or figures which are common to each term of the divisors, must be thrown out of them, or must divide them, before they are used in the operation.

EXAMPLES.

1. To find the greatest common measure of $\frac{ab+b^2}{ac^2+bc^2}$.

$$\begin{array}{r} ab+b^2) ac^2+bc^2 \\ \text{or } a+b) ac^2+bc^2 \quad (c^2 \\ \hline ac^2+bc^2 \end{array}$$

Therefore the greatest common measure is $a+b$.

2. To find the greatest common measure of $\frac{a^3-ab^2}{a^2+2ab+b^2}$

$$\begin{array}{r} a^2+2ab+b^2) a^3-ab^2 \quad (a \\ \hline a^3+2a^2b+ab^2 \\ \hline -2a^2b-2ab^2) a^2+2ab+b^2 \\ \text{or } a+b) a^2+2ab+b^2 \quad (a+b \\ \hline a^2+ab \\ \hline ab+b^2 \\ \hline ab+b^2 \end{array}$$

Therefore $a+b$ is the greatest common divisor.

3. To find the greatest common divisor of $\frac{a^2-4}{ab+2b}$

Ans. $a-2$.

4. To

4. To find the greatest common divisor of $\frac{a^5 - a^3b^2}{a^4 - b^4}$.

Ans. $a^2 - b^2$.

5. Find the greatest com. measure of $\frac{a^3x + 2a^2x^2 + 2ax^3 + x^4}{5a^5 + 10a^4x + 5a^2x^2}$.

CASE V.

To Reduce a Fraction to its Lowest Terms.

FIND the greatest common measure, as in the last problem. Then divide both the terms of the fraction by the common measure thus found, and it will reduce it to its lowest terms at once, as was required. Or divide the terms by any quantity which it may appear will divide them both, as in arithmetical fractions.

EXAMPLES.

1. Reduce $\frac{ab+b^2}{ac^2+bc^2}$ to its lowest terms.

$$\begin{array}{l} ab+b^2 \\ \text{or } a+b \end{array} \left. \begin{array}{l} ac^2+bc^2 \\ ac^2+bc^2 \\ ac^2+bc^2 \end{array} \right\} (c^2)$$

Here $ab+b^2$ is divided by the common factor b .

Therefore $a+b$ is the greatest common measure, and hence

$a+b) \frac{ab+b^2}{ac^2+bc^2} = \frac{b}{c^2}$, is the fraction required.

2. To reduce $\frac{c^3 - b^2c}{c^2 + 2bc + b^2}$ to its least terms.

$$\begin{array}{l} c^3 - b^2c \\ c^3 + 2bc^2 + b^2c \end{array} \left(\begin{array}{l} c \\ c^2 + 2bc + b^2 \end{array} \right)$$

$$\begin{array}{l} -2bc^2 - 2b^2c \\ \text{or } c+b \end{array} \left(\begin{array}{l} c^2 + 2bc + b^2 \\ c^2 + 2bc + b^2 \\ c^2 + bc \end{array} \right)$$

$$\begin{array}{l} bc + b^2 \\ bc + b^2 \end{array}$$

Therefore

Therefore $c+b$ is the greatest common measure, and hence $c+b$) $\frac{c^3-b^2c}{c^2+2bc+b^2} = \frac{c^2-bc}{c+b}$ is the fraction required.

3. Reduce $\frac{c^3-b^3}{c^4-b^2c^2}$ to its lowest terms. Ans. $\frac{c^2+bc+b^2}{c^3+bc^2}$.

4. Reduce $\frac{a^2-b^2}{a^4-b^4}$ to its lowest terms. Ans. $\frac{1}{a^2+b^2}$.

5. Reduce $\frac{a^4-b^4}{a^3-3a^2b+3ab^2-b^3}$ to its lowest terms.

6. Reduce $\frac{3a^5+6a^4c+3a^3c^2}{a^3c+3a^2c^2+3ac^3+c^4}$ to its lowest terms.

7. Reduce $\frac{a^3-ab^2}{a^2+2ab+b^2}$ to its lowest terms.

CASE VI.

To add Fractional Quantities together.

If the fractions have a common denominator, add all the numerators together; then under their sum set the common denominator, and it is done.

If they have not a common denominator, reduce them to one, and then add them as before.

EXAMPLES.

1. Let $\frac{a}{3}$ and $\frac{a}{4}$ be given, to find their sum.

Here $\frac{a}{3} + \frac{a}{4} = \frac{4a}{12} + \frac{3a}{12} = \frac{7a}{12}$ is the sum required.

2. Given $\frac{a}{b}$, $\frac{b}{c}$, and $\frac{c}{d}$, to find their sum.

Here $\frac{a}{b} + \frac{b}{c} + \frac{c}{d} = \frac{acd}{bcd} + \frac{bbd}{bcd} + \frac{bcc}{bcd} = \frac{acd+bbd+bcc}{bcd}$ the sum required.

3. Let

* 3. Let $a - \frac{3x^2}{b}$ and $b + \frac{2ax}{c}$ be added together.

$$\begin{aligned} \text{Here } a - \frac{3x^2}{b} + b + \frac{2ax}{c} &= a - \frac{3cx^2}{bc} + b + \frac{2abx}{bc} \\ &= a + b + \frac{2abx - 3cx^2}{bc}, \text{ the sum required.} \end{aligned}$$

4. Add $\frac{4x}{3a}$ and $\frac{2x}{5b}$ together.

$$\text{Ans. } \frac{20bx + 6ax}{15ab}$$

5. Add $\frac{a}{3}$, $\frac{a}{4}$ and $\frac{a}{5}$ together.

$$\text{Ans. } \frac{47a}{60}$$

6. Add $\frac{2a-3}{4}$ and $\frac{5a}{8}$ together.

$$\text{Ans. } \frac{9a-6}{8}$$

7. Add $2a + \frac{a+3}{6}$ to $4a + \frac{2a-5}{4}$.

$$\text{Ans. } 6a + \frac{14a-13}{20}$$

8. Add $6a$, and $\frac{3a^2}{4b}$ and $\frac{a+b}{3b}$ together.

9. Add $\frac{5a}{4}$, and $\frac{6a}{5}$ and $\frac{3a+2}{7}$ together.

10. Add $2a$, and $\frac{3a}{8}$ and $3 + \frac{a}{6}$ together.

11. Add $8a + \frac{3a}{4}$ and $2a - \frac{5a}{8}$ together.

CASE VII.

* *To Subtract one Fractional Quantity from another.*

REDUCE the fractions to a common denominator, as in addition, if they have not a common denominator

Subtract the numerators from each other, and under their difference set the common denominator, and the work is done.

* In the addition of mixed quantities, it is best to bring the fractional parts only to a common denominator, and to annex their sum to the sum of the integers, with the proper sign. And the same rule may be observed for mixed quantities in subtraction also.

EXAMPLES.

EXAMPLES.

1. To find the difference of $\frac{3a}{4}$ and $\frac{4a}{7}$.

Here $\frac{3a}{4} - \frac{4a}{7} = \frac{21a}{28} - \frac{16a}{28} = \frac{5a}{28}$ is the difference required.

2. To find the difference of $\frac{2a-b}{4c}$ and $\frac{3a-4b}{3b}$.

Here $\frac{2a-b}{4c} - \frac{3a-4b}{3b} = \frac{6ab-3bb}{12bc} - \frac{12ac-16bc}{12bc} = \frac{6ab-3bb-12ac+16bc}{12bc}$ is the difference required.

3. Required the difference of $\frac{10a}{9}$ and $\frac{4a}{7}$.

4. Required the difference of $6a$ and $\frac{3a}{4}$.

5. Required the difference of $\frac{5a}{4}$ and $\frac{2a}{3}$.

6. Subtract $\frac{2b}{c}$ from $\frac{3a+c}{b}$.

7. Take $\frac{2a+6}{9}$ from $\frac{4a+8}{5}$.

8. Take $2a - \frac{a-3b}{c}$ from $4a + \frac{2a}{c}$.

CASE VIII.

To Multiply Fractional Quantities together.

MULTIPLY the numerators together for a new numerator, and the denominators for a new denominator.*

* 1 When the numerator of one fraction, and the denominator of the other, can be divided by some quantity, which is common to both, the quotients may be used instead of them.

2. When a fraction is to be multiplied by an integer, the product is found either by multiplying the numerator, or dividing the denominator by it; and if the integer be the same with the denominator, the numerator may be taken for the product.

EXAMPLES.

EXAMPLES.

1. Required to find the product of $\frac{a}{8}$ and $\frac{2a}{5}$.

Here $\frac{a \times 2a}{8 \times 5} = \frac{2a^2}{40} = \frac{a^2}{20}$ the product required.

2. Required the product of $\frac{a}{3}$, $\frac{3a}{4}$, and $\frac{6a}{7}$.

$\frac{a \times 3a \times 6a}{3 \times 4 \times 7} = \frac{18a^3}{84} = \frac{3a^3}{14}$ the product required.

3. Required the product of $\frac{2a}{b}$ and $\frac{a+b}{2a+c}$.

Here $\frac{2a \times (a+b)}{b \times (2a+c)} = \frac{2aa+2ab}{2ab+bc}$ the product required.

4. Required the product of $\frac{4a}{3}$ and $\frac{6a}{5c}$.

5. Required the product of $\frac{3a}{4}$ and $\frac{4b^2}{3a}$.

6. To multiply $\frac{3a}{b}$ and $\frac{8ac}{b}$ and $\frac{4ab}{3c}$ together.

7. Required the product of $2a + \frac{ab}{2c}$ and $\frac{3a^2}{b}$.

8. Required the product of $\frac{2a^2 - 2b^2}{3bc}$ and $\frac{4a^2 + 2b^2}{a+b}$.

9. Required the product of $3a$, and $\frac{2a+1}{a}$, and $\frac{2a-1}{2a+b}$.

10. Multiply $a + \frac{x}{2a} - \frac{x^2}{4a^2}$ by $x - \frac{a}{2x} + \frac{a^2}{4x^2}$.

CASE IX.

To Divide one Fractional Quantity by another.

DIVIDE the numerators by each other, and the denominators by each other, if they will exactly divide. But, if not, then invert the terms of the divisor, and multiply by it exactly as in multiplication.*

EXAMPLES.

* 1. If the fractions to be divided have a common denominator, take the numerator of the dividend for a new numerator, and the numerator of the divisor for the new denominator.

2. When a fraction is to be divided by any quantity, it is the same thing whether the numerator be divided by it, or the denominator multiplied by it.

3. When

EXAMPLES.

1. Required to divide $\frac{a}{4}$ by $\frac{3a}{8}$

Here $\frac{a}{4} \div \frac{3a}{8} = \frac{a}{4} \times \frac{8}{3a} = \frac{8a}{12a} = \frac{2}{3}$ the quotient.

2. Required to divide $\frac{3a}{2b}$ by $\frac{5c}{4d}$.

Here $\frac{3a}{2b} \div \frac{5c}{4d} = \frac{3a}{2b} \times \frac{4d}{5c} = \frac{12ad}{10bc} = \frac{6ad}{5bc}$ the quotient.

3. To divide $\frac{2a+b}{3a-2b}$ by $\frac{3a+2b}{4a+b}$. Here,

$\frac{2a+b}{3a-2b} \times \frac{4a+b}{3a+2b} = \frac{8a^2+6ab+b^2}{9a^2-4b^2}$ the quotient required.

4. To divide $\frac{3a^2}{a^2+b^2}$ by $\frac{2a}{2a+2b}$.

Here, $\frac{3a^2}{a^2+b^2} \times \frac{a+b}{a} = \frac{3a^2 \times (a+b)}{(a^2+b^2) \times a} = \frac{3a}{a^2-ab+b^2}$ is the quotient required.

5. To divide $\frac{3x}{4}$ by $\frac{11}{12}$.

6. To divide $\frac{6x^2}{5}$ by $3x$.

7. To divide $\frac{3x+1}{9}$ by $\frac{4x}{3}$.

8. To divide $\frac{4x}{2x-1}$ by $\frac{x}{3}$.

9. To divide $\frac{4x}{5}$ by $\frac{3a}{5b}$.

10. To divide $\frac{2a-b}{4cd}$ by $\frac{5ac}{6d}$.

11. Divide $\frac{5a^4-5b^4}{2a^2-4ab+2b^2}$ by $\frac{6a^2+5ab}{4a-4b}$.

INVOLU-

3. When the two numerators, or the two denominators, can be divided by some common quantity, let that be done, and the quotients used instead of the fractions first proposed.

INVOLUTION.

INVOLUTION is the raising of powers from any proposed root ; such as finding the square, cube, biquadrate, &c. of any given quantity. The method is as follows ;

* MULTIPLY the root or given quantity by itself, as many times as there are units in the index less one, and the last product will be the power required.—Or, in literals, multiply the index of the root by the index of the power, and the result will be the power, the same as before,

Note. When the sign of the root is +, all the powers of it will be + ; but when the sign is —, all the even powers will be +, and all the odd powers — ; as is evident from multiplication.

EXAMPLES.

a , the root
 $a^2 =$ square.
 $a^3 =$ cube
 $a^4 =$ 4th power
 $a^5 =$ 5th power
 &c.

— $2a$, the root
 + $4a^2 =$ square
 — $8a^3 =$ cube
 + $16a^4 =$ 4th power
 — $32a^5 =$ 5th power

— $\frac{2ax^2}{3b}$, the root
 + $\frac{4a^2x^4}{9b^2} =$ square
 — $\frac{8a^3x^6}{27b^3} =$ cube
 + $\frac{16a^4x^8}{81b^4} =$ 4th power.

a^2 , the root
 $a^4 =$ square
 $a^6 =$ cube
 $a^8 =$ 4th power
 $a^{10} =$ 5th power
 &c.

— $3ab^2$, the root
 + $9a^2b^4 =$ square
 — $27a^3b^6 =$ cube
 + $81a^4b^8 =$ 4th power
 — $243a^5b^{10} =$ 5th power

$\frac{a}{2b}$, the root
 $\frac{a^2}{4b^2} =$ square
 $\frac{a^3}{8b^3} =$ cube
 $\frac{a^4}{16b^4} =$ biquadrate

* Any power of the product of two or more quantities, is equal to the same power of each of the factors, multiplied together.

And any power of a fraction, is equal to the same power of the numerator, divided by the like power of the denominator.

Also, powers or roots of the same quantity, are multiplied by one another, by adding their exponents ; or divided, by subtracting their exponents.

Thus $a^3 \times a^2 = a^{3+2} = a^5$. And $a^3 \div a^2$ or $\frac{a^3}{a^2} = a^{3-2} = a$.

$$x - a = \text{root}$$

$$x - a$$

$$x^2 - ax$$

$$-ax + a^2$$

$$x^2 - 2ax + a^2 \text{ square}$$

$$x - a$$

$$x^3 - 2ax^2 + a^2x$$

$$-ax^2 + 2a^2x - a^3$$

$$x^3 - 3ax^2 + 3a^2x - a^3$$

$$x + a = \text{root}$$

$$x + a$$

$$x^2 + ax$$

$$+ax + a^2$$

$$x^2 + 2ax + a^2$$

$$x + a$$

$$x^3 + 2ax^2 + a^2x$$

$$+ax^2 + 2a^2x + a^3$$

$$x^3 + 3ax^2 + 3a^2x + a^3$$

(the cubes, or third powers, of $x - a$ and $x + a$.)

EXAMPLES FOR PRACTICE.

1. Required the cube or third power of $3a^2$.
2. Required the 4th power of $2a^2b$.
3. Required the 3d power of $-4a^2b^3$.
4. To find the biquadrate of $-\frac{a^2x}{2b^2}$.
5. Required the 5th power of $a - 2x$.
6. To find the 6th power of $2a^{\frac{1}{2}}$.

SIR ISAAC NEWTON'S RULE for raising a Binomial to any Power whatever.*

1. To find the terms without the Co-efficients. The index of the first, or leading quantity, begins with the index of the given power, and in the succeeding terms decreases continually by 1, in every term to the last; and in the 2d or following quantity, the indices of the terms are 0, 1, 2, 3, 4, &c. increasing always by 1. That is, the first term will contain only the first part of the root with the same index, or of

* This rule expressed in general terms is as follows;

$$(a+x)^n = a^n + n \cdot a^{n-1}x + n \cdot \frac{n-1}{2} a^{n-2}x^2 + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} a^{n-3}x^3 \text{ \&c.}$$

$$(a-x)^n = a^n - n \cdot a^{n-1}x + n \cdot \frac{n-1}{2} a^{n-2}x^2 - n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} a^{n-3}x^3 \text{ \&c.}$$

Note. The sum of the co-efficients, in every power, is equal to the number 2, when raised to that power. Thus $1 + 1 = 2$ in the first power; $1 + 2 + 1 = 4 = 2^2$ in the square; $1 + 3 + 3 + 1 = 8 = 2^3$ in the cube, or third power; and so on.

the

the same height as the intended power : and the last term of the series will contain only the 2d part of the given root, when raised also to the same height of the intended power ; but all the other or intermediate terms will contain the products of some powers of both the members of the root, in such sort, that the powers or indices of the 1st or leading member will always decrease by 1, while those of the 2d member always increase by 1.

2. *To find the Co-efficients.* The first co-efficient is always 1, and the second is the same as the index of the intended power ; to find the third co-efficient, multiply that of the 2d term by the index of the leading letter in the same term, and divide the product by 2 ; and so on, that is, multiply the co-efficient of the term last found by the index of the leading quantity in that term, and divide the product by the number of terms to that place, and it will give the co-efficient of the term next following ; which rule will find all the co-efficients, one after another.

Note. The whole number of terms will be 1 more than the index of the given power : and when both terms of the root are +, all the terms of the power will be + ; but if the second term be -, all the odd terms will be +, and all the even terms -, which causes the terms to be + and - alternately. Also the sum of the two indices, in each term, is always the same number, viz. the index of the required power : and counting from the middle of the series, both ways, or towards the right and left, the indices of the two terms are the same figures at equal distances, but mutually changed places. Moreover, the co-efficients are the same numbers at equal distances from the middle of the series, towards the right and left ; so by whatever numbers they increase to the middle, by the same in the reverse order they decrease to the end.

EXAMPLES.

1. Let $a+x$ be involved to the 5th power.

The terms without the co-efficients, by the 1st rule, will be

$$a^5, a^4x, a^3x^2, a^2x^3, ax^4, x^5,$$

and the co-efficients, by the 2d rule, will be

$$1, 5, \frac{5 \times 4}{2}, \frac{10 \times 3}{3}, \frac{10 \times 2}{4}, \frac{5 \times 1}{5} ;$$

$$\text{or } 1, 5, 10, 10, 5, 1 ;$$

Therefore the 5th power altogether is

$$a^5 + 5a^4x + 10a^3x^2 + 10a^2x^3 + 5ax^4 + x^5.$$

But it is best to set down both the co-efficients and the powers of the letters at once, in one line, without the intermediate lines in the above example, as in the example here below.

2. Let $a-x$ be involved to the 6th power.

The terms with the co-efficients will be

$$a^6 - 6a^5x + 15a^4x^2 - 20a^3x^3 + 15a^2x^4 - 6ax^5 + x^6.$$

3. Required the 4th power of $a-x$.

$$\text{Ans. } a^4 - 4a^3x + 6a^2x^2 - 4ax^3 + x^4.$$

And thus any other powers may be set down at once, in the same manner; which is the best way.

EVOLUTION.

EVOLUTION is the reverse of Involution, being the method of finding the square root, cube root, &c. of any given quantity whether simple or compound.

CASE I.

To find the Roots of Simple Quantities.

EXTRACT the root of the co-efficient for the numeral part; and divide the index of the letter or letters, by the index of the power, and it will give the root of the literal part; then annex this to the former, for the whole root sought.*

* Any even root of an affirmative quantity, may be either $+$ or $-$: thus the square root of $+a^2$ is either $+a$, or $-a$; because $+a \times +a = +a^2$, and $-a \times -a = +a^2$ also.

But an odd root of any quantity will have the same sign as the quantity itself: thus the cube root of $+a^3$ is $+a$ and the cube root of a^3 is $-a$; for $+a \times +a \times +a = +a^3$, and $-a \times -a \times -a = -a^3$.

Any even root of a negative quantity is impossible: for neither $+a \times +a$, nor $-a \times -a$ can produce $-a^2$.

Any root of a product, is equal to the like root of each of the factors multiplied together. And for the root of a fraction, take the root of the numerator, and the root of the denominator.

EXAMPLES.

EXAMPLES.

1. The square root of $4a^2$, is $2a$.
2. The cube root of $8a^3$, is $2a^{\frac{3}{3}}$ or $2a$.
3. The square root of $\frac{5a^2b^2}{9c^2}$, or $\sqrt{\frac{5a^2b^2}{9c^2}}$, is $\frac{ab}{3c} \sqrt{5}$.
4. The cube of $-\frac{16a^4b^6}{27c^3}$, is $-\frac{2ab^2}{3c} \sqrt[3]{2a}$.
5. To find the square root of $2a^2b^4$. Ans. $ab^2 \sqrt{2}$.
6. To find the cube root of $-64a^3b^6$. Ans. $-4ab^2$.
7. To find the square root of $\frac{8a^2b^2}{3c^3}$. Ans. $\frac{2ab}{c} \sqrt{\frac{2}{3c}}$.
8. To find the 4th root of $81a^4b^6$. Ans. $3ab \sqrt[4]{b}$.
9. To find the 5th root of $-32a^5b^6$. Ans. $-2ab \sqrt[5]{b}$.

CASE II.

To find the Square root of a Compound Quantity:

THIS is performed like as in numbers, thus :

1. Range the quantities according to the dimensions of one of the letters, and set the root of the first term in the quotient.

2. Subtract the square of the root, thus found, from the first term, and bring down the next two terms to the remainder for a dividend ; and take double the root for a divisor.

3. Divide the dividend by the divisor, and annex the result both to the quotient and to the divisor.

4. Multiply the divisor thus increased, by the term last set in the quotient, and subtract the product from the dividend.

And so on, always the same, as in common arithmetic.

EXAMPLES.

1. Extract the square root of $a^4 - 4a^3b + 6a^2b^2 - 4ab^3 + b^4$.
 $a^4 - 4a^3b + 4a^2b^2 - 4ab^3 + b^4$ ($a^2 - 2ab + b^2$ the root.)

$$\begin{array}{r} \hline 2a^2 - 2ab) - 4a^3b + 6a^2b^2 \\ \quad - 4a^3b + 4a^2b^2 \\ \hline \end{array}$$

$$\begin{array}{r} \hline 2a^2 - 4ab + b^2) 2a^2b^2 - 4ab^3 + b^4 \\ \quad 2a^2b^2 - 4ab^3 + b^4 \\ \hline \end{array}$$

2. Find

2. Find the root of $a^4 + 4a^3b + 10a^2b^2 + 12ab^3 + b^4$.
 $a^4 + 4a^3b + 10a^2b^2 + 12ab^3 + 9b^4 (a^2 + 2ab + 3b^2)$
 a^4

$$\begin{array}{r} 2a^2 + 2ab \) \ 4a^3b + 10a^2b^2 \\ \underline{4a^3b + 4a^2b^2} \end{array}$$

$$\begin{array}{r} 2a^2 + 4ab + 3b^2 \) \ 6a^2b^2 + 12ab^3 + 9b^4 \\ \underline{6a^2b^2 + 12ab^3 + 9b^4} \end{array}$$

3. To find the square root of $a^4 + 4a^3 + 6a^2 + 4a + 1$.

$$\text{Ans. } a^2 + 2a + 1.$$

4. Extract the square root of $a^4 - 2a^3 + 2a^2 - a + \frac{1}{4}$.

$$\text{Ans. } x^2 - x + \frac{1}{2}.$$

5. It is required to find the square root of $a^2 - ab$.

$$\text{Ans. } a - \frac{b}{2} - \frac{b^2}{8a} - \frac{b^3}{16a^2} - \&c.$$

CASE III.

To find the Roots of any Powers in General.

THIS is also done like the same roots in numbers, thus :

Find the root of the first term, and set it in the quotient, —Subtract its power from that term, and bring down the second term for a dividend.—Involve the root, last found, to the next lower power, and multiply it by the index of the given power, for a divisor.—Divide the dividend by the divisor, and set the quotient as the next term of the root.—Involve now the whole root to the power to be extracted ; then subtract the power thus arising from the given power, and divide the first term of the remainder by the divisor first found ; and so on till the whole is finished.*

EXAMPLES.

* As this method, in high powers, may be thought too laborious, it will not be improper to observe, that the roots of compound quantities may sometimes be easily discovered, thus :

Extract the roots of some of the most simple terms, and connect them together by the sign + or —, as may be judged most suitable for the purpose.—Involve the compound root, thus found, to the proper power ; then, if this be the same with the given quantity, it is the root required.—But if it be found to differ only in some of the signs, change them from + to —, or from — to +, till its power agrees with the given one throughout.

Thus

EXAMPLES.

1. To find the square root of $a^4 - 2a^3b + 3a^2b^2 - 2ab^3 + b^4$.

$$\begin{array}{r}
 a^4 - 2a^3b + 3a^2b^2 - 2ab^3 + b^4 \\
 \underline{a^4} \\
 2a^3) - 2a^3b \\
 \underline{2a^3} \\
 a^4 - 2a^3b + a^2b^2 =) a^2 - ab)^2 \\
 \underline{2a^2} \\
 2a^2) 2a^2b^2 \\
 \underline{2a^2} \\
 a^4 - 2a^3b + 3a^2b^2 - 2ab^3 + b^4 = (a^2 - ab + b^2)^2.
 \end{array}$$

2. Find the cube root of $a^6 - 6a^5 + 21a^4 - 44a^3 + 63a^2 - 54a + 27$.

$$a^6 - 6a^5 + 21a^4 - 44a^3 + 63a^2 - 54a + 27 = (a^2 - 2a + 3)^3.$$

$$3a^4) - 6a^5$$

$$a^6 - 6a^5 + 12a^4 - 2a^3 =) a^2 - 2a)^3$$

$$3a^4) + 9a^4$$

$$a^6 - 6a^5 + 21a^4 - 44a^3 + 63a^2 - 54a + 27 = (a^2 - 2a + 3)^3.$$

3. To find the square root of $a^2 - 2ab + 2ax + b^2 - 2bx + x^2$.

$$\text{Ans. } a - b + x.$$

4. Find the cube root of $a^6 - 3a^5 + 9a^4 - 13a^3 + 18a^2 - 12a + 8$.

$$\text{Ans. } a^2 - a + 2.$$

5. Find the 4th root of $81a^4 - 216a^3b + 216a^2b^2 - 96ab^3 + 16b^4$.

$$\text{Ans. } 3a - 2b.$$

6. Find the 5th root of $a^5 - 10a^4 + 40a^3 - 80a^2 + 80a - 32$.

$$\text{Ans. } a - 2.$$

7. Required the square root of $1 - x^2$.

8. Required the cube root of $1 - x^3$.

Thus, in the 5th example, the root $3a - 2b$, is the difference of the roots of the first and last terms; and in the third example, the root $a - b + x$, is the sum of the roots of the 1st, 4th, and 6th terms. The same may also be observed of the 6th example, where the root is found from the first and last terms.

SURDS.

SURDS.

SURDS are such quantities as have not exact values in numbers; and are usually expressed by fractional indices, or by means of the radical sign $\sqrt{\quad}$. Thus, $3^{\frac{1}{2}}$, or $\sqrt{3}$, denotes the square root of 3; and $2^{\frac{2}{3}}$ or $\sqrt[3]{2^2}$, or $\sqrt[3]{4}$, the cube root of the square of 2; where the numerator shows the power to which the quantity is to be raised, and the denominator its root.

PROBLEM I.

To Reduce a Rational Quantity to the Form of a Surd.

RAISE the given quantity to the power denoted by the index of the surd: then over or before this new quantity set the radical sign, and it will be the form required.

EXAMPLES.

1. To reduce 4 to the form of the square root.
First, $4^2 = 4 \times 4 = 16$; then $\sqrt{16}$ is the answer.
2. To reduce $3a^2$ to the form of the cube root.
First $3a^2 \times 3a^2 \times 3a^2 = (3a^2)^3 = 27a^6$;
then $\sqrt[3]{27a^6}$ or $(27a^6)^{\frac{1}{3}}$ is the answer.
3. Reduce 6 to the form of the cube root.
Ans. $(216)^{\frac{1}{3}}$ or $\sqrt[3]{216}$.
4. Reduce $\frac{1}{3}ab$ to the form of the square root.
Ans. $\sqrt{\frac{1}{9}a^2b^2}$.
5. Reduce 2 to the form of the 4th root. Ans. $(16)^{\frac{1}{4}}$
6. Reduce $a^{\frac{1}{3}}$ to the form of the 5th root.
7. Reduce $a+x$ to the form of the square root.
8. Reduce $a-x$ to the form of the cube root.

PROBLEM II.

To Reduce Quantities to a Common Index.

1. **REDUCE** the indices of the given quantities to a common denominator, and involve each of them to the power denoted by its numerator; then I set over the common denominator will form the common index. Or,

2. If

2. If the common index be given, divide the indices of the quantities by the given index, and the quotients will be the new indices for those quantities. Then over the said quantities, with their new indices, set the given index, and they will make the equivalent quantities sought.

EXAMPLES.

1. Reduce $3^{\frac{1}{2}}$ and $5^{\frac{1}{3}}$ to a common index.

Here $\frac{1}{2}$ and $\frac{1}{3} = \frac{5}{10}$ and $\frac{2}{10}$.

Therefore $3^{\frac{5}{10}}$ and $5^{\frac{2}{10}} = (3^5)^{\frac{1}{10}}$ and $(5^2)^{\frac{1}{10}} = \sqrt[10]{3^5}$ and $\sqrt[10]{5^2}$
 $5^2 = \sqrt[10]{243}$ and $\sqrt[10]{25}$.

2. Reduce a^3 and $b^{\frac{1}{3}}$ to the same common index $\frac{1}{3}$.

Here, $\frac{3}{1} \div \frac{1}{3} = \frac{3}{1} \times \frac{3}{1} = \frac{9}{1}$ the 1st index,

and $\frac{1}{3} \div \frac{1}{3} = \frac{1}{3} \times \frac{3}{1} = \frac{1}{1}$ the 2d index.

Therefore $(a^9)^{\frac{1}{3}}$ and $(b^{\frac{1}{3}})^{\frac{1}{1}}$, or $\sqrt[3]{a^9}$ and $\sqrt[3]{b^{\frac{1}{3}}}$ are the quantities.

3. Reduce $4^{\frac{1}{3}}$ and $5^{\frac{1}{2}}$ to the common index $\frac{1}{6}$.

Ans. $256^{\frac{1}{6}}$ and $25^{\frac{1}{6}}$.

4. Reduce $a^{\frac{1}{3}}$ and $x^{\frac{1}{4}}$ to the common index $\frac{1}{12}$.

Ans. $(a^4)^{\frac{1}{12}}$ and $(x^3)^{\frac{1}{12}}$.

5. Reduce a^2 and x^3 to the same radical sign.

Ans. $\sqrt{a^4}$ and $\sqrt{x^6}$.

6. Reduce $(a+x)^{\frac{1}{3}}$ and $(a-x)^{\frac{1}{2}}$ to a common index.

7. Reduce $(a+b)^{\frac{1}{2}}$ and $(a-b)^{\frac{1}{4}}$ to a common index.

PROBLEM III.

To Reduce Surds to more Simple Terms.

FIND out the greatest power contained in, or to divide the given surd; take its root, and set it before the quotient or the remaining quantities, with the proper radical sign between them.

EXAMPLES.

1. To reduce $\sqrt{32}$ to simpler terms.

Here $\sqrt{32} = \sqrt{16 \times 2} = \sqrt{16} \times \sqrt{2} = 4 \times \sqrt{2} = 4\sqrt{2}$.

2. To reduce $\sqrt[3]{320}$ to simpler terms.

$\sqrt[3]{320} = \sqrt[3]{64 \times 5} = \sqrt[3]{64} \times \sqrt[3]{5} = 4 \times \sqrt[3]{5} = 4\sqrt[3]{5}$.

3. Reduce

3. Reduce $\sqrt{75}$ to its simplest terms. Ans. $5\sqrt{3}$.
 4. Reduce $\sqrt{\frac{44}{75}}$ to simpler terms. Ans. $\frac{2}{15}\sqrt{33}$.
 5. Reduce $\sqrt[3]{189}$ to its simplest terms. Ans. $3\sqrt[3]{7}$.
 6. Reduce $\sqrt[3]{\frac{135}{2}}$ to its simplest terms. Ans. $\frac{3}{2}\sqrt[3]{10}$.
 7. Reduce $\sqrt{75a^2b}$ to its simplest terms. Ans. $5a\sqrt{3b}$.

Note. There are other cases of reducing algebraic surds to simpler forms, that are practised on several occasions; one instance of which, on account of its simplicity and usefulness, may be here noticed, viz. in fractional forms having compound surds in the denominator, multiply both numerator and denominator by the same terms of the denominator, but having one sign changed, from + to - or from - to +, which will reduce the fraction to a rational denominator.

Ex. To reduce $\frac{\sqrt{20} + \sqrt{12}}{\sqrt{5} - \sqrt{3}}$, multiply it by $\frac{\sqrt{5} + \sqrt{3}}{\sqrt{5} + \sqrt{3}}$, and it becomes $\frac{16 + 4\sqrt{15}}{2} = 8 + 2\sqrt{15}$. Also, if $\frac{3\sqrt{15} - 4\sqrt{5}}{\sqrt{15} + \sqrt{5}}$; multiply it by $\frac{\sqrt{15} - \sqrt{5}}{\sqrt{15} - \sqrt{5}}$, and it becomes $\frac{65 - 7\sqrt{75}}{15 - 5} = \frac{65 - 35\sqrt{3}}{10} = \frac{13 - 7\sqrt{3}}{2}$.

PROBLEM IV.

To add Surd Quantities together.

1. BRING all fractions to a common denominator, and reduce the quantities to their simplest terms, as in the last problem.—2. Reduce also such quantities as have unlike indices to other equivalent ones, having a common index.—3. Then, if the surd part be the same in them all, annex it to the sum of the rational parts, with the sign of multiplication, and it will give the total sum required.

But if the surd part be not the same in all the quantities, they can only be added by the signs + and -.

EXAMPLES.

1. Required to add $\sqrt{18}$ and $\sqrt{32}$ together.

First, $\sqrt{18} = \sqrt{9 \times 2} = 3\sqrt{2}$; and $\sqrt{32} = \sqrt{16 \times 2} = 4\sqrt{2}$:
 Then, $3\sqrt{2} + 4\sqrt{2} = (3 + 4)\sqrt{2} = 7\sqrt{2} =$ sum required.

2. It is required to add $\sqrt[3]{375}$, and $\sqrt[3]{192}$ together.

First, $\sqrt[3]{375} = \sqrt[3]{125 \times 3} = 5\sqrt[3]{3}$; and $\sqrt[3]{192} = \sqrt[3]{64 \times 3} = 4\sqrt[3]{3}$:
 Then, $5\sqrt[3]{3} + 4\sqrt[3]{3} = (5 + 4)\sqrt[3]{3} = 9\sqrt[3]{3} =$ sum required.

3. Required

3. Required the sum of $\sqrt{27}$ and $\sqrt{48}$. Ans. $7\sqrt{3}$.
 4. Required the sum of $\sqrt{50}$ and $\sqrt{72}$. Ans. $11\sqrt{2}$.
 5. Required the sum of $\sqrt{\frac{2}{3}}$ and $\sqrt{\frac{1}{15}}$.
 Ans. $4\sqrt{\frac{1}{15}}$ or $\frac{4}{\sqrt{15}}\sqrt{15}$.
 6. Required the sum of $\sqrt[3]{56}$ and $\sqrt[3]{189}$. Ans. $5\sqrt[3]{7}$.
 7. Required the sum of $\sqrt[3]{\frac{1}{4}}$ and $\sqrt[3]{\frac{1}{32}}$. Ans. $\frac{3}{4}\sqrt[3]{2}$.
 8. Required the sum of $3\sqrt{a^2b}$ and $5\sqrt{16a^2b}$.

PROBLEM V.

To find the Difference of Surd Quantities.

PREPARE the quantities the same way as in the last rule ; then subtract the rational parts, and to the remainder annex the common surd, for the difference of the surds required.

But if the quantities have no common surd, they can only be subtracted by means of the sign—.

EXAMPLES.

1. To find the difference between $\sqrt{320}$ and $\sqrt{80}$.

First, $\sqrt{320} = \sqrt{64 \times 5} = 8\sqrt{5}$; and $\sqrt{80} = \sqrt{16 \times 5} = 4\sqrt{5}$.
 Then $8\sqrt{5} - 4\sqrt{5} = 4\sqrt{5}$ the difference sought.

2. To find the difference between $\sqrt[3]{128}$ and $\sqrt[3]{54}$.

First, $\sqrt[3]{128} = \sqrt[3]{64 \times 2} = 4\sqrt[3]{2}$; and $\sqrt[3]{54} = \sqrt[3]{27 \times 2} = 3\sqrt[3]{2}$.
 Then $4\sqrt[3]{2} - 3\sqrt[3]{2} = \sqrt[3]{2}$, the difference required.

3. Required the difference of $\sqrt{75}$ and $\sqrt{48}$. Ans. $\sqrt{3}$.

4. Required the difference of $\sqrt[3]{256}$ and $\sqrt[3]{32}$. Ans. $2\sqrt[3]{4}$.

5. Required the difference of $\sqrt{\frac{2}{3}}$ and $\sqrt{\frac{5}{9}}$. Ans. $\frac{1}{3}\sqrt{\frac{2}{3}}$.

6. Required the difference of $\sqrt[3]{\frac{2}{3}}$ and $\sqrt[3]{\frac{25}{9}}$. Ans. $\frac{2}{3}\sqrt[3]{\frac{2}{3}}$.

7. Find the difference of $\sqrt{24a^2b^2}$ and $\sqrt{54b^4}$.
 Ans. $(2ab - 3b^2)\sqrt{6}$, or $(3b^2 - 2ab)\sqrt{6}$.

PROBLEM VI.

To Multiply Surd Quantities together.

REDUCE the surds to the same index, if necessary ; next multiply the rational quantities together, and the surds together ; then annex the one product to the other for the whole product required ; which may be reduced to more simple terms if necessary.

EXAMPLES

EXAMPLES.

1. Required to find the product of $4\sqrt{12}$, and $3\sqrt{2}$.
Here, $4 \times 3 \times \sqrt{12} \times \sqrt{2} = 12\sqrt{12} \times 2 = 12\sqrt{24} = 12\sqrt{4 \times 6}$
 $= 12 \times 2 \times \sqrt{6} = 24\sqrt{6}$, the product required.
2. Required to multiply $\frac{1}{4}\sqrt[3]{\frac{3}{4}}$ by $\frac{1}{3}\sqrt[3]{\frac{3}{8}}$.
Here $\frac{1}{4} \times \frac{1}{3} \sqrt[3]{\frac{3}{4}} \times \sqrt[3]{\frac{3}{8}} = \frac{1}{12} \times \sqrt[3]{\frac{9}{32}} = \frac{1}{12} \times \sqrt[3]{\frac{18}{64}} = \frac{1}{12} \times \frac{1}{4} \times \sqrt[3]{18}$
 $= \frac{1}{48} \sqrt[3]{18}$, the product required.
3. Required the product of $3\sqrt{2}$ and $2\sqrt{8}$. Ans. 24.
4. Required the product of $\frac{1}{3}\sqrt[3]{4}$ and $\frac{3}{4}\sqrt[3]{12}$. Ans. $\frac{1}{2}\sqrt[3]{6}$.
5. To find the product of $\frac{5}{3}\sqrt{\frac{3}{8}}$ and $\frac{9}{10}\sqrt{\frac{2}{5}}$. Ans. $\frac{3}{20}\sqrt{15}$.
6. Required the product of $2\sqrt[3]{14}$ and $3\sqrt[3]{4}$. Ans. $12\sqrt[3]{7}$.
7. Required the product of $2a^{\frac{2}{3}}$ and $a^{\frac{4}{3}}$. Ans. $2a$.
8. Required the product of $(a+b)^{\frac{1}{3}}$ and $(a+b)^{\frac{3}{4}}$.
9. Required the product of $2x + \sqrt{b}$ and $2x - \sqrt{b}$.
10. Required the product of $(a+2\sqrt{b})^{\frac{1}{2}}$ and $(a-2\sqrt{b})^{\frac{1}{2}}$.
11. Required the product of $2x^{\frac{1}{n}}$ and $3x^{\frac{1}{m}}$.
12. Required the product of $4x^{\frac{1}{n}}$ and $2y^{\frac{1}{n}}$.

PROBLEM VII.

To Divide one Surd Quantity by another.

REDUCE the surds to the same index, if necessary; then take the quotient of the rational quantities, and annex it to the quotient of the surds, and it will give the whole quotient required; which may be reduced to more simple terms if requisite.

EXAMPLES.

1. Required to divide $6\sqrt{96}$ by $3\sqrt{8}$.
Here $6 \div 3 = 2$, $\sqrt{(96 \div 8)} = 2\sqrt{12} = 2\sqrt{(4 \times 3)} = 2 \times 2\sqrt{3} = 4\sqrt{3}$, the quotient required.
2. Required to divide $12\sqrt[3]{280}$ by $3\sqrt[3]{5}$.
Here $12 \div 3 = 4$, and $280 \div 5 = 56 = 8 \times 7 = 2^3 \cdot 7$.
Therefore $4 \times 2 \times \sqrt[3]{7} = 8\sqrt[3]{7}$, is the quotient required.

3. Let

3. Let $4\sqrt{50}$ be divided by $2\sqrt{5}$. Ans. $2\sqrt{10}$.
 4. Let $6\sqrt[3]{100}$ be divided by $3\sqrt[3]{5}$. Ans. $2\sqrt[3]{20}$.
 5. Let $\frac{5}{8}\sqrt{\frac{1}{50}}$ be divided by $\frac{3}{4}\sqrt{\frac{2}{5}}$. Ans. $\frac{1}{16}\sqrt{5}$
 6. Let $\frac{3}{4}\sqrt[3]{\frac{3}{16}}$ be divided by $\frac{2}{5}\sqrt[3]{\frac{2}{5}}$. Ans. $\frac{5}{16}\sqrt[3]{30}$.
 7. Let $\frac{4}{5}\sqrt{a}$, or $\frac{4}{5}a^{\frac{1}{2}}$, be divided by $\frac{2}{3}a^{\frac{1}{3}}$. Ans. $\frac{6}{5}a^{\frac{1}{6}}$.
 8. Let $a^{\frac{4}{3}}$ be divided by $a^{\frac{2}{3}}$.
 9. To divide $3a^{\frac{1}{n}}$ by $4a_m$.

PROBLEM VIII

To Involve or Raise Surd Quantities to any Power.

RAISE both the rational part and the surd part. Or multiply the index of the quantity by the index of the power to which it is to be raised, and to the result annex the power of the rational parts, which will give the power required.

EXAMPLES.

1. Required to find the square of $\frac{3}{4}a^{\frac{1}{2}}$.

First, $(\frac{3}{4})^2 = \frac{3}{4} \times \frac{3}{4} = \frac{9}{16}$, and $(a^{\frac{1}{2}})^2 = a^{\frac{1}{2}} \times 2 = a^{\frac{2}{2}} = a$.

Therefore $(\frac{3}{4}a^{\frac{1}{2}})^2 = \frac{9}{16}a$, is the square required.

2. Required to find the square of $\frac{1}{2}a^{\frac{2}{3}}$.

First, $\frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$ and $(a^{\frac{2}{3}})^2 = a^{\frac{4}{3}} = a\sqrt[3]{a}$;

Therefore $(\frac{1}{2}a^{\frac{2}{3}})^2 = \frac{1}{4}a\sqrt[3]{a}$ is the square required.

3. Required to find the cube of $\frac{2}{3}\sqrt{6}$ or $\frac{2}{3} \times 6^{\frac{1}{2}}$.

First, $(\frac{2}{3})^3 = \frac{2}{3} \times \frac{2}{3} \times \frac{2}{3} = \frac{8}{27}$ and $(6^{\frac{1}{2}})^3 = 6^{\frac{3}{2}} = 6\sqrt{6}$.

Theref. $(\frac{2}{3}\sqrt{6})^3 = \frac{8}{27} \times 6\sqrt{6} = \frac{16}{9}\sqrt{6}$, the cube required.

4. Required the square of $2\sqrt[3]{2}$. Ans. $4\sqrt[3]{4}$.

5. Required the cube of $3^{\frac{1}{2}}$, or $\sqrt{3}$. Ans. $3\sqrt{3}$.

6. Required the 3d power of $\frac{1}{3}\sqrt{3}$. Ans. $\frac{1}{9}\sqrt{3}$.

7. Required to find the 4th power of $\frac{1}{2}\sqrt{2}$. Ans. $\frac{1}{4}$.

8. Required

8. Required to find the m th power of $a^{\frac{1}{n}}$.
 9. Required to find the square of $2 + \sqrt{3}$.

PROBLEM IX.

*To Evolve or Extract the Roots of Surd Quantities.**

EXTRACT both the rational part and the surd part. Or divide the index of the given quantity by the index of the root to be extracted; then to the result annex the root of the rational part, which will give the root required.

EXAMPLES

1. Required to find the square root of $16\sqrt{6}$.
 First, $\sqrt{16} = 4$, and $(6^{\frac{1}{2}})^{\frac{1}{2}} = 6^{\frac{1}{2} \div 2} = 6^{\frac{1}{4}}$;
 theref. $(16\sqrt{6})^{\frac{1}{2}} = 4 \cdot 6^{\frac{1}{4}} = 4\sqrt[4]{6}$, is the square root required.
2. Required to find the cube root of $\frac{1}{27}\sqrt{3}$.
 First, $\sqrt[3]{\frac{1}{27}} = \frac{1}{3}$, and $(\sqrt{3})^{\frac{1}{3}} = 3^{\frac{1}{2} \div 3} = 3^{\frac{1}{6}}$;
 theref. $(\frac{1}{27}\sqrt{3})^{\frac{1}{3}} = \frac{1}{3} \cdot 3^{\frac{1}{6}} = \frac{1}{3}\sqrt[6]{3}$, is the cube root required.
3. Required the square root of 6^3 . Ans. $6\sqrt{6}$.
 4. Required the cube root of $\frac{1}{8}a^3b$. Ans. $\frac{1}{2}a\sqrt[3]{6b}$.
 5. Required the fourth root of $16a^2$. Ans. $2\sqrt{a}$.
 6. Required to find the m th root of $x^{\frac{1}{n}}$.
 7. Required the square root of $a^2 - 6a\sqrt{b} + 9b$.

* The square root of a binomial or residual surd, $a + b$, or $a - b$ may be found thus: Take $\sqrt{a^2 - b^2} = c$;

$$\text{then } \sqrt{a+b} = \sqrt{\frac{a+c}{2}} + \sqrt{\frac{a-c}{2}};$$

$$\text{and } \sqrt{a-b} = \sqrt{\frac{a+c}{2}} - \sqrt{\frac{a-c}{2}}.$$

Thus the square root of $4 + 2\sqrt{3} = 1 + \sqrt{3}$;
 and the square root of $6 - 2\sqrt{5} = \sqrt{5} - 1$.

Note. For the cube root or any higher root see Hutton's Dictionary under the word *Binomial Surd*, Newton's Universal Arithmetic, and the treatises of Algebra by Mc. Laurin, Wood, and Bonnycastle.

INFINITE

INFINITE SERIES.

An Infinite Series is formed either from division, dividing by a compound divisor, or by extracting the root of a compound surd quantity; and is such as, being continued, would run on infinitely, in the manner of a continued decimal fraction.

But by obtaining a few of the first terms, the law of the progression will be manifest; so that the series may thence be continued, without actually performing the whole operation.

PROBLEM I.

To Reduce Fractional Quantities into Infinite Series by Division.

DIVIDE the numerator by the denominator, as in common division; then the operation, continued as far as may be thought necessary, will give the infinite series required.

EXAMPLES.

1. To change $\frac{2ab}{a+b}$ into an infinite series.

$$a+b) 2ab \dots \left(2b - \frac{2b^2}{a} + \frac{2b^3}{a^2} - \frac{2b^4}{a^3} + \&c. \right)$$

$$\underline{2ab + 2b^2}$$

$$-2b^2$$

$$\underline{-2b^2 - \frac{2b^3}{a}}$$

$$\underline{2b^3}$$

$$\underline{\frac{2b^3}{a}}$$

$$\underline{\frac{2b^3}{a} + \frac{2b^4}{a^2}}$$

$$\underline{2b^4}$$

$$\underline{\frac{2b^4}{a^2}}$$

$$\underline{\frac{2b^4}{a^2} - \frac{2b^5}{a^3}}$$

$$\underline{\frac{2b^5}{a^3}, \&c.}$$

2. Let

2. Let $\frac{1}{1-a}$ be changed into an infinite series.
 $1-a) 1 \dots\dots (1+a+a^2+a^3+a^4+\&c.$
 $\frac{1}{1-a}$

$$\frac{a}{a-a^2}$$

$$\frac{a^2}{a^2-a^3}$$

$$\frac{a^3}{a^3-a^4}$$

3. Expand $\frac{b}{a+c}$ into an infinite series.

$$\text{Ans. } \frac{b}{a} \times (1 - \frac{c}{a} + \frac{c^2}{a^2} - \frac{c^3}{a^3} + \&c.)$$

4. Expand $\frac{a}{a-b}$ into an infinite series.

$$\text{Ans. } 1 + \frac{b}{a} + \frac{b^2}{a^2} + \frac{b^3}{a^3} + \&c.$$

5. Expand $\frac{1-x}{1+x}$ into an infinite series.

$$\text{Ans. } 1 - 2x + 2x^2 - 2x^3 + 2x^4, \&c.$$

6. Expand $\frac{a^2}{(a+b)^2}$ into an infinite series.

$$\text{Ans. } 1 - \frac{2b}{a} + \frac{3b^2}{a^2} - \frac{4b^3}{a^3}, \&c.$$

7. Expand $\frac{1}{1+1} = \frac{1}{2}$, into an infinite series.

PROBLEM II.

To Reduce a Compound Surd into an Infinite Series.

EXTRACT the root as in common arithmetic; then the operation, continued as far as may be thought necessary, will give the series required. But this method is chiefly of use in extracting the square root, the operation being too tedious for the higher powers.

EXAMPLES.

EXAMPLES.

1. Extract the root of $a^2 - x^2$ in an infinite series

$$\begin{array}{r}
 a^2 - x^2 \left(a - \frac{x^2}{2a} - \frac{x^4}{8a^3} - \frac{x^6}{16a^5} - \frac{5x^8}{128a^7} \right) \&c. \\
 \hline
 a^2 \\
 \hline
 2a - \frac{x^2}{2a} - x^2 \\
 \qquad -x^2 + \frac{x^4}{4a^2} \\
 \hline
 2a - \frac{x^2}{a} - \frac{x^4}{8a^3} - \frac{x^4}{4a^2} \\
 \qquad - \frac{x^4}{4a^2} + \frac{x^6}{8a^4} + \frac{x^6}{64a^6} \\
 \hline
 2a - \frac{x^2}{a} - \frac{x^4}{4a^3} \&c. - \frac{x^6}{8a^4} - \frac{x^8}{64a^6} \\
 \qquad - \frac{x^8}{8a^4} + \frac{x^8}{16a^6} \&c. \\
 \hline
 - \frac{5x^8}{64a^6} \&c.
 \end{array}$$

2. Expand $\sqrt{1+1} = \sqrt{2}$, into an infinite series.

Ans. $1 + \frac{1}{2} - \frac{1}{8} + \frac{1}{16} - \frac{5}{128} \&c.$

3. Expand $\sqrt{1-1}$ into an infinite series.

Ans. $1 - \frac{1}{2} - \frac{1}{8} - \frac{1}{16} - \frac{5}{128} \&c.$

4. Expand $\sqrt{a^2+x}$ into an infinite series.

5. Expand $\sqrt{a^2-2bx-x^2}$ to an infinite series.

PROBLEM III.

To Extract any Root of a Binomial: or to Reduce a Binomial Surd into an Infinite Series.

THIS will be done by substituting the particular letters of the binomial, with their proper signs, in the following general theorem or formula, viz.

$$(P + PQ)^{\frac{m}{n}} = P^{\frac{m}{n}} + \frac{m}{n} A Q + \frac{m-n}{2n} B Q^2 + \frac{m-2n}{3n} C Q^3 + \&c.$$

and

and it give will the root required : observing that p denotes the first term, q the second term divided by the first, $\frac{m}{n}$ the index of the power or root ; and $A, B, C, D, \&c.$, denote the several foregoing terms with their proper signs.

EXAMPLES,

1. To extract the sq. root of $a^2 + b^2$, in an infinite series.

Here $p = a^2$, $q = \frac{b^2}{a^2}$ and $\frac{m}{n} = \frac{1}{2}$; therefore

$p \frac{m}{n} = (a^2)^{\frac{1}{2}} = a = A$, the 1st term of the series.

$\frac{m}{n} Aq = \frac{1}{2} \times a \times \frac{b^2}{a^2} = \frac{b^2}{2a} = B$, the 2d term.

$\frac{m-n}{2n} Bq = \frac{1-2}{4} \times \frac{b^2}{2a} \times \frac{b^2}{a^2} = -\frac{b^4}{2.4a^3} = C$, the 3d term.

$\frac{m-2n}{3n} Cq = \frac{1-4}{6} \times -\frac{b^4}{2.4a^3} \times \frac{b^2}{a^2} = \frac{3b^6}{2.4.6a^5} = D$ the 4th.

Hence $a + \frac{b^2}{2a} - \frac{b^4}{2.4a^3} + \frac{3.b^6}{2.4.6a^5} - \&c.$ or

$a + \frac{b^2}{2a} - \frac{b^4}{8a^3} + \frac{b^6}{16a^5} - \frac{5b^8}{128a^7} \&c.$ is the series required.

2. To find the value of $\frac{1}{(a-x)}$, or its equal $(a-x)^{-2}$, in an infinite series.*

* *Note.* To facilitate the application of the rule to fractional examples, it is proper to observe, that any surd may be taken from the denominator of a fraction and placed in the numerator, and vice versa, by only changing the sign of its index. Thus,

$\frac{1}{x^2} = 1 \times x^{-2}$ or only x^{-2} ; and $\frac{1}{(a+b)^2} = 1 \times (a+b)^{-2}$ or

$(a+b)^{-2}$; and $\frac{a^2}{(a+x)^2} = a^2 (a+x)^{-2}$; and $\frac{x^{\frac{1}{2}}}{x^{\frac{1}{3}}} = x^{\frac{1}{2}} \times x^{-\frac{1}{3}}$; also

$\frac{(a^2+x^2)^{\frac{1}{2}}}{(a^2-x^2)^{\frac{1}{2}}} = (a^2+x^2)^{\frac{1}{2}} \times (a^2-x^2)^{-\frac{1}{2}}$; &c.

Here

Here $p=a$, $q=\frac{-x}{a}=-a^{-1}x$, and $\frac{m}{n}=\frac{-2}{1}=-2$; theref.

$\frac{m}{n} = (a)^{-2} = \frac{1}{a^2} = A$, the 1st term of the series.

$\frac{m}{n} Aq = -2 \times \frac{1}{a^2} \times \frac{-x}{a} = \frac{2x}{a^3} = 2a^{-3}x = B$, the 2d term.

$\frac{m-n}{2n} Bq = -\frac{3}{2} \times \frac{2x}{a^3} \times \frac{-x}{a} = \frac{3x^2}{a^4} = 3a^{-4}x^2 = C$, the 3d.

$\frac{m-2n}{3n} Cq = -\frac{4}{3} \times \frac{3x^2}{a^4} \times \frac{-x}{a} = \frac{4x^3}{a^5} = 4a^{-5}x^3 = D$.

Hence $a^{-2} + 2a^{-3}x + 3a^{-4}x^2 + 4a^{-5}x^3 + \&c.$ or

$\frac{1}{a} + \frac{2x}{a^3} + \frac{3x^2}{a^4} + \frac{4x^3}{a^5} + \frac{5x^4}{a^6} \&c.$ is the series required.

3. To find the value of $\frac{a^2}{a-x}$, in an infinite series.

$$\text{Ans. } a + x + \frac{x^2}{a} + \frac{x^3}{a^2} + \frac{x^4}{a^3} \&c.$$

4. To expand $\sqrt{\frac{1}{(a^2+x^2)}}$ or $\frac{1}{(a^2+x^2)^{\frac{1}{2}}}$ in a series.

$$\text{Ans. } \frac{1}{a} - \frac{x^2}{2a^3} + \frac{3x^4}{8a^5} - \frac{5x^6}{16a^7} \&c.$$

5. To expand $\frac{a^2}{(a-b)^2}$ in an infinite series.

$$\text{Ans. } 1 + \frac{2b}{a} + \frac{3b^2}{a^2} + \frac{4b^3}{a^3} + \frac{5b^4}{a^4} \&c.$$

6. To expand $\sqrt{a^2-x^2}$ or $(a^2-x^2)^{\frac{1}{2}}$ in a series.

$$\text{Ans. } a - \frac{x^2}{2a} - \frac{x^4}{8a^3} - \frac{x^6}{16a^5} - \frac{5x^8}{128a^7} \&c.$$

7. Find the value of $\sqrt[3]{(a^3-b^3)}$ or $(a^3-b^3)^{\frac{1}{3}}$ in a series.

$$\text{Ans. } a - \frac{b^3}{3a^2} - \frac{b^6}{9a^5} - \frac{5b^9}{81a^8} \&c.$$

8. To find the value of $\sqrt[5]{(a^5+x^5)}$ or $(a^5+x^5)^{\frac{1}{5}}$ in a series.

$$\text{Ans. } a + \frac{x^5}{5a^4} - \frac{2x^{10}}{25a^9} + \frac{6x^{15}}{125a^{14}} \&c.$$

9. To find the square root of $\frac{a-b}{a+b}$ in an infinite series.

$$\text{Ans. } 1 - \frac{b}{a} + \frac{x^2}{2a^2} - \frac{x^3}{2a^3} \&c.$$

10. Find the cube root of $\frac{a^3}{a^3+b^3}$, in a series.

$$\text{Ans. } 1 - \frac{b^3}{3a^3} + \frac{2b^6}{9a^6} - \frac{14b^9}{81a^9} \&c.$$

ARITHMETICAL PROGRESSION.

ARITHMETICAL PROGRESSION is when a series of quantities have all the same common difference, or when they either increase or decrease by the same common difference. Thus, 2, 4, 6, 8, 10, 12, &c. are in arithmetical progression, having the common difference 2; and $a, a+d, a+2d, a+3d, a+4d, a+5d, \&c.$ are series in arithmetical progression, the common difference being d .

The most useful part of arithmetical proportion is contained in the following theorems:

1. When four quantities are in Arithmetical Proportion, the sum of the two extremes is equal to the sum of the two means. Thus, in the arithmetical 4, 6, 7, 9, the sum $4+9=6+7=13$: and in the arithmeticals, $a, a+d, b, b+d$, the sum $a+b+d=a+b+d$.

1. In any continued arithmetical progression, the sum of the two extremes is equal to the sum of any two terms at an equal distance from them.

Thus.

Thus, if the series be 1, 3, 5, 7, 9, 11, &c.

Then $1+11=3+9=5+7=12$.

2. The last term of any increasing arithmetical series, is equal to the first term increased by the product of the common difference multiplied by the number of terms less one; but in a decreasing series, the last term is equal to the first term lessened by the said product.

Thus, the 20th term of the series, 1, 3, 5, 7, 9, &c. is $= 1+2(20-1)=1+2\times 19=1+38=39$.

And the n th term of $a, a-d, a-2d, a-3d, a-4d, \&c.$ is $= a-(n-1)\times d=a-(n-1)d$.

3. The sum of all the terms in any series in arithmetical progression, is equal to half the sum of the two extremes multiplied by the number of terms.

Thus, the sum of 1, 3, 5, 7, 9, &c, continued to the 10th term, is $= \frac{(1+19)\times 10}{2} = \frac{20\times 10}{2} = 10\times 10=100$.

And the sum of n terms of $a, a+d, a+2d, a+3d, \text{ to } a+md,$ is $= (a+a+md) \cdot \frac{n}{2} = (a+\frac{1}{2}md) n$.

EXAMPLES FOR PRACTICE.

1. The first term of an increasing arithmetical series is 1, the common difference 2, and the number of terms 21; required the sum of the series?

First, $1+2\times 20=1+40=41$, is the last term.

Then $\frac{1+41}{2}\times 20=21\times 20=420$, the sum required.

2. The first term of a decreasing arithmetical series is 199, the common difference 3, and the number of terms 67; required the sum of the series?

First, $199-3\cdot 66=199-198=1$. is the last term.

Then $\frac{199+1}{2}\times 67=100\times 67=6700$, the sum required.

3. To find the sum of 100 terms of the natural numbers 1, 2, 3, 4, 5, 6, &c. Ans.. 5050.

4. Required

4. * Required the sum of 99 terms of the odd numbers 1, 3, 5, 7, 9, &c. Ans. 9811.

5. The first term of a decreasing arithmetical series is 10, the common difference $\frac{1}{3}$, and the number of terms 21; required the sum of the series? Ans. 140.

6. One hundred stones being placed on the ground, in a straight line, at the distance of 2 yards from each other: how far will a person travel, who shall bring them one by one to a basket, which is placed 2 yards from the first stone?

Ans. 11 miles and 840 yards.

APPLICATION OF ARITHMETICAL PROGRESSION TO MILITARY AFFAIRS.

QUESTION I.

A TRIANGULAR Battalion,† consisting of thirty ranks, in which the first rank is formed of one man only, the second

* The sum of any number (n) of terms of the arithmetical series of odd number 1, 3, 5, 7, 9, &c, is equal to the square (n^2) of that number. That is, If 1, 3, 5, 7, 9, &c. be the numbers, then will

1², 2², 3², 4², 5², &c. be the sums of 1, 2, 3, &c, terms.

Thus, 0 + 1 = 1 or 1², the sum of 1 term,

1 + 3 = 4 or 2², the sum of 2 terms,

4 + 5 = 9 or 3², the sum of 3 terms,

9 + 7 = 16 or 4², the sum of 4 terms, &c.

For, by the 3d theorem, 1 + 2 ($n-1$) = 1 + 2 $n-2$ = 2 $n-1$ is the last term, when the number of terms is n ; to this last term 2 $n-1$, add the first term 1, gives 2 n the sum of the extremes, or n half the sum of the extremes; then, by the 4th theorem, $n \times n = n^2$ is the sum of all the terms. Hence it appears in general, that half the sum of the extremes, is always the same as the number of the terms n ; and that the sum of all the terms, is the same as the square of the same number, n^2 .

See more on Arithmetical Progression in the Arithmetic, p. 111.

† By triangular battalion, is to be understood, a body of troops, ranged in the form of a triangle, in which the ranks exceed each other by an equal number of men:

second of 3, the third of 5, and so on : What is the strength of such a triangular battalion ? Answer, 900 men.

QUESTION II.

A detachment having 12 successive days to march, with orders to advance the first day only 2 leagues, the second $3\frac{1}{2}$, and so on increasing $1\frac{1}{2}$ league each day's march : What is the length of the whole march, and what is the last day's march ?

Answer the last day's march is $18\frac{1}{2}$ leagues, and 123 leagues is the length of the whole march.

QUESTION III.

A brigade of sappers,* having carried on 15 yards of sap the first night, the second only 13 yards, and so on, decreasing 2 yards every night, till at last they carried on in one night only 3 yards : What is the number of nights they were employed ; and what is the whole length of the sap ?

Answer, they were employed 7 nights, and the length of the whole sap was 63 yards.

men ; if the first rank consist of one man only, and the difference between the ranks be also 1, then its form is that of an equilateral triangle? and when the difference between the ranks is more than 1, its form may then be an insceles or scalene triangle. The practice of forming troops in this order, which is now laid aside, was formerly held in greater esteem than forming them in a solid square as admitting of a greater front, especially when the troops were to make simply a stand on all sides.

* A brigade of sappers, consists generally of 8 men divided equally into two parties. While one of these parties is advancing the sap, the other is furnishing the gabions, fascines, and other necessary implements, and when the first party is tired, the second takes its place, and so on till each man in turn has been at the head of the sap. A sap is a small ditch, between three and four feet in breadth and depth ; and is distinguished from the trench by its breadth only, the trench having between 10 and 15 feet breadth. As an encouragement to sappers, the pay for all the work carried on by the whole brigade, is given to the survivors.

QUESTION

QUESTION IV.

A number of gabions* being given to be placed in six ranks, one above the other, in such a manner as that each rank exceeding one another equally, the first may consist of 4 gabions, and the last of 9 : What is the number of gabions in the six ranks ; and what is the difference between each rank ?

Answer, the difference between the ranks will be 1, and the number of gabions in the six ranks will be 39.

QUESTION V.

Two detachments, distant from each other 57 leagues, and both designing to occupy an advantageous post equi-distant from each other's camp, set out at different times ; the first detachment increasing every day's march 1 league and a half, and the second detachment increasing each day's march 2 leagues : both the detachments arrive at the same time ; the first after 5 days' march, and the second after 4 days' march : What is the number of leagues marched by each detachment each day ?

The progression $\frac{7}{10}, 2\frac{2}{10}, 3\frac{7}{10}, 5\frac{2}{10}, 6\frac{7}{10}$, answers the conditions of the first detachment : and the progression $1\frac{5}{8}, 3\frac{5}{8}, 5\frac{5}{8}, 7\frac{5}{8}$, answers the conditions of the second detachment,

QUESTION VI.

A deserter, in his flight, travelling at the rate of 8 leagues a day ; and a detachment of dragoons being sent after him, with orders to march the first day only 2 leagues, the second 5 leagues, the third 8 leagues, and so on : What is the number of days necessary for the detachment to overtake the deserter, and what will be the number of leagues marched before he is overtaken ?

Answer, 5 days are necessary to overtake him ; and consequently 40 leagues will be the extent of the march.


* Gabions are baskets, open at both ends, made of ozier twigs, and of a cylindrical form ; those made use of at the trenches are 2 feet wide, and about 3 feet high ; which, being filled with earth, serve as a shelter from the enemy's fire ; and those made use of to construct batteries, are generally higher and broader. There is another sort of gabion, made use of to raise a low parapet ; its height is from 1 to 2 feet, and 1 foot wide at top, but somewhat less at bottom, to give room for placing the muzzle of a firelock between them ; these gabions serve instead of sand bags. A sand bag is generally made to contain about a cubical foot of earth.

QUESTION VII.


A convoy* distant 35 leagues, having orders to join its camp, and to march at the rate of 5 leagues per day : its escort departing at the same time, with orders to march the first day only half a league, and the last day $9\frac{1}{2}$ leagues ; and both the escort and convoy arriving at the same time : At what distance is the escort from the convoy at the end of each march ?

OF COMPUTING SHOT OR SHELLS IN A FINISHED PILE.

SHOT and shells are generally piled in three different forms, called triangular, square, or oblong piles, according as their base is either a triangle, a square, or a rectangle.



ABCD, fig. 1, is a triangular pile,
EFGH, fig. 2, is a square pile.



ABCDEF, fig. 3, is an oblong pile.

* By convoy is generally meant a supply of ammunition or provisions, conveyed to a town or army. The body of men that guard this supply is called escort.

A triangular

A triangular pile is formed by the continual laying of triangular horizontal courses of shot one above another, in such a manner, as that the sides of these courses, called rows, decrease by unity from the bottom row to the top row, which ends always in 1 shot.

A square pile is formed by the continual laying of square horizontal courses of shot one above another, in such a manner, as that the sides of these courses decrease by unity from the bottom to the top row, which ends also in 1 shot.

In the triangular and the square piles, the sides or faces being equilateral triangles, the shot contained in those faces form an arithmetical progression, having for first term unity, and for last term and number of terms, the shot contained in the bottom row; for the number of horizontal rows, or the number counted on one of the angles from the bottom to the top, is always equal to those counted on one side in the bottom: the sides or faces in either the triangular or square piles, are called arithmetical triangles; and the numbers contained in these, are called triangular numbers: $\triangle ABC$, fig. 1, EFG , fig. 2, are arithmetical triangles.

The oblong pile may be conceived as formed from the square pile $ABCD$: to one side or face of which, as AD , a number of arithmetical triangles equal to the face have been added: and the number of arithmetical triangles added to the square pile, by means of which the oblong pile is formed, is always one less than the shot in the top row; or, which is the same, equal to the difference between the bottom row of the greater side and that of the lesser.

QUESTION VIII.

To find the shot in the triangular pile $ABCD$, fig. 1, the bottom row AB consisting of 8 shot.

SOLUTION.

The proposed pile consisting of 8 horizontal courses, each of which forms an equilateral triangle; that is, the shot contained in these being in an arithmetical progression, of which the first and last term, as also the number of terms, are known; it follows, that the sum of these particular courses, or of the 8 progressions, will be the shot contained in the proposed pile; then

The

The shot of the first or lower } triangular course will be }	<hr style="width: 100%;"/> $8 + 1 \times 4 = 36$
the second - - - - -	<hr style="width: 100%;"/> $7 + 1 \times 3\frac{1}{2} = 28$
the third - - - - -	<hr style="width: 100%;"/> $6 + 1 \times 3 = 21$
the fourth - - - - -	<hr style="width: 100%;"/> $5 + 1 \times 2\frac{1}{2} = 15$
the fifth - - - - -	<hr style="width: 100%;"/> $4 + 1 \times 2 = 10$
the sixth - - - - -	<hr style="width: 100%;"/> $3 + 1 \times 1\frac{1}{2} = 6$
the seventh - - - - -	<hr style="width: 100%;"/> $2 + 1 \times 1 = 3$
the eighth - - - - -	<hr style="width: 100%;"/> $1 + 1 \times \frac{1}{2} = 1$
	<hr style="width: 100%;"/> Total . 120 shot in the pile proposed.

QUESTION IX.

To find the shot of the square pile EFGH, fig. 2, the bottom row EF consisting of 8 shot.

SOLUTION.

The bottom row containing 8 shot, the second only 7; that is, the rows forming the progression 8, 7, 6, 5, 4, 3, 2, 1, in which each of the terms being the square root of the shot contained in each separate square course employed in forming the square pile, it follows, that the sum of the squares of these roots will be the shot required: and the sum of the squares of 8, 7, 6, 5, 4, 3, 2, 1, being 204, expresses the shot in the proposed pile.

QUESTION X.

To find the shot of the oblong pile ABCDEF, fig. 3; in which BF=16, and BC=7.

SOLUTION.

The oblong pile proposed consisting of the square pile ABCD, whose bottom row is 7 shot; besides 9 arithmetical triangles or progressions, in which the first and last term, as also the number of terms, are known; it follows that,

if to the contents of the square pile	- 140
we add the sum of 9 times the progression	- 252

their total gives the contents required - 392 shot.

REMARK I.

The shot in the triangular and the square piles, as also the shot, in each horizontal course, may at once be ascertained

tained by the following table: the vertical column **A**, contains the shot in the bottom row, from 1 to 20 inclusive; the column **B** contains the triangular numbers, or number of each course; the column **C** contains the sum of the triangular numbers, that is, the shot contained in a triangular pile, commonly called pyramidal numbers; the column **D** contains the square of the numbers of the column **A**, that is, the shot contained in each square horizontal course; and the column **E** contains the sum of these squares or shot in a square pile.

C	B	A	D	E
Pyramidal numbers.	Triangular numbers.	Natural numbers.	Square of the natural numbers.	Sum of these square numbers.
1	1	1	1	1
4	3	2	4	5
10	6	3	9	14
20	10	4	16	30
35	15	5	25	55
56	21	6	36	91
84	28	7	49	140
120	36	8	64	204
165	45	9	81	285
220	55	10	100	385
286	66	11	121	506
364	78	12	144	650
455	91	13	169	819
560	105	14	196	1015
680	120	15	225	1240
816	136	16	256	1496
969	153	17	289	1785
1140	171	18	324	2109
1330	190	19	361	2470
1540	210	20	400	2870

Thus, the bottom row in a triangular pile, consisting of 9 shot, the contents will be 165; and when of 9 in the square pile, 285.—In the same manner, the contents either of a square or triangular pile being given, the shot in the bottom row may be easily ascertained.

The contents of any oblong pile by the preceding table may be also with little trouble ascertained, the less side not exceeding 20 shot, nor the difference between the less and the greater side 20. Thus, to find the shot in an oblong pile, the

the less side being 15, and the greater 35; we are first to find the contents of the square pile, by means of which the oblong pile may be conceived to be formed; that is, we are to find the contents of a square pile, whose bottom row is 15 shot; which being 1240, we are, secondly, to add these 1240 to the product 2400 of the triangular number 120, answering to 15, the number expressing the bottom row of the arithmetical triangle, multiplied by 20, the number of those triangles; and their sum, being 3640, expresses the number of shot in the proposed oblong pile.

REMARK II.

The following algebraical expressions, deduced from the investigations of the sums of the powers of numbers in arithmetical progression, which are seen upon many gunners' callipers*, serve to compute with ease and expedition the shot or shells in any pile.

That serving to compute any triangular pile, is represented by $\left. \begin{array}{l} n+2 \times n+1 \times n \\ 6 \end{array} \right\}$

That serving to compute any square pile, is represented by $\left. \begin{array}{l} n+1 \times 2n+1 \times n \\ 6 \end{array} \right\}$

In each of these, the letter n represents the number in the bottom row: hence, in a triangular pile, the number in the bottom row being 30; then this pile will be $\frac{30+2 \times 30+1}{6} = 4960$ shot or shells. In a square pile, the number in the bottom row being also 30; then this pile will be $\frac{30+1 \times 60+1 \times 30}{6} = 9455$ shot or shells.

That serving to compute any oblong pile, is represented by $\frac{2n+1+3m \times n+1 \times n}{6}$, in which the letter n denotes the

* Callipers are large compasses, with bowed shanks, serving to take the diameters of convex and concave bodies. The gunners' callipers consist of two thin rules or plates, which are moveable quite round a joint, by the plates folding one over the other: the length of each rule or plate is 6 inches, the breadth about 1 inch. It is usual to represent, on the plates, a variety of scales, tables, proportions, &c. such as are esteemed useful to be known by persons employed about artillery; but, except the measuring of the caliber of shot and cannon, and the measuring of salient and re-entering angles, none of the articles, with which the callipers are usually filled, are essential to that instrument

number of courses, and the letter m the number of shot, less one, in the top row: hence, in an oblong pile the number of courses being 30, and the top row 31; this pile will be $60+1+90 \times 30+1 \times \frac{30}{6} = 23405$ shot or shells.

PROPORTION AND GEOMETRICAL PROGRESSION.

PROPORTION contemplates the relation of quantities considered as to what part or what multiple one is of another, or how often one contains, or is contained in, another.—Of two quantities compared together, the first is called the Antecedent, and the second the Consequent. Their ratio is the quotient which arises from dividing the one by the other.

Four Quantities are proportional, when the two couples have equal ratios, or when the first is the same part or multiple of the second, as the third is of the fourth. Thus, 3, 6, 4, 8, and a, ar, b, br , are geometrical proportionals. For $\frac{6}{3} = \frac{8}{4} = 2$, and $\frac{ar}{a} = \frac{br}{b} = r$. And they are stated thus, 3 : 6 :: 4 : 8, &c.

Direct Proportion is when the same relation subsists between the first term and the second, as between the third and the fourth: As in the terms above. But Reciprocal, or Inverse Proportion, is when one quantity increases in the same proportion, as another diminishes: As in these, 3, 6, 8, 4; and these, a, ar, br, b .

Quantities are in geometrical progression, or continuous proportion, when every two terms have always the same ratio, or when the first has the same ratio to the second, as the second to the third, and the third to the fourth, &c. Thus, 2, 4, 8, 16, 32, 64, &c, and $a; ar, ar^2, ar^3, ar^4, ar^5, \&c.$ are series in geometrical progression.

The most useful part of Proportion and Geometrical Proportion is contained in the following theorems.

1. When

1. When four quantities are in proportion, the product of the two extremes is equal to the product of the two means. As in these, 3, 6, 4, 8, where $3 \times 8 = 6 \times 4 = 24$; and in these, a, ar, b, br , where $a \times br = ar \times b = abr$.

2. When four quantities are in proportion, the product of the means divided by either of the extremes gives the other extreme. Thus if $3 : 6 :: 4 : 8$, then $\frac{6 \times 4}{3} = 8$, and $\frac{6 \times 4}{8} = 3$; also if $a : ar :: b : br$, then $\frac{abr}{a} = br$, or $\frac{abr}{br} = a$. And this is the foundation of the Rule of Three.

3. If any continued geometrical progression, the product of the two extremes, and that of any other two terms, equally distant from them, are equal to each other, or equal to the square of the middle term when there is an odd number of them. So in the series, 1, 2, 4, 8, 16, 32, 64, &c. it is $1 \times 64 = 2 \times 32 = 4 \times 16 = 8 \times 8 = 64$.

4. In any continued geometrical series, the last term is equal to the first multiplied by such a power of the ratio as is denoted by 1 less than the number of terms. Thus, in the series, 3, 6, 12, 24, 48, 96, &c. it is $3 \times 2^5 = 96$.

5. The sum of any series in geometrical progression, is found by multiplying the last term by the ratio, and dividing the difference of this product and the first term by the difference between 1 and the ratio. Thus, the sum of 3, 6, 12, 24, 48, 96, 192, is $\frac{192 \times 2 - 3}{2 - 1} = 384 - 3 = 381$. And the sum of n terms of the series, $a, ar, ar^2, ar^3, ar^4, \&c.$ to ar^{n-1} , is $\frac{ar^{n-1} \times r - a}{r - 1} = \frac{ar^n - a}{r - 1} = \frac{r^n - 1}{r - 1} a$.

6. When four quantities, a, ar, b, br , or 2, 6, 4, 12, are proportional; then any of the following forms of those quantities are also proportional, viz.

1. Directly, $a : ar :: b : br$; or $2 : 6 :: 4 : 12$.
2. Inversely, $ar : a :: br : b$; or $6 : 2 :: 12 : 4$.
3. Alternately, $a : b :: ar : br$; or $2 : 4 :: 6 : 12$.
4. Compoundedly, $a : a + ar :: b : b + br$; or $2 : 8 :: 4 : 16$.
5. Dividedly, $a : ar - a :: b : br - b$, or $2 : 4 :: 4 : 8$.
6. Mixed,

6. Mixed, $ar+a : ar-a :: br+b : br-b$; or $3 : 4 :: 16 : 5$.
 7. Multiplication, $ac : arc :: bc : brc$; or $2.3 : 6.3 :: 4 : 12$.
 8. Division, $\frac{a}{c} : \frac{ar}{c} :: b : br$; or $1 : 3 :: 4 : 12$.
 9. The numbers a, b, c, d , are in harmonical proportion, when $a : d :: a \propto b : c \propto d$.

EXAMPLES.

1. Given the first term of a geometric series 1, the ratio 2, and the number of terms 12; to find the sum of the series; First, $1 \times 2^{11} = 1 \times 2048$, is the last term.

Then $\frac{2048 \times 2 - 1}{2 - 1} = \frac{4096 - 1}{1} = 4095$, the sum required.

2. Given the first term of a geometrical series $\frac{1}{3}$, the ratio $\frac{1}{2}$, and the number of terms 8; to find the sum of the series? First, $\frac{1}{3} \times (\frac{1}{2})^7 = \frac{1}{3} \times \frac{1}{128} = \frac{1}{384}$, is the last term.

Then, $(\frac{1}{3} - \frac{1}{384} \times \frac{1}{2}) \div (1 - \frac{1}{2}) = (\frac{1}{3} - \frac{1}{768}) \div \frac{1}{2} = \frac{255}{768} \times \frac{2}{1} = \frac{255}{384}$, the sum required.

3. Required the sum of 12 terms of the series, 1, 3, 9, 27, 31, &c. Ans. 265720.

4. Required the sum of 12 terms of the series 1, $\frac{1}{3}$, $\frac{1}{9}$, $\frac{1}{27}$, $\frac{1}{31}$, &c. Ans. $\frac{265720}{177147}$.

5. Required the sum of 100 terms of the series 1, 2, 4, 8, 16, 32, &c. Ans. 1267650600228229401496703205375.

See more of Proportion in the Arithmetic.

SIMPLE EQUATIONS.

An Equation is the expression of two equal quantities, with the sign of equality (=) placed between them. Thus, $10 - 4 = 6$ is an equation, denoting the equality of the quantities $10 - 4$ and 6.

Equations

Equations, are either simple or compound. A Simple Equation, is that which contains only one power of the unknown quantity, without including different powers. Thus, $x - a = b + c$, or $ax^2 = b$, is a simple equation, containing only one power of the unknown quantity x . But $x^2 - 2ax = b^2$ is a compound one.

GENERAL RULE.

Reduction of Equations, is the finding the value of the unknown quantity. And this consists in disengaging that quantity from the known ones; or in ordering the equation so, that the unknown letter or quantity may stand alone on one side of the equation, or of the mark of equality, without a co-efficient: and all the rest, or the known quantities, on the other side.—In general, the unknown quantity is disengaged from the known ones, by performing always the reverse operations. So if the known quantities are connected with it by $+$ or addition, they must be subtracted; if by minus ($-$), or subtraction, they must be added; if by multiplication, we must divide by them; if by division, we must multiply; when it is in any power, we must extract the root; and when in any radical we must raise it to the power. As in the following particular rules; which are founded on the general principle of performing equal operations on equal quantities; in which case it is evident that the results must still be equal, whether by equal additions, or subtractions, or multiplications, or divisions, or roots, or powers.

PARTICULAR RULE I.

WHEN known quantities are connected with the unknown by $+$ or $-$; transpose them to the other side of the equation, and change their signs. Which is only adding or subtracting the same quantities on both sides, in order to get all the unknown terms on one side of the equation, and all the known ones on the other side.*

Thus,

* Here it is earnestly recommended that the pupil be accustomed, at every line or step in the reduction of the equations, to name the particular operation to be performed on the equation in the line, in order to produce the next form or state of the equation, in applying each of these rules, according as the particular forms of the equation may require; applying them according to the order in which they are

Thus, if $x+5=8$; then transposing 5 gives $x=8-5=3$.
 And, if $x-3+7=9$; then transposing the 3 and 7, gives
 $x=9+3-7=5$.

Also, if $x-a+b=cd$: then by transposing a and b , it is
 $x=a-b+cd$.

In like manner, if $5x-6=4x+10$, then by transposing 6
 and $4x$, it is $5x-4x=10+6$, or $x=16$.

RULE II.

WHEN the unknown term is multiplied by any quantity ;
 divide all the terms of the equation by it.

Thus, if $ax=ab-4a$; then dividing by a , gives $x=b-4$.

And, if $3x+5=20$; then first transposing 5 gives $3x=15$;
 and then by dividing by 3, it is $x=5$.

In like manner, if $ax+3ab=4c^2$; then by dividing by a , it
 is $x+3b=\frac{4c^2}{a}$; and then transposing $3b$, gives $x=\frac{4c^2}{a}-3b$.

RULE III.

WHEN the unknown term is divided by any quantity ; we
 must then multiply all the terms of the equation by that di-
 visor ; which takes it away.

Thus, if $\frac{x}{4}=3+2$: then mult. by 4, gives $x=12+8=20$.

And, if $\frac{x}{a}=3b+2c-d$:

then by mult. a , it gives $x=3ab+2ac-ad$.

Also, if $\frac{3x}{5}-3=5+2$:

Then by transposing 3, it is $\frac{3}{5}x=10$.

And multiplying by 5, it is $3x=50$.

Lastly dividing by 3 gives $x=16\frac{2}{3}$.

are here placed; and beginning every line with the words *Then by*, as in the following specimens of Examples; which two words will always bring to his recollection, that he is to pronounce what particular operation he is to perform on the last line, in order to give the next; allotting always a single line for each operation, and ranging the equations neatly just under each other, in the several lines, as they are successively produced.

RULE

RULE IV.

WHEN the unknown quantity is included in any root or surd; transpose the rest of the terms, if there be any, by Rule 1; then raise each side to such a power as is denoted by the index of the surd; viz. square each side when it is the square root; cube each side when it is the cube root; &c. which clears that radical.

Thus, if $\sqrt{x-3}=4$; then transposing 3, gives $\sqrt{x}=7$;
And squaring both sides gives $x=49$.

And, if $\sqrt{2x+10}=8$:

Then by squaring, it becomes $2x+10=64$;

And by transposing 10, it is $2x=54$;

Lastly, dividing by 2, gives $x=27$.

Also, if $\sqrt[3]{3x+4}+3=6$:

Then by transposing 3, it is $\sqrt[3]{3x+4}=3$;

And by cubing, it is $3x+4=27$;

Also, by transposing 4, it is $3x=23$;

Lastly, dividing by 3, gives $x=7\frac{2}{3}$.

RULE V.

WHEN that side of the equation which contains the unknown quantity is a complete power, or can easily be reduced to one, by rule 1, 2, or 3; then extract the root of the said power on both sides of the equation; that is, extract the square root when it is a square power, or the cube root when it is a cube, &c.

Thus, if $x^2+8x+16=36$, or $(x+4)^2=36$:

Then by extracting the roots, it is $x+4=6$;

And by transposing 4, it is $x=6-4=2$.

And if $3x^2-19=21+35$.

Then, by transposing 19, it is $3x^2=75$;

And dividing by 3, gives $x^2=25$;

And extracting the root, gives $x=5$.

Also, if $\frac{3}{4}x^2-6=21$.

Then transposing 6, gives $\frac{3}{4}x^2=30$;

And multiplying by 4, gives $3x^2=120$;

Then dividing by 3, gives $x^2=40$;

Lastly, extracting the root, gives $x=\sqrt{40}=6.324555$.

RULE VI.

WHEN there is any analogy or proportion, it is to be changed into an equation, by multiplying the two extreme terms together, and the two means together, and making the one product equal to the other.

Thus, if $2x : 9 :: 3 : 5$.

Then, mult. the extremes and means, gives $10x=27$;

And dividing by 10, gives $x=2\frac{7}{10}$.

And if $\frac{3}{4}x : a :: 5b : 2c$.

Then mult. extremes and means gives $\frac{3}{2}cx=5ab$;

And multiplying by 2, gives $3cx=10ab$;

Lastly, dividing by $3c$, gives $x = \frac{10ab}{3c}$.

Also, if $10-x : \frac{2}{3}x :: 3 : 1$.

Then mult. extremes and means, gives $10-x=2x$;

And transposing x , gives $10=3x$;

Lastly, dividing by 3, gives $3\frac{1}{3}=x$.

RULE VII.

WHEN the same quantity is found on both sides of an equation, with the same sign, either plus or minus, it may be left out of both : and when every term in an equation is either multiplied or divided by the same quantity it may be struck out of them all.

Thus, if $3x+2a=2a+b$:

Then by taking away $2a$, it is $3x=b$.

And dividing by 3, it is $x=\frac{1}{3}b$.

Also if there be $4ax+6ab=7ac$.

Then striking out or dividing by a , gives $4x+6b=7c$.

Then, by transposing $6b$, it becomes $4x=7c-6b$;

And then dividing by 4 gives $x=\frac{7}{4}c-\frac{3}{2}b$.

Again, if $\frac{2}{3}x-\frac{7}{3}=\frac{10}{3}-\frac{7}{3}$.

Then, taking away the $\frac{7}{3}$, it becomes $\frac{2}{3}x=\frac{10}{3}$;

And taking away the 3's, it is $2x=10$;

Lastly, dividing by 2 gives $x=5$.

MISCELLANEOUS EXAMPLES.

1. Given $7x-18=4x+6$; to find the value of x .

First, transposing 18 and $5x$ gives $3x=24$;

Then dividing by 3, gives $x=8$.

2. Given

2. Given $20 - 4x - 12 = 92 - 10x$; to find x .
 First transposing 20 and 12 and $10x$, gives $6x = 84$;
 Then dividing by 6, gives $x = 14$.
3. Let $4ax - 5b = 3dx + 2c$ be given; to find x .
 First, by trans. $5b$ and $3dx$, it is $4ax - 3dx = 5b + 2c$;
 Then dividing by $4a - 3d$, gives $x = \frac{5b + 2c}{4a - 3d}$.
4. Let $5x^2 - 12x = 9x + 2x^2$ be given; to find x .
 First, by dividing by x , it is $5x - 12 = 9 + 2x$;
 Then transposing 12 and $2x$, gives $3x = 21$;
 Lastly, dividing by 3, gives $x = 7$.
5. Given $9ax^3 - 15abx^2 = 6ax^3 + 12ax^2$; to find x .
 First, dividing by $3ax$, gives $3x - 5b = 2x + 4$;
 Then transposing $5b$ and $2x$, gives $x = 5b + 4$.
6. Let $\frac{x}{3} - \frac{x}{4} + \frac{x}{5} = 2$ be given, to find x .
 First, multiplying by 3, gives $x - \frac{3}{4}x + \frac{3}{5}x = 6$;
 Then multiplying by 4, gives $x + \frac{1}{5}x = 24$.
 Also multiplying by 5, gives $17x = 120$.
 Lastly, dividing by 17, gives $x = 7\frac{1}{17}$.
7. Given $\frac{x-5}{3} + \frac{x}{2} = 12 - \frac{x-10}{3}$; to find x .
 First, mult. by 3, gives $x - 5 + \frac{3}{2}x = 36 - x + 10$.
 Then transposing 5 and x , gives $2x + \frac{3}{2}x = 51$;
 And multiplying by 2, gives $7x = 102$.
 Lastly, dividing by 7, gives $x = 14\frac{2}{7}$.
8. Let $\sqrt{\frac{3x}{4}} + 7 = 10$, be given; to find x .
 First, transposing 7, gives $\sqrt{\frac{3}{4}x} = 3$;
 Then squaring the equation, gives $\frac{3}{4}x = 9$;
 Then dividing by 3, gives $\frac{1}{4}x = 3$;
 Lastly, multiplying by 4, gives $x = 12$.
9. Let $2x + 2\sqrt{a^2 + x^2} = \frac{5a^2}{\sqrt{a^2 + x^2}}$, be given; to find x .
 First, mult. by $\sqrt{a^2 + x^2}$, gives $2x\sqrt{a^2 + x^2} + 2a^2 + 2x^2 = 5a^2$
 Then trans. $2a^2$ and $2x^2$, gives $2x\sqrt{a^2 + x^2} = 3a^2 - 2x^2$;

Then

Then by squaring, it is $4x^2 \times \overline{a^2 + x^2} = \overline{3a^2 - 2x^2}^2$;
 That is, $4a^2x^2 + 4x^4 = 9a^4 - 12a^2x^2 + 4x^4$;
 By taking $4x^4$ from both sides, it is $4a^2x^2 = 9a^4 - 12a^2x^2$;
 Then transposing $12a^2x^2$, gives $16a^2x^2 = 9a^4$;
 Dividing by a^2 , gives $16x^2 = 9a^2$;
 And dividing by 16, gives $x^2 = \frac{9}{16}a^2$;
 Lastly extracting the root, gives $x = \frac{3}{4}a$.

EXAMPLES FOR PRACTICE.

- Given $2x - 5 + 16 = 21$; to find x . Ans. $x = 5$.
- Given $9x - 15 = x + 6$; to find x . Ans. $x = 2\frac{5}{8}$.
- Given $8 - 3x + 12 = 30 - 5x + 4$; to find x . Ans. $x = 7$.
- Given $x + \frac{1}{3}x - \frac{1}{4}x = 13$; to find x . Ans. $x = 12$.
- Given $3x + \frac{1}{2}x + 2 = 5x - 4$; to find x . Ans. $x = 4$.
- Given $4ax + \frac{1}{3}a - 2 = ax - bx$; to find x .

$$\text{Ans. } x = \frac{6-a}{9a+3b}$$

- Given $\frac{1}{3}x - \frac{1}{4}x + \frac{1}{5}x = \frac{1}{2}$; to find x . Ans. $x = \frac{30}{17}$.
- Given $\sqrt{4+x} = 4 - \sqrt{x}$; to find x . Ans. $x = 2\frac{1}{4}$.
- Given $4a+x = \frac{x^2}{4a+x}$; to deter. x . Ans. $x = -2a$.

- Given $\sqrt{4a^2+x^2} = \sqrt[4]{4b^4+x^4}$; to find x .

$$\text{Ans. } x = \sqrt{\frac{b^4-4a^4}{2a^2}}$$

- Given $\sqrt{x} + \sqrt{2a+x} = \frac{4a}{\sqrt{2a+x}}$; to find x .

$$\text{Ans. } x = \frac{2}{3}a$$

- Given $\frac{a}{1+2x} + \frac{a}{1-2x} = 2b$; to find x .

$$\text{Ans. } x = \frac{1}{2}\sqrt{\frac{b-a}{b}}$$

- Given $a+x = \sqrt{a^2+x}\sqrt{4b^2+x^2}$; to find x .

$$\text{Ans. } x = \frac{b^2}{a} - a$$

OF

OF REDUCING DOUBLE, TRIPLE, &c. EQUATIONS, CONTAINING
TWO, THREE, OR MORE UNKNOWN QUANTITIES.

PROBLEM I.

To Exterminate Two Unknown Quantities; Or, to Reduce the
Two Simple Equations containing them, to a Single one.

RULE I.

FIND the value of one of the unknown letters, in terms of the other quantities, in each of the equations, by the methods already explained. Then put those two values equal to each other for a new equation, with only one unknown quantity in it, whose value is to be found as before.

Note. It is evident that we must first begin to find the values of that letter which are easiest to be found in the two proposed equations.

EXAMPLES.

1. Given $\left\{ \begin{array}{l} 2x+3y=17 \\ 5x-2y=14 \end{array} \right\}$; to find x and y .

In the 1st equat. transp. $3y$ and div. by 2, gives $x = \frac{17-3y}{2}$;

In the 2d transp. $2y$ and div. by 5 gives $x = \frac{14+2y}{5}$;

Putting these two values equal, gives $\frac{14+2y}{5} = \frac{17-3y}{2}$;

Then mult. by 5 and 2, gives $28+4y=85-15y$;

Transposing 28 and 15y, gives $19y=57$;

And dividing by 19, gives $y=3$.

And hence $x=4$.

Or, to do the same by finding two values of y , thus:

In the 1st equat. tr. $2x$ and div. by 3, gives $y = \frac{17-2x}{3}$;

In the 2d tr. $2y$ and 14, and div. by 2, gives $y = \frac{5x-14}{2}$;

Putting these two values equal, gives $\frac{5x-14}{2} = \frac{17-2x}{3}$;

Mult. by 2 and by 3, gives $15x-42=34-4x$;

Transp.

Transp. 42 and $4x$, gives $19x=76$;

Dividing by 19 , gives $x=4$.

Hence $y=3$, as before.

2. Given $\left\{ \begin{array}{l} \frac{1}{2}x + 2y = a \\ \frac{1}{2}x - 2y = b \end{array} \right\}$; to find x and y .

Ans. $x=a+b$, and $y=\frac{1}{4}a-\frac{1}{4}b$.

3. Given $3x+y=22$, and $3y+x=18$; to find x and y .

Ans. $x=6$, and $y=4$.

4. Given $\left\{ \begin{array}{l} \frac{1}{2}x + \frac{1}{3}y = 4 \\ \frac{1}{3}x + \frac{1}{2}y = 3\frac{1}{2} \end{array} \right\}$; to find x and y .

Ans. $x=6$, and $y=3$.

5. Given $\frac{2x}{3} + \frac{3y}{5} = \frac{22}{5}$, and $\frac{3x}{5} + \frac{2y}{3} = \frac{67}{15}$; to find x and y .

Ans. $x=3$, and $y=4$.

6. Given $x+2y=s$, and $x^2-4y^2=d^2$; to find x and y .

Ans. $x = \frac{s^2+d^2}{2s}$, and $y = \frac{s^2-d^2}{4s}$.

7. Given $x-2y=d$, and $x : y :: a : b$; to find x and y .

Ans. $x = \frac{ad}{a-2b}$, and $y = \frac{bd}{a-2b}$.

RULE II.

FIND the value of one of the unknown letters, in only one of the equations, as in the former rule ; and substitute this value instead of that unknown quantity in the other equation, and there will arise a new equation, with only one unknown quantity, whose value is to be found as before.

Note. It is evident that it is best to begin first with that letter whose value is easiest found in the given equations.

EXAMPLES.

1. Given $\left\{ \begin{array}{l} 2x+3y=17 \\ 5x-2y=14 \end{array} \right\}$; to find x and y .

This will admit of four ways of solution ; thus : First,

In the 1st eq. trans. $3y$ and div. by 2 , gives $x = \frac{17-3y}{2}$;

This val. subs. for x in the 2d, gives, $\frac{85-15y}{2} - 2y = 14$;

Mult. by 2 , this becomes $85-15y-4y=28$;

Transp.

Transp. $15y$ and $4y$ and 28 , gives $57=19y$;
And dividing by 19 , gives $3=y$.

$$\text{Then } x = \frac{17-3y}{2} = 4.$$

2dly, in the 2d trans. $2y$ and div. by 5 , gives $x = \frac{14+2y}{5}$;

This subst. for x in the 1st, gives $\frac{28+4y}{5} + 3y = 17$;

Mult. by 5 , gives $28+4y+15y=85$;

Transpos. 28 , gives $19y=57$;

And dividing by 19 , gives $y=3$.

$$\text{Then } x = \frac{14+2y}{5} = 4, \text{ as before.}$$

3dly, in the 1st trans. $2x$ and div. by 3 , gives $y = \frac{17-2x}{3}$;

This subst. for y in the 2d, gives, $5x - \frac{34-4x}{3} = 14$;

Multiplying by 3 gives $15x-34+4x=42$;

Transposing 34 , gives $19x=76$;

And dividing by 19 , gives $x=4$.

$$\text{Hence } y = \frac{17-2x}{3} = 3, \text{ as before.}$$

4thly, in the 2d tr. $2y$ and 14 and div. by 2 , gives $y = \frac{5x-14}{2}$;

This substituted in the 1st, gives $2x + \frac{15x-42}{2} = 17$;

Multiplying by 2 , gives $19x-42=34$;

Transposing 42 , gives $19x=76$;

And dividing by 19 , gives $x=4$.

$$\text{Hence } y = \frac{5x-14}{2} = 3, \text{ as before.}$$

2. Given $2x+3y=29$, and $3x-2y=11$; to find x and y .

Ans. $x=7$, and $y=5$.

3. Given $\begin{cases} x+y=14 \\ x-y=2 \end{cases}$; to find x and y .

Ans. $x=8$, and $y=6$.

4. Given

4. Given $\left\{ \begin{array}{l} x : y :: 3 : 2 \\ x^2 - y^2 = 20 \end{array} \right\}$; to find x and y .

Ans. $x=6$, and $y=4$.

5. Given $\frac{x}{3} + 3y = 21$, and $\frac{y}{3} + 3x = 29$; to find x and y .

Ans. $x=9$, and $y=6$.

6. Given $10 - \frac{x}{2} = \frac{y}{3} + 4$, and $\frac{x-y}{2} + \frac{x}{4} - 2 = \frac{3y-x}{5} - 1$; to find x and y .

Ans. $x=8$, and $y=6$.

7. Given $x : y :: 4 : 3$, and $x^3 - y^3 = 37$; to find x and y .

Ans. $x=4$, and $y=3$.

RULE III.

LET the given equations be so multiplied, or divided, &c. and by such numbers or quantities, as will make the terms which contain one of the unknown quantities the same in both equations; if they are not the same when first proposed.

Then by adding or subtracting the equations, according as the signs may require, there will remain a new equation, with only one unknown quantity, as before. That is, add the two equations, when the signs are unlike, but subtract them when the signs are alike, to cancel that common term.

Note. To make two unequal terms become equal, as above, multiply each term by the co-efficient of the other.

EXAMPLES.

Given $\left\{ \begin{array}{l} 5x - 3y = 9 \\ 2x + 5y = 16 \end{array} \right\}$; to find x and y .

Here we may either make the two first terms, containing x , equal, or the two 2d terms, containing y equal. To make the two first terms equal, we must multiply the 1st equation by 2, and the 2d by 5; but to make the two 2d terms equal, we must multiply the 1st equation by 5, and the 2d by 3; as follows.

1. By

1. By making the two first terms equal:
 Mult. the 1st equ. by 2, gives $10x - 6y = 18$;
 And mult. the 2d by 5, gives $10x + 25y = 80$;
 Subtr. the upper from the under, gives $31y = 62$;
 And dividing by 31, gives $y = 2$.
 Hence, from the 1st given equ. $x = \frac{9+3y}{5} = 3$.
2. By making the two 2d terms equal:
 Mult. the 1st equat. by 5, gives $25x - 15y = 45$;
 And mult. the 2d by 3, gives $6x + 15y = 48$;
 Adding these two, gives $31x = 93$;
 And dividing by 31, gives $x = 3$;
 Hence, from the 1st equ. $y = \frac{5x-9}{3} = 2$.

MISCELLANEOUS EXAMPLES.

1. Given $\frac{x+8}{4} + 6y = 21$, and $\frac{y+6}{3} + 5x = 23$; to find x and y .
 Ans. $x=4$, and $y=3$.
2. Given $\frac{3x-y}{4} + 10 = 13$, and $\frac{3y+x}{2} + 5 = 12$; to find x and y .
 Ans. $x=5$, and $y=3$.
3. Given $\frac{3x+4y}{5} + \frac{x}{4} = 10$, and $\frac{6x-2y}{3} + \frac{y}{6} = 14$; to find x and y .
 Ans. $x=8$, and $y=4$.
4. Given $3x + 4y = 38$, and $4x - 3y = 9$; to find x and y .
 Ans. $x = 6$, and $y = 5$.

PROBLEM II.

To Exterminate Three or More Unknown Quantities; Or, to Reduce the Simple Equations, containing them, to a Single one.

RULE.

THIS may be done by any of the three methods in the last problem: viz.

1. AFTER the manner of the first rule in the last problem, find the value of one of the unknown letters in each of the given equations: next put two of these values equal to each other, and then one of these and a third value equal, and so on for all the values of it; which gives a new set of equations,

with which the same process is to be repeated, and so on till there is only one equation, to be reduced by the rules for a single equation.

2. Or, as in the 2d rule of the same problem, find the value of one of the unknown quantities in one of the equations only; then substitute this value instead of it in the other equations; which gives a new set of equations to be resolved as before, by repeating the operation.

3. Or, as in the 3d rule, reduce the equations, by multiplying or dividing them, so as to make some of the terms to agree: then, by adding or subtracting them, as the signs may require, one of the letters may be exterminated, &c. as before.

EXAMPLES.

1. Given $\begin{cases} x + y + z = 9 \\ x + 2y + 3z = 16 \\ x + 3y + 4z = 31 \end{cases}$; to find x , y , and z .

1. By the 1st method:

Transp. the terms containing y and z in each equa. gives

$$x = 9 - y - z,$$

$$x = 16 - 2y - 3z,$$

$$x = 21 - 3y - 4z;$$

Then putting the 1st and 2d values equal, and the 2d and 3d values equal, give

$$9 - y - z = 16 - 2y - 3z,$$

$$16 - 2y - 3z = 21 - 3y - 4z;$$

In the 1st trans. 9, z , and $2y$, gives $y = 7 - 2z$;

In the 2d trans. 16, $3z$ and $3y$, gives $y = 5 - z$;

Putting these two equal, gives $5 - z = 7 - 2z$;

Trans. 5 and $2z$, gives $z = 2$.

Hence $y = 5 - z = 3$, and $x = 9 - y - z = 4$.

2dly. By the 2d method:

From the 1st equa. $x = 9 - y - z$;

This value of x substit. in the 2d and 3d, gives

$$9 + y + 2z = 16,$$

$$9 + 2y + 3z = 21;$$

In the 1st trans. 9 and $2z$, gives $y = 7 - 2z$;

This substit. in the last, gives $23 - z = 21$;

Trans. z and 21, gives $2 = z$.

Hence again $y = 7 - 2z = 3$, and $x = 9 - y - z = 4$.

3dly. By

3dly. By the 3d method ; subtracting the 1st equ. from the 2d, and the 2d from the 3d, gives

$$\begin{aligned}y + 2z &= 7, \\y + z &= 5;\end{aligned}$$

Subtr. the latter from the former, gives $z=2$.

Hence $y=5-z=3$, and $x=9-y-z=4$.

2. Given $\left\{ \begin{aligned}x + y + z &= 18 \\x + 3y + 2z &= 38 \\x + \frac{1}{3}y + \frac{1}{2}z &= 10\end{aligned} \right\}$; to find x , y , and z .

Ans. $x=4$, $y=6$, $z=8$.

3. Given $\left\{ \begin{aligned}x + \frac{1}{2}y + \frac{1}{3}z &= 27 \\x + \frac{1}{3}y + \frac{1}{4}z &= 20 \\x + \frac{1}{4}y + \frac{1}{5}z &= 16\end{aligned} \right\}$; to find x , y , and z .

Ans. $x=1$, $y=12$, $z=60$.

4. Given $x-y=2$, $x-z=3$, and $y+z=9$; to find x , y , and z .

Ans. $x=7$; $y=5$; $z=4$.

5. Given $\left\{ \begin{aligned}2x + 3y + 4z &= 34 \\3x + 4y + 5z &= 46 \\4x + 6y + 8z &= 58\end{aligned} \right\}$; to find x , y , and z .

A COLLECTION OF QUESTIONS PRODUCING SIMPLE EQUATIONS.

QUEST. 1. To find two numbers, such, that their sum shall be 10, and their difference 6.

Let x denote the greater number, and y the less.*

Then, by the 1st condition $x + y = 10$,

And by the 2d - - - $x - y = 6$,

Transp. y in each, gives $x = 10 - y$,

and $x = 6 + y$;

Put these two values equal, gives $6 + y = 10 - y$;

Transpos. 6 and $-y$, gives - $2y = 4$;

Dividing by 2, gives - - $y = 2$.

And hence - - - $x = 6 + y = 8$.

* In all these solutions, as many unknown letters are always used as there are unknown numbers to be found, purposely the better to exercise the modes of reducing the equations: avoiding the short ways of notation, which though giving a shorter solution, are for that reason less useful to the pupil, as affording less exercise in practising the several rules in reducing equations.

QUEST. 2. Divide 100*l.* among A, B, C, so that A may have 20*l.* more than B, and B 10*l.* more than C.

Let $x = A$'s share, $y = B$'s, and $z = C$'s.

Then $x + y + z = 100$,

$$x = y + 20,$$

$$y = z + 10.$$

In the 1st substit. $y + 20$ for x , gives $2y + z + 20 = 100$;

In this substituting $z + 10$ for y , gives $3z + 40 = 100$;

By transposing 40, gives - - $3z = 60$;

And dividing by 3, gives - - $z = 20$.

Hence $y = z + 10 = 30$, and $x = y + 20 = 50$.

QUEST. 3. A prize of 500*l.* is to be divided between two persons, so as their shares may be in proportion as 7 to 8 ; required the share of each.

Put x and y for the two shares ; then by the question,

$7 : 8 :: x : y$, or mult. the extremes

and the means, $7y = 8x$,

$$\text{and } x + y = 500 ;$$

Transposing y , gives $x = 500 - y$;

This substituted in the 1st, gives $7y = 4000 - 8y$;

By transposing $8y$, it is $15y = 4000$;

By dividing by 15, it gives $y = 266\frac{2}{3}$;

And hence $x = 500 - y = 233\frac{1}{3}$.

QUEST. 4. What number is that whose 4th part exceeds its 5th part by 10 ?

Let x denote the number sought.

Then by the question $\frac{1}{4}x - \frac{1}{5}x = 10$;

By mult. by 4, it becomes $x - \frac{4}{5}x = 40$;

By mult. by 5, it gives $x = 200$, the number sought.

QUEST. 5. What fraction is that to the numerator of which if 1 be added, the value will be $\frac{1}{2}$; but if one be added to the denominator, its value will be $\frac{1}{3}$?

Let $\frac{x}{y}$ denote the fraction.

Then by the quest. $\frac{x+1}{y} = \frac{1}{2}$, and $\frac{x}{y+1} = \frac{1}{3}$.

The 1st mult. by 2 and y , gives $2x + 2 = y$;

The 2d mult. by 3 and $y + 1$ is $3x = y + 1$;

The upper taken from the under leaves $x - 2 = 1$;

By transpos. 2, it gives $x = 3$.

And hence $y = 2x + 2 = 8$; and the fraction is $\frac{3}{8}$.

QUEST. 6.

But $5 : 2 :: 2x^2$ (the whole number of eggs) : $\frac{4}{5}x$;
 Hence $\frac{4}{5}x =$ price of both sorts, at 5 for 2 pence ;
 Then by the question $\frac{1}{2}x + \frac{1}{3}x - \frac{4}{5}x = 3$;
 Mult. by 2, gives $x + \frac{2}{3}x - \frac{8}{5}x = 6$;
 And mult. by 3, gives $5x - \frac{8}{5}x = 18$;
 Also mult. by 5, gives $x = 90$, the number of eggs of
 each sort.

QUEST. 10. Two persons, A and B, engage at play. Before they begin, A has 80 guineas, and B has 60. After a certain number of games won and lost between them, A rises with three times as many guineas as B. Query, how many guineas did A win of B ?

Let x denote the number of guineas A won.
 Then A rises with $80+x$;
 And B rises with $60-x$;
 Theref. by the quest. $80+x = 180 - 3x$;
 Transp. 80 and $3x$, gives $4x = 100$;
 And dividing by 4, gives $x = 25$, the guineas won.

QUESTIONS FOR PRACTICE.

1. To determine two numbers such, that their difference may be 4, and the difference of their squares 64.

Ans. 6 and 10.

2. To find two numbers with these conditions, viz. that half the first with a 3d part of the second may make 9, and that a 4th part of the first with a 5th part of the second may make 5.

Ans. 8 and 15.

3. To divide the number 20 into two such parts, that a 3d of the one part added to a fifth of the other, may make 6.

Ans. 15 and 5.

4. To find three numbers such, that the sum of the 1st and 2d shall be 7, the sum of the 1st and 3d 8, and the sum of the 2d and 3d 9.

Ans. 3, 4, 5.

5. A father, dying, bequeathed his fortune, which was 2800*l.* to his son and daughter, in this manner ; that for every half crown the son might have, the daughter was to have a shilling. What then were their two shares ?

Ans. The son 2000*l.* and the daughter 800*l.*

6. Three persons, A, B, C, make a joint contribution, which in the whole amounts to 400*l.* : of which sum B contributes

tributes twice as much as *A* and 20*l.* more; and *C* as much as *A* and *B* together. What sum did each contribute?

Ans. *A* 60*l.* *B* 140*l.* and *C* 200*l.*

7. A person paid a bill of 100*l.* with half guineas and crowns, using in all 202 pieces; how many pieces were there of each sort?

Ans. 180 half guineas, and 22 crowns.

8. Says *A* to *B*, if you give me 10 guineas of your money I shall then have twice as much as you will have left: but says *B* to *A*, give me 10 of your guineas, and then I shall have 3 times as many as you. How many had each?

Ans. *A* 22, *B* 26.

9. A person goes to a tavern with a certain quantity of money in his pocket, where he spends 2 shillings; he then borrows as much money as he had left, and going to another tavern, he there spends 2 shillings also; then borrowing again as much money as was left, he went to a third tavern, where likewise he spent 2 shillings; and thus repeating the same at a fourth tavern, he then had nothing remaining. What sum had he at first?

Ans. 3*s.* 9*d.*

10. A man with his wife and child dine together at an inn. The landlord charged 1 shilling for the child; and for the woman he charged as much as for the child and $\frac{1}{4}$ as much as for the man; and for the man he charged as much as for the woman and child together. How much was that for each?

Ans. The woman 20*d.* and the man 32*d.*

11. A cask, which held 60 gallons, was filled with a mixture of brandy, wine, and cyder, in this manner, viz. the cyder was 6 gallons more than the brandy, and the wine was as much as the cyder and $\frac{1}{5}$ of the brandy. How much was there of each.

Ans. Brandy 15, cyder 21, wine 24.

12. A general, disposing his army into a square form, finds that he has 284 men more than a perfect square; but increasing the side by 1 man, he then wants 25 men to be a complete square. Then how many men had he under his command?

Ans. 24000.

13. What number is that, to which if 3, 5, and 8, be severally added, the three sums shall be in geometrical progression?

Ans. 1.

14. The stock of three traders amounted to 760*l.* the shares of the first and second exceeded that of the third by

by 240; and the sum of the 2d and 3d exceeded the first by 260. What was the share of each?

Ans. The 1st 200, the 2d 300, the 3d 260.

15. What two numbers are those, which, being in the ratio of 3 to 4, their product is equal to 12 times their sum?

Ans. 21 and 28.

16. A certain company at a tavern, when they came to settle their reckoning, found that had there been 4 more in company, they might have paid a shilling a-piece less than they did; but that if there had been 3 fewer in company, they must have paid a shilling a-piece more than they did. What then was the number of persons in company, what each paid, and what was the whole reckoning?

Ans. 24 persons, each paid 7s. and the whole reckoning 8 guineas.

17. A jocky has two horses: and also two saddles, the one valued at 18*l* the other at 3*l*. Now when he sets the better saddle on the 1st horse, and the worst on the 2d, it makes the first horse worth double the 2d: but when he places the better saddle on the 2d horse, and the worse on the first, it makes the 2d horse worth three times the 1st. What then were the values of the two horses?

Ans. The 1st 6*l*. and the 2d 9*l*.

18. What two numbers are as 2 to 3, to each of which if 6 be added, the sums will be as 4 to 5?

Ans. 6 and 9.

19. What are those two numbers, of which the greater is to the less as their sum is to 20, and as their difference is to 10?

Ans. 15 and 45.

20. What two numbers are those, whose difference, sum, and product, are to each other, as the three numbers 2, 3, 5?

Ans. 2 and 10.

21. To find three numbers in arithmetical progression, of which the first is to the third as 5 to 9, and the sum of all three is 63?

Ans. 15, 21, 27.

22. It is required to divide the number 24 into two such parts, that the quotient of the greater part divided by the less, may be to the quotient of the less part divided by the greater, as 4 to 1.

Ans. 16 and 8.

23. A gentleman being asked the age of his two sons, answered, that if to the sum of their ages 18 be added, the result will be double the age of the elder; but if 6 be taken

taken from the difference of their ages, the remainder will be equal to the age of the younger. What then were their ages ?

Ans. 30 and 12.

24. To find four numbers such, that the sum of the 1st, 2d, and 3d, shall be 13 ; the sum of the 1st, 2d, and 4th, 15 ; the sum of the 1st, 3d, and 4th, 18 ; and lastly the sum of the 2d, 3d, and 4th, 20.

Ans. 2, 4, 7, 9.

25. To divide 48 into 4 such parts, that the 1st increased by 3, the second diminished by 3, the third multiplied by 3, and the 4th divided by 3, may be all equal to each other.

Ans. 6, 12, 3, 27.

QUADRATIC EQUATIONS.

QUADRATIC Equations are either simple or compound.

A simple quadratic equation, is that which involves the square of the unknown quantity only. As $ax^2=b$. And the solution of such quadratics has been already given in simple equations.

A compound quadratic equation, is that which contains the square of the unknown quantity in one term, and the first power in another term. As $ax^2+bx=c$.

All compound quadratic equations, after being properly reduced, fall under the three following forms, to which they must always be reduced by preparing them for solution.

1. $x^2+ax=b$

2. $x^2-ax=b$

3. $x^2-ax=-b$.

The general method of solving quadratic equations, is by what is called completing the square, which is as follows :

1. REDUCE the proposed equation to a proper simple form, as usual, such as the forms above ; namely, by transposing all the terms which contain the unknown quantity to one side of the equation, and the known terms to the other ; placing the square term first, and the single power second ;

dividing

dividing the equation by the co-efficient of the square or first term, if it has one, and changing the signs of all the terms, when that term happens to be negative, as that term must always be made positive before the solution. Then the proper solution is by completing the square as follows, viz.

2. Complete the unknown side to a square, in this manner, viz. Take half the co-efficient of the second term, and square it; which square add to both sides of the equation, then that side which contains the unknown quantity will be a complete square.

3. Then extract the square root on both sides of the equation,* and the value of the unknown quantity will be determined,

* As the square root of any quantity may be either $+$ or $-$, therefore all quadratic equations admit of two solutions. Thus, the square root of $+n^2$ is either $+n$ or $-n$; for $+n \times +n$ and $-n \times -n$ are each equal to $+n^2$. But the square root of $-n^2$, or $\sqrt{-n^2}$, is imaginary or impossible, as neither $+n$ nor $-n$, when squared, gives $-n^2$.

So, in the first form, $x^2 + ax = b$, where $x + \frac{1}{2}a$ is found $= \sqrt{b + \frac{1}{4}a^2}$, the root may be either $+\sqrt{b + \frac{1}{4}a^2}$, or $-\sqrt{b + \frac{1}{4}a^2}$, since either of them being multiplied by itself produces $b + \frac{1}{4}a^2$. And this ambiguity is expressed by writing the uncertain or double sign \pm before $\sqrt{b + \frac{1}{4}a^2}$; thus $x = \pm \sqrt{b + \frac{1}{4}a^2} - \frac{1}{2}a$.

In this form, where $x = \pm \sqrt{b + \frac{1}{4}a^2} - \frac{1}{2}a$, the first value of x , viz. $x = +\sqrt{b + \frac{1}{4}a^2} - \frac{1}{2}a$, is always affirmative; for since $\frac{1}{4}a^2 + b$ is greater than $-\frac{1}{4}a^2$, the greater square must necessarily have the greater root; therefore $\sqrt{b + \frac{1}{4}a^2}$ will always be greater than $\sqrt{\frac{1}{4}a^2}$, or its equal $\frac{1}{2}a$; and consequently $+\sqrt{b + \frac{1}{4}a^2} - \frac{1}{2}a$ will always be affirmative.

The second value, viz. $x = -\sqrt{b + \frac{1}{4}a^2} - \frac{1}{2}a$ will always be negative, because it is composed of two negative terms. Therefore when $x^2 + ax = b$, we shall have $x = +\sqrt{b + \frac{1}{4}a^2} - \frac{1}{2}a$ for the affirmative value of x , and $x = +\sqrt{b + \frac{1}{4}a^2} - \frac{1}{2}a$ for the negative value of x .

In the second form, where $x = \pm \sqrt{b + \frac{1}{4}a^2} + \frac{1}{2}a$ the first value, viz. $x = +\sqrt{b + \frac{1}{4}a^2} + \frac{1}{2}a$ is always affirmative, since it is composed of two affirmative terms. But the second value, viz. $x = -\sqrt{b + \frac{1}{4}a^2} + \frac{1}{2}a$, will always be negative; for since $b + \frac{1}{4}a^2$ is greater than $\frac{1}{4}a^2$, therefore $\sqrt{b + \frac{1}{4}a^2}$ will be greater than $\sqrt{\frac{1}{4}a^2}$, or its equal $\frac{1}{2}a$; and consequently $-\sqrt{b + \frac{1}{4}a^2} + \frac{1}{2}a$ is always a negative quantity.

Therefore,

determined, making the root of the known side either $+$ or $-$, which will give two roots of the equation, or two values of the unknown quantity.

Note, 1. The root of the first side of the equation, is always equal to the root of the first term, with half the coefficient of the second term joined to it; with its sign, whether $+$ or $-$.

2. All equations, in which there are two terms including the unknown quantity, and which have the index of the one just double that of the other, are resolved like quadratics, by completing the square, as above.

Thus, $x^4 + ax^2 = b$, or $x^{2n} + ax^n = b$, or $x + ax^{\frac{1}{2}} = b$, are the same as quadratics, and the value of the unknown quantity may be determined accordingly.

Therefore, when $x^2 - ax = b$, we shall have $x = +\sqrt{b + \frac{1}{4}a^2} + \frac{1}{2}a$ for the affirmative value of x ; and $x = -\sqrt{b + \frac{1}{4}a^2} + \frac{1}{2}a$ for the negative value of x ; so that in both the first and second forms, the unknown quantity has always two values, one of which is positive, and the other negative.

But in the third form, where $x = \pm\sqrt{\frac{1}{4}a^2 - b} + \frac{1}{2}a$, both the values of x will be positive when $\frac{1}{4}a^2$ is greater than b . For the first value, viz. $x = +\sqrt{\frac{1}{4}a^2 - b} + \frac{1}{2}a$ will then be affirmative, being composed of two affirmative terms.

The second value, viz. $x = -\sqrt{\frac{1}{4}a^2 - b} + \frac{1}{2}a$ is affirmative also; for since $\frac{1}{4}a^2$ is greater than $\frac{1}{4}a^2 - b$, therefore $\sqrt{\frac{1}{4}a^2}$ or $\frac{1}{2}a$ is greater than $\sqrt{\frac{1}{4}a^2 - b}$; and consequently $-\sqrt{\frac{1}{4}a^2 - b} + \frac{1}{2}a$ will always be an affirmative quantity. So that, when $x^2 - ax = b$, we shall have $x = +\sqrt{\frac{1}{4}a^2 - b} + \frac{1}{2}a$, and also $x = -\sqrt{\frac{1}{4}a^2 - b} + \frac{1}{2}a$, for the values of x , both positive.

But in this third form, if b be greater than $\frac{1}{4}a^2$, the solution of the proposed question will be impossible. For since the square of any quantity (whether that quantity be affirmative or negative) is always affirmative, the square root of a negative quantity is impossible, and cannot be assigned. But when b is greater than $\frac{1}{4}a^2$, then $\frac{1}{4}a^2 - b$ is a negative quantity; and therefore its root $\sqrt{\frac{1}{4}a^2 - b}$ is impossible, or imaginary; consequently, in that case, $x = \frac{1}{2}a \pm \sqrt{\frac{1}{4}a^2 - b}$, or the two roots or values of x , are both impossible, or imaginary quantities.

EXAMPLES.

1. Given $x^2 + 4x = 60$; to find x .

First, by completing the square $x^2 + 4x + 4 = 64$;

Then by extracting the root, $x + 2 = \pm 8$;

Then, transpos. 2, gives, $x = 6$ or -10 , the two roots.

2. Given $x^2 - 6x + 10 = 65$; to find x .

First trans. 10 gives $x^2 - 6x = 55$;

Then by complet. the sq. it is $x^2 - 6x + 9 = 64$;

And by extr. the root, gives $x - 3 = \pm 8$;

Then trans. 3, gives $x = 11$ or -5 .

3. Given $2x^2 + 8x - 30 = 60$; to find x .

First by transpos. 20, it is $2x^2 + 8x = 90$;

Then div. by 2, gives $x^2 + 4x = 45$;

And by compl. the sq. it is $x^2 + 4x + 4 = 49$;

Then extr. the root, it is $x + 2 = \pm 7$;

And transp. 2, gives $x = 5$ or -9 .

4. Given $3x^2 - 3x + 9 = 8\frac{1}{3}$; to find x .

First div. by 3, gives $x^2 - x + 3 = 2\frac{7}{9}$;

Then transpos. 3, gives $x^2 - x = -\frac{2}{9}$;

And compl. the sq. gives $x^2 - x + \frac{1}{4} = \frac{1}{36}$;

Then extr. the root gives $x - \frac{1}{2} = \pm \frac{1}{6}$;

And transp. $\frac{1}{2}$, gives $x = \frac{2}{3}$ or $\frac{1}{3}$.

5. Given $\frac{1}{2}x^2 - \frac{1}{3}x + 30\frac{1}{2} = 52\frac{2}{3}$, to find x .

First by transpos. $30\frac{1}{2}$, it is $\frac{1}{2}x^2 - \frac{1}{3}x = 22\frac{1}{3}$;

Then mult. by 2, gives $x^2 - \frac{2}{3}x = 44\frac{2}{3}$;

And by compl. the sq. it is $x^2 - \frac{2}{3}x + \frac{1}{9} = 44\frac{4}{9}$;

Then extr. the root, gives $x - \frac{1}{3} = \pm 6\frac{2}{3}$;

And transp. $\frac{1}{3}$, gives $x = 7$ or $-6\frac{1}{3}$.

6. Given $ax^2 - bx = c$; to find x .

First by div. by a , it is $x^2 - \frac{b}{a}x = \frac{c}{a}$;

Then compl. the sq. gives $x^2 - \frac{b}{a}x + \frac{b^2}{4a^2} = \frac{c}{a} + \frac{b^2}{4a^2}$;

And extrac. the root, gives $x - \frac{b}{2a} = \pm \sqrt{\frac{4ac + b^2}{4a^2}}$;

Then transp. $\frac{b}{2a}$, gives $x = \pm \sqrt{\frac{4ac + b^2}{4a^2}} + \frac{b}{2a}$.

7. Given $x^4 - 2ax^2 = b$; to find x .

First by compl. the sq. gives $x^4 - 2ax^2 + a^2 = a^2 + b$;

And

And extract. the root, gives $x^2 - a = \pm \sqrt{a^2 + b}$;

Then transpos. a , gives $x^2 = \pm \sqrt{a^2 + b + a}$;

And extract. the root, gives $x = \pm \sqrt{a \pm \sqrt{a^2 + b}}$.

And thus, by always using similar words at each line, the pupil will resolve the following examples.

EXAMPLES FOR PRACTICE.

1. Given $x^2 - 6x - 7 = 33$; to find x . Ans. $x = 10$.
2. Given $x^2 - 5x - 10 = 14$; to find x . Ans. $x = 8$.
3. Given $5x^2 + 4x - 90 = 114$; to find x . Ans. $x = 6$.
4. Given $\frac{1}{2}x^2 - \frac{1}{4}x + 2 = 9$; to find x . Ans. $x = 4$.
5. Given $3x^2 - 2x^2 = 40$; to find x . Ans. $x = 2$.
6. Given $\frac{1}{3}x - \frac{1}{2}\sqrt{x} = 1\frac{1}{2}$; to find x . Ans. $x = 9$.
7. Given $\frac{1}{2}x^2 + \frac{2}{3}x = \frac{3}{4}$; to find x . Ans. $x = .727766$.
8. Given $x^6 + 4x^3 = 12$; to find x .
Ans: $x = \sqrt[3]{2} = 1.259921$.
9. Given $x^2 + 4x = a^2 + 2$; to find x .
Ans. $x = \sqrt{a^2 + 6} - 2$.

QUESTIONS PRODUCING QUADRATIC EQUATIONS.

1. To find two numbers whose difference is 2, and product 80.

Let x and y denote the two required numbers*.

Then the first condition gives $x - y = 2$.

And the second gives $xy = 80$.

Then transp. y in the 1st gives $x = y + 2$;

This value of x substitut. in the 2d, is $y^2 + 2y = 80$

Then comp. the square gives $y^2 + 2y + 1 = 81$;

And extrac. the root gives $y + 1 = 9$;

And transpos. 1 gives $y = 8$;

And therefore $x = y + 2 = 10$.

* These questions, like those in simple equations, are also solved by using as many unknown letters, as are the numbers required, for the better exercise in reducing equations; not aiming at the shortest modes of solution, which would not afford so much useful practice.

2. To divide the number 14 into two such parts, that their product may be 48.

Let x and y denote the two numbers.

Then the 1st condition gives $x+y=14$,

And the 2d gives $xy=48$.

Then transp. y in the 1st gives $x=14-y$;

This value subst. for x in the 2d, is $14y-y^2=48$;

Changing all the signs, to make the square positive,
gives $y^2-14y=-48$;

Then compl. the square gives $y^2-14y+49=1$;

And extrac. the root gives $y-7=\pm 1$;

Then transpos. 7, gives $y=8$ or 6 , the two parts.

3. Given the sum of two numbers $= 9$, and the sum of their squares $= 45$; to find those numbers.

Let x and y denote the two numbers

Then by the 1st condition $x+y=9$.

And by the 2d $x^2+y^2=45$.

Then transpos. y in the 1st gives $x=9-y$;

This value subst. in the 2d gives $81-18y+2y^2=45$;

Then transpos. 81, gives $2y^2-18y=-36$;

And dividing by 2 gives $y^2-9y=-18$;

Then compl. the sq. gives $y^2-9y+\frac{81}{4}=\frac{9}{4}$;

And extrac. the root gives $y-\frac{9}{2}=\pm\frac{3}{2}$;

Then transpos. $\frac{9}{2}$ gives $y=6$ or 3 , the two numbers.

4. What two numbers are those, whose sum, product, and difference of their squares, are all equal to each other?

Let x and y denote the two numbers.

Then the 1st and 2d expression give $x+y=xy$.

And the 1st and 3d give $x+y=x^2-y^2$.

Then the last equa. div. by $x+y$, gives $1=x-y$;

And transpos. y , gives $y+1=x$;

This val. substit. in the 1st gives $2y+1=y^2+y$;

And transpos. $2y$, gives $1=y^2-y$;

Then complet. the sq. gives $\frac{5}{4}=y^2-y+\frac{1}{4}$;

And extracting the root gives $\frac{1}{2}\sqrt{5}=y-\frac{1}{2}$;

And transposing $\frac{1}{2}$ gives $\frac{1}{2}\sqrt{5}+\frac{1}{2}=y$;

And therefore $x=y+1=\frac{1}{2}\sqrt{5}+\frac{3}{2}$.

And if these expressions be turned into numbers, by extracting the root of 5, &c. they give $x=2.6180+$, and $y=1.6180+$.

5. There are four numbers in arithmetical progression, of which

which the product of the two extremes is 22, and that of the means 40; what are the numbers?

Let x = the less extreme,
 and y = the common difference;
 Then $x, x+y, x+2y, x+3y$, will be the four numbers.
 Hence by the 1st condition $x^2 + 3xy = 22$,
 And by the 2d $x^2 + 3xy + 2y^2 = 40$.
 Then subtracting the first from the 2d gives $2y^2 = 18$;
 And dividing by 2 gives $y^2 = 9$;
 And extracting the root gives $y = 3$.
 Then substit. 3 for y in the 1st. gives $x^2 + 9x = 22$;
 And completing the square gives $x^2 + 9x + \frac{81}{4} = \frac{169}{4}$;
 Then extracting the root gives $x + \frac{9}{2} = \frac{13}{2}$;
 And transposing $\frac{9}{2}$ gives $x = 2$ the least number.
 Hence the four numbers are 2, 5, 8, 11.

6. To find 3 numbers in geometrical progression, whose sum shall be 7, and the sum of their squares 21.

Let x, y , and z , denote the three numbers sought.
 Then by the 1st condition $xz = y^2$,
 And by the 2d $x + y + z = 7$,
 And by the 3d $x^2 + y^2 + z^2 = 21$.
 Transposing y in the 2d gives $x + z = 7 - y$;
 Sq. this equa. gives $x^2 + 2xz + z^2 = 49 - 14y + y^2$;
 Substi. $2y^2$ for $2xz$, gives $x^2 + 2y^2 + z^2 = 49 - 14y + y^2$;
 Subtr. y^2 from each side, leaves $x^2 + y^2 + z^2 = 49 - 14y$;
 Putting the two values of $x^2 + y^2 + z^2$ }
 equal to each other, gives $21 = 49 - 14y$;
 Then transposing 21 and 14y, gives $14y = 28$;
 And dividing by 14, gives $y = 2$.
 Then substit. 2 for y in the 1st equa. gives $xz = 4$.
 And in the 4th, it gives $x + z = 5$;
 Transposing z in the last, gives $x = 5 - z$;
 This substit. in the next above, gives $5z - z^2 = 4$;
 Changing all the signs, gives $z^2 - 5z = -4$;
 Then completing the square, gives $z^2 - 5z + \frac{25}{4} = \frac{9}{4}$;
 And extracting the root gives $z - \frac{5}{2} = \pm \frac{3}{2}$;
 Then transposing $\frac{5}{2}$ gives z and $x = 4$ and 1, the two other numbers;
 So that the three numbers are 1, 2, 4.

QUESTIONS FOR PRACTICE.

1. WHAT number is that which added to its square makes
 42? Ans. 6.
2. To

2. To find two numbers such, that the less may be to the greater as the greater is to 12, and that the sum of their squares may be 45. Ans. 3 and 6.

3. What two numbers are those, whose difference is 2, and the difference of their cubes 98? Ans. 3 and 5.

4. What two numbers are those whose sum is 6, and the sum of their cubes 72? Ans. 2 and 4.

5. What two numbers are those, whose product is 20, and the difference of their cubes 61; Ans. 4 and 5.

6. To divide the number 11 into two such parts, that the product of their squares may be 784. Ans. 4 and 7.

7. To divide the number 5 into two such parts, that the sum of their alternate quotients may be $4\frac{1}{4}$, that is of the two quotients of each part divided by the other. Ans. 1 and 4.

8. To divide 12 into two such parts, that their product may be equal to 8 times their difference. Ans. 4 and 8.

9. To divide the number 10 into two such parts, that the square of 4 times the less part, may be 112 more than the square of 2 times the greater. Ans. 4 and 6.

10. To find two numbers such, that the sum of their squares may be 89, and their sum multiplied by the greater may produce 104. Ans. 5 and 8.

11. What number is that, which being divided by the product of its two digits, the quotient is $5\frac{1}{3}$; but when 9 is subtracted from it, there remains a number having the same digits inverted? Ans. 32.

12. To divide 20 into three parts, such that the continual product of all three may be 270 and that the difference of first and second may be 2 less than the difference of the second and third. Ans. 5, 6, 9.

13. To find three numbers in arithmetical progression, such that the sum of their squares may be 56, and the sum arising by adding together once the first and 2 times the second and 3 times the third, may amount to 28. Ans. 2, 4, 6.

14. To divide the number 13 into three such parts, that their squares may have equal differences, and that the sum of those squares may be 75. Ans. 1, 5, 7.

15. To find three numbers having equal differences, so that their sum may be 12, and the sum of their fourth powers 962. Ans. 3, 4, 5.

16. To find three numbers having equal differences, and such that the square of the least added to the product of the two greater may make 28, but the square of the greatest added to the product of the two less may make 44. Ans. 2, 4, 6.

17. Three

17. Three merchants, A, B, C, on comparing their gains find, that among them all they have gained 144*l.*; and that B's gain added to the square root of A's made 92*l.*; but if added to the square root of C's it made 91*l.* What were their several gains? Ans. A 400, B 900, C 144.

18. To find three numbers in arithmetical progression, so that the sum of their squares shall be 93; also if the first be multiplied by 3, the second by 4, and the third by 5, the sum of the products may be 66. Ans. 2, 5, 8.

19. To find four numbers such, that the first may be to the second as the third to the fourth; and that the first may be to the fourth as 1 to 5; also the second to the third as 5 to 9; and the sum of the second and fourth may be 20. Ans. 3, 5, 9, 15.

20. To find two numbers such that their product added to their sum may make 47, and their sum taken from the sum of their squares may leave 62. Ans. 5 and 7.

RESOLUTION OF CUBIC AND HIGHER EQUATIONS.

A **CUBIC** Equation, or Equation of the 3d degree or power, is one that contains the third power of the unknown quantity. As $x^3 - ax^2 + bx = c$.

A **Biquadratic**, or Double Quadratic, is an equation that contains the 4th Power of the unknown quantity :

$$\text{As } x^4 - ax^3 + bx^2 - cx = d.$$

An Equation of the 5th Power or Degree, is one that contains the 5th power of the unknown quantity :

$$\text{As } x^5 - ax^4 + bx^3 - cx^2 + dx = e.$$

And

And so on, for all other higher powers. Where it is to be noted, however, that all the powers, or terms in the equation, are supposed to be freed from surds or fractional exponents.

There are many particular and prolix rules usually given for the solution of some of the above-mentioned powers or equations. But they may be all readily solved by the following easy rule of Double Position, sometimes called Trial-and-error.

RULE.

1. FIND, by trial, two numbers, as near the true root as you can, and substitute them separately in the given equation, instead of the unknown quantity; and find how much the terms collected together, according to their signs $+$ or $-$, differ from the absolute known term of the equation, marking whether these errors are in excess or defect.

2. Multiply the difference of the two numbers, found or taken by trial, by either of the errors, and divide the product by the difference of the errors, when they are alike, but by their sum when they are unlike. Or say, As the difference or sum of the errors, is to the difference of the two numbers, so is either error to the correction of its supposed number.

3. Add the quotient, last found, to the number belonging to that error, when its supposed number is too little, but subtract it when too great, and the result will give the true root *nearly*.

4. Take this root and the nearest of the two former, or any other that may be found nearer; and, by proceeding in like manner as above, a root will be had still nearer than before. And so on to any degree of exactness required.

Note 1. It is best to employ always two assumed numbers that shall differ from each other only by unity in the last figure on the right hand; because then the difference, or multiplier, is only 1. It is also best to use always the least error in the above operation.

Note 2. It will be convenient also to begin with a single figure at first, trying several single figures till there be found the

the two nearest the truth, the one too little, and the other too great; and in working with them, find only one more figure. Then substitute this corrected result in the equation, for the unknown letter, and if the result prove too little, substitute also the number next greater for the second supposition; but contrariwise, if the former prove too great, then take the next less number for the second supposition; and in working with the second pair of errors, continue the quotient only so far as to have the corrected number to four places of figures. Then repeat the same process again with this last corrected number, and the next greater or less, as the case may require, carrying the third corrected number to eight figures; because each new operation commonly doubles the number of true figures. And thus proceed to any extent that may be wanted.

EXAMPLES.

Ex. 1. To find the root of the cubic equation $x^3+x^2+x=100$, or the value of x in it.

Here it is soon found that x lies between 4 and 5. Assume therefore these two numbers, and the operation will be as follows:

1st Sup.		2d. Sup.
4	- x	5
16	- x^2	25
64	- x^3	125
<hr/>		
84	- sums	155
100	but should be.	100
<hr/>		
-16	- errors	+55

the sum of which is 71.
Then as $71 : 1 :: 16 : \cdot 2$.
Hence $x=4\cdot 2$ nearly.

Again, suppose $4\cdot 2$ and $4\cdot 3$; and repeat the work as follows:

1st Sup.		2d Sup.
4·2	- x	4·3
17·64	- x^2	18·49
74·088	- x^3	79·507
<hr/>		
95·928	sums	102·297
100		100
<hr/>		
-4·072	errors	+ 2·297

the sum of which is 6·369.
As $6\cdot 369 : 1 :: 2\cdot 297 : 0\cdot 036$
This taken from - 4·300

leaves x nearly $=4\cdot 264$

Again,

Again, suppose 4·264, and 4·265, and work as follows :

4·264	-	x	-	4·265
18·181696	-	x^2	-	18·190225
77·526752	-	x^3	-	77·581310
<hr/>				
99·972448	-	sums	-	100·036535
100				100
<hr/>				
-0·027552	.	errors	-	+0·036535
the sum of which is ·064087.				
Then as ·064087 : ·001 :: ·027552 : 0·0004299				
To this adding				
				<u>4·264</u>

gives x very nearly = 4·2644299

The work of the example above might have been much shortened, by the use of the Table of Powers in the Arithmetic, which would have given two or three figures by inspection. But the example has been worked out so particularly as it is, the better to show the method.

Ex. 2. To find the root of the equation $x^3 - 15x^2 + 63x = 50$, or the value of x in it.

Here it soon appears that x is very little above 1.

Suppose therefore 1·0 and 1·1 and work as follows :

1·0	-	x	-	1·1
63·0	-	$63x$	-	69·3
-15	-	$-15x^2$	-	-18·15
1	-	x^3	-	1·331
<hr/>				
49	-	sums	-	52·481
50				50
<hr/>				
-1	-	errors	-	+2·481
3·481 sum of the errors.				
As 3·481 : 1 :: 1 : ·03 correct.				
				<u>1·00</u>

Hence $x = 1·03$ nearly

Again, suppose the two numbers 1·03 and 1·02, &c. as follows :

1·03	-	x	-	1·02
64·89	-	$63x$	-	64·26
-15·9135	-	$-15x^2$	-	-15·6060
1·092727	-	x^3	-	1·061208
<hr/>				
50·069227	-	sums	-	49·715208
50				50
<hr/>				
+·069227	-	errors	-	-·284792
·284792				
As ·354019 : ·01 :: ·069227 :				
				<u>·0019555</u>
This taken from 1·03				
<hr/>				
leaves x nearly = 1·02804				

Note 3.

Note 3. Every equation has as many roots as it contains dimensions, or as there are units in the index of its highest power. That is a simple equation has only one value of the root; but a quadratic equation has two values or roots, a cubic equation has three roots, a biquadratic equation has four roots, and so on.

And when one of the roots of an equation has been found by approximation, as above, the rest may be found as follows. Take, for a dividend, the given equation, with the known term transposed, with its sign changed, to the unknown side of the equation; and, for a divisor, take x minus the root just found. Divide the said dividend by the divisor, and the quotient will be the equation depressed a degree lower than the given one.

Find a root of this new equation by approximation, as before, or otherwise, and it will be a second root of the original equation. Then, by means of this root, depress the second equation one degree lower, and from thence find a third root, and so on, till the equation be reduced to a quadratic; then the two roots of this being found, by the method of completing the square, they will make up the remainder of the roots. Thus in the foregoing equation, having found one root to be 1.02804. connect it by minus with x for a divisor, and the equation for a dividend, &c. as follows:

$$x - 1.02804) x^3 - 15x^2 + 63x - 50 (x^2 - 13.97196x + 48.63627 = 0$$

Then the two roots of this quadratic equation, or $x^2 - 13.97196x - 48.63627$, by completing the square, are 6.57653 and 7.39543, which are also the other two roots of the given cubic equation. So that all the three roots of that equation, viz. $x^3 - 15x^2 + 63x = 50$.

are 1.02804	}	and the sum of all the roots is found to be 15, being equal to the co-efficient of the 2d term of the equation, which the sum of the roots always ought to be, when they are right.
and 6.57653		
and 7.39543		
sum <u>16.00000</u>		

Note 4. It is also a particular advantage of the foregoing rule, that it is not necessary to prepare the equation, as for other rules, by reducing it to the usual final form and state of equations.

equations. Because the rule may be applied at once to an unreduced equation, though it be ever so much embarrassed by surd and compound quantities. As in the following example :

Ex. 3. Let it be required to find the root x of the equation $\sqrt{144x^2 - (x^2 + 20)^2} + \sqrt{196x^2 - (x^2 + 24)^2} = 114$, or the value of x in it.

By a few trials, it is soon found that the value of x is but little above 7. Suppose therefore first that x is = 7, and then $x=8$.

First, when $x=7$.

Second, when $x=8$.

47·906	-	$\sqrt{144x^2 - (x^2 + 20)^2}$	-	46·476
65·384	-	$\sqrt{196x^2 - (x^2 + 24)^2}$	-	69·283
113·290	-	the sums of these	-	115·759
114·000	-	the true number	-	114·000
-0·710	-	the two errors	-	+1·759
+1·759	-		-	

As 2·469 : 1 :: 0·710 : 0·2 nearly
7·0

Therefore $x=7·2$ nearly

Suppose again $x=7·2$, and then, because it turns out too great suppose x also = 7·1, &c. as follows :

Supp. $x=7·2$

Supp. $x=7·1$

47·990	-	$\sqrt{144x^2 - (x^2 + 20)^2}$	-	47·973
66·402	-	$\sqrt{196x^2 - (x^2 + 24)^2}$	-	65·904
114·392	-	the sums of these	-	113·877
114·000	-	the true number	-	114·000
+0·392	-	the two errors	-	-0·123
0·123	-		-	

As

As $\cdot 515 : \cdot 1 :: \cdot 123 : \cdot 024$ the correction,
 $7 \cdot 100$ add

Therefore $x = 7 \cdot 124$ nearly the root required.

Note 5. The same rule also among other more difficult forms of equation, succeeds very well in what are called exponential ones, or those which have an unknown quantity in the exponent of the power; as in the following example:

Ex. 4. To find the value of x in the exponential equation $x^x = 100$.

For more easily resolving such kinds of equations, it is convenient to take the logarithms of them, and then compute the terms by means of a table of logarithms. Thus, the logarithms of the two sides of the present equation are $x \times \log.$ of $x=2$ the $\log.$ of 100. Then, by a few trials, it is soon perceived that the value of x is somewhere between the two numbers 3 and 4, and indeed nearly in the middle between them, but rather nearer the latter than the former. Taking therefore first $x=3 \cdot 5$, and then $=3 \cdot 6$, and working with the logarithms, the operation will be as follows:

First supp. $x=3 \cdot 5$ Log. of $3 \cdot 5 = 0 \cdot 544068$ then $3 \cdot 5 \times \log. 3 \cdot 5 = 1 \cdot 904238$ the true number $2 \cdot 000000$ <hr style="width: 50%; margin-left: 0;"/> error, too little, $- \cdot 095762$ $\cdot 002689$		Second Supp. $x = 3 \cdot 6$ Log. of $3 \cdot 6 = 0 \cdot 556303$ then $3 \cdot 6 \times \log. 3 \cdot 6 = 2 \cdot 002689$ the true number $2 \cdot 000000$ <hr style="width: 50%; margin-left: 0;"/> error, too great $+ \cdot 002689$
--	--	---

$\cdot 098451$ sum of the errors. Then,

As $\cdot 098451 : \cdot 1 :: \cdot 002689 : 0 \cdot 00273$ the correction
 taken from $3 \cdot 60000$

leaves $- 3 \cdot 59727 = x$ nearly.*

Ex. 5. To find the value of x in the equation $x^3 + 10x^2 + 5x = 260$.
 Ans. $x = 4 \cdot 1179857$.

Ex. 6. To find the value of x in the equation $x^3 - 2x = 50$.
 Ans. $3 \cdot 8648854$.

* By repeating the operations with a larger table of logarithms, we find $x = 3 \cdot 59729502354$. Ed.

Ex. 7.

Ex. 7. To find the value of x in the equation $x^2 + 2x^2 - 23x = 70$.
 Ans. $x = 5.13457$.

Ex. 8. To find the value of x in the equation $x^3 - 17x^2 + 54x = 350$.
 Ans. $x = 14.95407$.

Ex. 9. To find the value of x in the equation $x^4 - 3x^2 - 75x = 10000$.
 Ans. $x = 10.2609$.

Ex. 10. To find the value of x in the equation $2x^4 - 16x + 40x^2 - 30x = -1$.
 Ans. $x = 1.284724$.

Ex. 11. To find the value of x in the equation $x^5 + 2x^4 + 3x^3 + 4x^2 + 5x = 54321$.
 Ans. $x = 8.414455$.

Ex. 12. To find the value of x in the equation $x^x = 123456789$.
 Ans. $x = 8.6400268$.

Ex. 13. Given $2x - 7x^3 + 11x^2 - 3x = 11$, to find x .

Ex. 14. To find the value of x in the equation
 $(3x^2 - 2\sqrt{x} + 1)^{\frac{3}{5}} - (x^2 - 4x\sqrt{x} + 3\sqrt{x})^{\frac{5}{9}} = 56$.
 Ans. $x = 18.360877$.

To resolve Cubic Equations by Carden's Rule.

THOUGH the foregoing general method, by the application of Double Position, be the readiest way, in real practice, of finding the roots in numbers of cubic equations, as well as of all the higher equations universally, we may here add the particular method commonly called Carden's Rule, for resolving cubic equations, in case any person should choose occasionally to employ that method.

The form that a cubic equation must necessarily have to be resolved by this rule, is this, viz. $z^3 + az = b$, that is, wanting the second term, or the term of the 2d power z^2 . Therefore, after any cubic equation has been reduced down to its final usual form, $x^3 + px^2 + qx = r$, freed from the co-efficient of its first term, it will then be necessary to take away the 2d term px^2 ; which is to be done in this manner: Take $\frac{1}{3}p$, or $\frac{1}{2}$ of the co-efficient of the second term, and annex it with the contrary sign to another unknown letter z , thus $z - \frac{1}{3}p$; then substitute this for x , the unknown letter in the original equation $x^3 + px^2 + qx = r$, and there will result this reduced equation $z^3 + az = b$, of the form proper for applying the following, or Carden's rule. Or take $e = \frac{1}{3}a$, and $d = \frac{1}{2}b$, by which the reduced equation takes this form $z^3 + 3cz = 2d$.

Then

Then substitute the values of c and d in this

$$\text{form } z = \sqrt[3]{d + \sqrt{(d^2 + c^3)}} + \sqrt[3]{d - \sqrt{(d^2 + c^3)}},$$

$$\text{or } z = \sqrt[3]{d + \sqrt{(d^2 + c^3)}} - \frac{\sqrt[3]{d + \sqrt{(d^2 + c^3)}}}{c},$$

and the value of the root z , of the reduced equation $z^3 + az = b$, will be obtained. Lastly, take $x = z - \frac{1}{3}p$, which will give the value of x , the required root of the original equation $x^3 + px^2 + qx = r$, first proposed.

One root of this equation being thus obtained, then depressing the original equation one degree lower, after the manner described p. 260 and 261, the other two roots of that equation will be obtained by means of the resulting quadratic equation.

Note. When the co-efficient a , or c , is negative, and c^3 is greater than d^2 , this is called the irreducible case, because then the solution cannot be generally obtained by this rule.

Ex. To find the roots of the equation $x^3 - 6x^2 + 10x = 8$.

First, to take away the 2d term, its co-efficient being -6 , its 3d part is -2 ; put therefore $x = z + 2$, then

$$\begin{array}{r} x^3 = z^3 + 6z^2 + 12z + 8 \\ -6x^2 = -6z^2 - 24z - 24 \\ +10x = +10z + 20 \end{array}$$

$$\text{theref. the sum } z^3 * -2z + 4 = 8$$

$$\text{or } z^3 * -2z = 4$$

$$\text{Here then } a = -2, b = 4, c = -\frac{2}{3}, d = 2.$$

$$\text{Theref. } \sqrt[3]{d + \sqrt{(d^2 + c^3)}} = \sqrt[3]{2 + \sqrt{(4 - \frac{8}{27})}} = \sqrt[3]{2 + \sqrt{\frac{100}{27}}} = \sqrt[3]{2 + \frac{10}{3}\sqrt{3}} = 1.57735$$

$$\text{and } \sqrt[3]{d - \sqrt{(d^2 + c^3)}} = \sqrt[3]{2 - \sqrt{(4 - \frac{8}{27})}} = \sqrt[3]{2 - \sqrt{\frac{100}{27}}} = \sqrt[3]{2 - \frac{10}{3}\sqrt{3}} = 0.42265$$

then the sum of these two is the value of $z = 2$.

Hence $x = z + 2 = 4$, one root of x in the eq. $x^3 - 6x^2 + 10x = 8$.

To find the two other roots, perform the division, &c. as in p. 261, thus:

$$\begin{array}{r} x-4) x^3 - 6x^2 + 10x - 8 \\ \underline{x^3 - 4x^2} \\ -2x^2 + 10x \\ \underline{-2x^2 + 8x} \\ 2x - 8 \\ \underline{2x - 8} \\ 0 \end{array}$$

Hence $x^2 - 2x = -2$, or $x^2 - 2x + 1 = -1$, and $x - 1 = \pm \sqrt{-1}$; $x = 1 + \sqrt{-1}$ or $= 1 - \sqrt{-1}$, the two other sought.

Ex. 2. To find the roots of $x^3 - 9x^2 + 28x = 30$.

Ans. $x = 3$, or $= 3 + \sqrt{-1}$, or $= 3 - \sqrt{-1}$.

Ex. 3. To find the roots of $x^3 - 7x^2 + 14x = 20$.

Ans. $x = 5$, or $= 1 + \sqrt{-3}$, or $= 1 - \sqrt{-3}$.

OF SIMPLE INTEREST.

As the interest of any sum, for any time, is directly proportional to the principal sum, and to the time; therefore the interest of 1 pound, for 1 year, being multiplied by any given principal sum, and by the time of its forbearance, in years and parts, will give its interest for that time. That is, if there be put

r = the rate of interest of 1 pound per annum,

p = any principal sum lent,

t = the time it is lent for, and

a = the amount or sum of principal and interest; then is $p + prt$ = the interest of the sum p , for the time t , and consequent $p + prt$ or $p \times (1 + rt) = a$, the amount for that time.

From this expression, other theorems can easily be deduced, for finding any of the quantities above mentioned; which theorems collected together, will be as below:

1st, $a = p + prt$, the amount,

2d, $p = \frac{a}{1 + rt}$, the principal,

3d, $r = \frac{a - p}{pt}$, the rate,

4th, $t = \frac{a - p}{pr}$, the time.

For Example. Let it be required to find, in what time any principal sum will double itself, at any rate of simple interest.

In this case, we must use the first theorem, $a = p + prt$, in which the amount a must be made $= 2p$, or double the principal, that is, $p + prt = 2p$, or $prt = p$, or $rt = 1$; and hence $t = \frac{1}{r}$.

Here,

Here, r being the interest of $1l.$ for 1 year, it follows, that the doubling at simple interest, is equal to the quotient of any sum divided by its interest for 1 year. So, if the rate of interest be 5 per cent. then $100 \div 5 = 20$, is the time of doubling at that rate.

Or the 4th theorem gives at once

$$t = \frac{a-p}{pr} = \frac{2p-p}{pr} = \frac{2-1}{r} = \frac{1}{r}, \text{ the same as before.}$$

COMPOUND INTEREST.

BESIDES the quantities concerned in Simple Interest namely,

p = the principal sum,

r = the rate or interest of $1l.$ for 1 year,

a = the whole amount of the principal and interest,

t = the time,

there is another quantity employed in Compound Interest, viz. the ratio of the rate of interest. which is the amount of $1l.$ for 1 time of payment. and which here let be denoted by R . viz.

$R = 1 + r$. the amount of $1l.$ for 1 time.

Then the particular amounts for the several times may be thus computed, viz. As $1l.$ is to its amount for any time, so is any proposed principal sum, to its amount for the same time; that is. as

$1l. : R :: p : pR$, the 1st year's amount,

$1l. : R :: pR : pR^2$, the 2d year's amount,

$1l. : R :: pR^2 : pR^3$, the 3d year's amount,

and so on,

Therefore, in general, $pR^t = a$ is the amount for the t year, or t time of payment. Whence the following general theorems are deduced :

1st, $a = pR^t$ the amount,

2d, $p = \frac{a}{R^t}$, the principal,

3d, $R = \sqrt[t]{\frac{a}{p}}$, the ratio,

4th, $t = \frac{\log. \text{ of } a - \log. \text{ of } p}{\log. \text{ of } R}$, the time.

From which, any one of the quantities may be found, when the rest are given.

As to the whole interest, it is found by barely subtracting the principal p from the amount a

Example. Suppose it be required to find, in how many years any principal sum will double itself, at any proposed rate of compound interest.

In this case the 4th theorem must be employed, making $a=2p$; and then it is,

$$t = \frac{\log. a - \log. p}{\log. R} = \frac{\log. 2p - \log. p}{\log. R} = \frac{\log. 2}{\log. R}$$

So, if the rate of interest be 5 per cent. per annum; then $R = 1 + .05 = 1.05$; and hence

$$t = \frac{\log. 2}{\log. 1.05} = \frac{.301030}{.021189} = 14.2067 \text{ nearly;}$$

that is, any sum doubles itself in $14\frac{1}{5}$ years nearly, at the rate of 5 per cent. per annum compound interest.

Hence, and from the like question in Simple Interest, above given, are deduced the times in which any sum doubles itself, at several rates of interest, both simple and compound; viz.

At		At Simp. Int.	At Comp. Int.
2	} per cent. per annum interest, 1 <i>l.</i> or any other sum, will double itself in the following years.	in 50	in 35.0028
2½		40	28.0701
3		33½	23.4498
3½		28½	20.1488
4		25	17.6730
4½		22½	15.7473
5		20	14.2067
6		16½	11.8957
7		14½	10.2448
8		12½	9.0065
9	11½	8.0432	
10	10	7.2725	

The following Table will very much facilitate calculations of compound interest on any sum, for any number of years, at various rates of interest.

The amounts of 1*l.* in any number of years.

Yrs.	3	3½	4	4½	5	6
1	1.0300	1.0350	1.0400	1.0450	1.0500	1.0600
2	1.0609	1.0712	1.0816	1.0920	1.1025	1.1236
3	1.0927	1.1087	1.1249	1.1412	1.1576	1.1910
4	1.1255	1.1475	1.1699	1.1925	1.2155	1.2625
5	1.1593	1.1877	1.2167	1.2462	1.2763	1.3382
6	1.1941	1.2293	1.2653	1.3023	1.3401	1.4185
7	1.2299	1.2723	1.3159	1.3609	1.4071	1.5036
8	1.2668	1.3168	1.3686	1.4221	1.4775	1.5939
9	1.3048	1.3629	1.4233	1.4861	1.5513	1.6895
10	1.3439	1.4106	1.4802	1.5530	1.6289	1.7909
11	1.3842	1.4600	1.5395	1.6229	1.7103	1.8983
12	1.4258	1.5111	1.6010	1.6939	1.7959	2.0122
13	1.4685	1.5640	1.6651	1.7722	1.8856	2.1329
14	1.5126	1.6187	1.7317	1.8519	1.9799	2.2609
15	1.5580	1.6753	1.8009	1.9353	2.0789	2.3966
16	1.6047	1.7340	1.8730	2.0224	2.1829	2.5404
17	1.6528	1.7947	1.9479	2.1134	2.2920	2.6928
18	1.7024	1.8575	2.0258	2.2085	2.4066	2.8543
19	1.7535	1.9225	2.1068	2.3079	2.5270	3.0256
20	1.8061	1.9898	2.1911	2.4117	2.6535	3.2071

The use of this Table, which contains all the powers, R^t , to the 20th power, or the amounts of 1*l.* is chiefly to calculate the interest, or the amount of any principal sum, for any time, not more than 20 years.

For example, let it be required to find, to how much 52*l.* will amount in 15 years, at the rate of 5 per cent. per annum compound interest.

In the table, on the line 15, and in the column 5 per cent.

is the amount of 1*l.* viz. - - - 2.0789

this multiplied by the principal - - - 523

gives the amount - - - 1087.2647

or - - - 1087*l.* 5*s.* ¾*d.*

and therefore the interest is - - - 564*l.* 5*s.* ¾*d.*

Note 1. When the rate of interest is to be determined to any other time than a year; as suppose to ½ a year, or ¼ a year, &c; the rules are still the same; but then *t* will express

express that time, and R must be taken the amount for that time also.

Note 2. When the compound interest, or amount, of any sum, is required for the parts of a year; it may be determined in the following manner:

1st. For any time which is some aliquot part of a year:— Find the amount of $1l.$ for 1 year, as before; then that root of it which is denoted by the aliquot part, will be the amount of $1l.$ This amount being multiplied by the principal sum, will produce the amount of the given sum as required.

2d. When the time is not an aliquot part of a year:—Reduce the time into days, and take the 365th root of the amount of $1l.$ for 1 year which will give the amount of the same for 1 day. Then raise this amount to that power whose index is equal to the number of days, and it will be the amount for that time. Which amount being multiplied by the principal sum, will produce the amount of that sum as before.—And in these calculations, the operation by logarithms will be very useful.

OF ANNUITIES.

ANNUITY is a term used for any periodical income, arising from money lent, or from houses, lands, salaries, pensions, &c. payable from time to time, but mostly by annual payments.

Annuities are divided into those that are in Possession, and those in Reversion: the former meaning such as have commenced; and the latter such as will not begin till some particular event has happened, or till after some certain time has elapsed.

When an annuity is forborn for some years, or the payments not made for that time, the annuity is said to be in Arrears.

An annuity may also be for a certain number of years; or it may be without any limit, and then it is called a Perpetuity.

The Amount of an annuity, forborn for any number of years, is the sum arising from the addition of all the annuities for that number of years, together with the interest due upon each after it becomes due.

The

The Present Worth or Value of an annuity, is the price or sum which ought to be given for it, supposing it to be bought off or paid all at once.

- Let a = the annuity, pension, or yearly rent ;
- n = the number of years forborn, or lent for ;
- R = the amount of l . for 1 year ;
- m = the amount of the annuity ;
- v = its value, or its present worth.

Now, 1 being the present value of the sum R , by proportion the present value of any other sum a , is thus found :

as $R : 1 :: a : \frac{a}{R}$ the present value of a due 1 year hence.

In like manner $\frac{a}{R^2}$ is the present value of a due 2 years,

hence ; for $R : 1 :: \frac{a}{R} : \frac{a}{R^2}$. So also $\frac{a}{R^3}, \frac{a}{R^4}, \frac{a}{R^5}$, &c. will be

the present values of a , due at the end of 3, 4, 5, &c. years respectively. Consequently the sum of all these, or

$\frac{a}{R} + \frac{a}{R^2} + \frac{a}{R^3} + \frac{a}{R^4} + \dots = (\frac{1}{R} + \frac{1}{R^2} + \frac{1}{R^3} + \frac{1}{R^4} + \dots) \times a$, conti-

nued to n terms, will be the present value of all the n years' annuities. And the value of the perpetuity, is the sum of the series to infinity.

But this series, it is evident, is a geometrical progression, *See note* having $\frac{1}{R}$ both for its first term and common ratio, and the

number of its terms n ; therefore the sum v of all the terms, or the present value of all the annual payments, will be

$$v = \frac{\frac{1}{R} - \frac{1}{R} \times \frac{1}{R^n}}{1 - \frac{1}{R}} \times a, \text{ or } = \frac{R^n - 1}{R - 1} \times \frac{a}{R^n}$$

When the annuity is a perpetuity ; n being infinite, R^n is also infinite, and therefore the quantity $\frac{1}{R^n}$ becomes $= 0$,

therefore $\frac{a}{R - 1} \times \frac{1}{R^n}$ also $= 0$; consequently the expression then

becomes

becomes barely $v = \frac{a}{R-1}$; that is, any annuity divided by the interest of 1*l.* for 1 year, gives the value of the perpetuity. So, if the rate of interest be 5 per cent.

Then $100a \div 5 = 20a$ is the value of the perpetuity at 5 per cent: Also $100a \div 4 = 25a$ is the value of the perpetuity at 4 per cent: And $100a \div 3 = 33\frac{1}{3}a$ is the value of the perpetuity at 3 per cent: and so on.

Again, because the amount of 1*l.* in n years, is R^n , its increase in that time will be $R^n - 1$; but its interest for one single year, or the annuity answering to that increase, is $R - 1$; therefore as $R - 1$ is to $R^n - 1$, so is a to m ; that is, $m = \frac{R^n - 1}{R - 1} \times a$. Hence, the several cases relating to Annuities in Arrear, will be resolved by the following equations:

$$m = \frac{R^n - 1}{R - 1} \times a = vR^n \quad \text{See note 2^d}$$

$$v = \frac{R^n - 1}{R - 1} \times \frac{a}{R^n} = \frac{m}{R^n};$$

$$a = \frac{R - 1}{R^n - 1} \times m = \frac{R - 1}{R - 1} \times vR^n;$$

$$n = \frac{\log. m - \log. v}{\log. R} = \frac{\log. \frac{mR - m + a}{a}}{\log. R}; \quad \text{See note 3^d}$$

$$\text{Log. } R = \frac{\log. m - \log. v}{n};$$

$$r = \left(\frac{1}{R^p} - \frac{1}{R^n} \right) \times \frac{a}{R - 1};$$

In this last theorem, r denotes the present value of an annuity in reversion, after p years, or not commencing till after the first p years, being found by taking the difference between the two values $\frac{R^n - 1}{R - 1} \times \frac{a}{R}$ and $\frac{R^p - 1}{R - 1} \times \frac{a}{R}$, for n years and p years.

But the amount and present value of any annuity for any number of years, up to 21, will be most readily found by the two following tables.

TABLE

ANNUITIES.

273

TABLE I.

The Amount of an Annuity of 1*l.* at Compound Interest.

Yrs.	at 3 per c.	2½ per c.	4 per c.	4½ per c.	5 per c.	6 per c.
1	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
2	2.0300	2.0350	2.0400	2.0450	2.0500	2.0600
3	3.0909	3.1062	3.1216	3.1370	3.1525	3.1836
4	4.1836	4.2149	4.2465	4.2782	4.3101	4.3746
5	5.3091	5.3625	5.4163	5.4707	5.5256	5.6371
6	6.4684	6.5502	6.6330	6.7169	6.8011	6.9753
7	7.6625	7.7794	7.8983	8.0192	8.1420	8.3938
8	8.8923	9.0517	9.2142	9.3800	9.5491	9.8975
9	10.1591	10.3685	10.5828	10.8021	11.0266	11.4913
10	11.4639	11.7314	12.0061	12.2882	12.5779	13.1808
11	12.8078	13.1420	13.4864	13.8412	14.2068	14.9716
12	14.1920	14.6020	15.0258	15.4640	15.9171	16.8699
13	15.6178	16.1130	16.6268	17.1599	17.7130	18.8821
14	17.0863	17.6770	18.2019	18.8321	19.5986	21.0151
15	18.5989	19.2957	20.8236	20.7841	21.5786	23.2760
16	20.1569	20.9710	21.8245	22.7193	23.6575	25.6725
17	21.7616	22.7050	23.6975	24.7417	25.8404	28.2129
18	23.4141	24.4997	25.6454	26.8551	28.1324	30.9057
19	25.1169	26.3572	27.6712	29.0636	30.5390	33.7600
20	26.8704	28.2797	29.7781	31.3714	33.0660	36.7856
21	28.6765	30.2695	31.9692	33.7831	35.7193	39.9927

TABLE II. The present Value of an Annuity of 1*l.*

Yrs.	at 3 per c.	3½ per c.	4 per c.	4½ per c.	5 per c.	6 per c.
1	0.9709	0.9662	0.9615	0.9569	0.9524	0.9434
2	1.9135	1.8997	1.8861	1.8727	1.8594	1.8334
3	2.8286	2.8016	2.7751	2.7490	2.7233	2.6730
4	3.7171	3.6731	3.6299	3.5875	3.5460	3.4651
5	4.5797	4.5151	4.4518	4.3900	4.3295	4.2124
6	5.4172	5.3286	5.2421	5.1579	5.0757	4.9173
7	6.2303	6.1145	6.0020	5.8927	5.7864	5.5824
8	7.0197	6.8740	6.7327	6.5959	6.4632	6.2098
9	7.7861	7.6077	7.4353	7.2688	7.1078	6.8017
10	8.5302	8.3166	8.1109	7.9127	7.7217	7.3601
11	9.2526	9.0116	8.7605	8.5289	8.3054	7.8869
12	9.9540	9.6633	9.3851	9.1186	8.8633	8.3838
13	10.6350	10.3027	9.9857	9.6829	9.3936	8.8527
14	11.2961	10.9205	10.5631	10.2228	9.8986	9.2950
15	11.9379	11.5174	11.1184	10.7396	10.3797	9.7123
16	12.5611	12.0941	11.6523	11.2340	10.8378	10.1059
17	13.1661	12.6513	12.1657	11.7072	11.2741	10.4773
18	13.7533	13.1897	12.6593	12.1600	11.6896	10.8276
19	14.3238	13.7098	13.1339	12.5933	12.0853	11.1581
20	14.8775	14.2124	13.5903	13.0079	12.4622	11.4699
21	15.4150	14.6980	14.0292	13.4047	12.8212	11.7641

To find the Amount of any annuity forborn a certain number of years.

TAKE out the amount of 1*l.* from the first table, for the proposed rate and time; then multiply it by the given annuity; and the product will be the amount, for the same number of years, and rate of interest.—And the converse to find the rate or time.

Exam. To find how much an annuity of 50*l.* will amount to in 20 years, at $3\frac{1}{2}$ per cent. compound interest.

On the line of 20 years, and in the column of $3\frac{1}{2}$ per cent. stands 28.2797, which is the amount of an annuity of 1*l.* for the 20 years. Then 28.2797×50 gives $1413.985l = 1413l. 19s. 8d.$ for the answer required.

To find the present Value of any annuity for any number of years.—Proceed here by the 2d table, in the same manner as above for the 1st table, and the present worth required will be found.

Exam. 1. To find the present value of an annuity of 50*l.* which is to continue 20 years, at $3\frac{1}{2}$ per cent.—By the table, the present value of 1*l.* for the given rate and time, is 14.2124; therefore $14.2124 \times 50 = 710.62l.$ or $710l. 12s. 4d.$ the present value required.

Exam. 2. To find the present value of an annuity of 20*l.* to commence 10 years hence, and then to continue for 11 years longer, or to terminate 21 years hence, at 4 per cent. interest.—In such cases as this, we have to find the difference between the present values of two equal annuities, for the two given times; which therefore will be done by subtracting the tabular value of the one period from that of the other, and then multiplying by the given annuity. Thus,

tabular value for 21 years 14.0292
ditto for - - 10 years 8.1109

the difference 5.9183
multiplied by 20

gives - - 118 366*l.*

or - - 118*l.* 7*s.* $3\frac{1}{2}d.$ the answer.

GEOMETRY.

DEFINITIONS.

1. **A POINT** is that which has position, but no magnitude, nor dimensions; neither length, breadth, nor thickness.

2. A **Line** is length, without breadth or thickness.

3. A **Surface** or **Superficies**, is an extension or a figure, of two dimensions, length and breadth; but without thickness.

4. A **Body** or **Solid**, is a figure of three dimensions, namely, length, breadth, and depth, or thickness.

5. **Lines** are either **Right**, or **Curved**, or **Mixed** of these two.

6. A **Right Line**, or **Straight Line**, lies all in the same direction, between its extremities; and is the shortest distance between two points.

When a line is mentioned simply, it means a **Right line**.

7. A **Curve** continually changes its direction between its extreme points.

8. **Lines** are either **Parallel**, **Oblique**, **Perpendicular**, or **Tangential**.

9. **Parallel Lines** are always at the same perpendicular distance; and they never meet though ever so far produced.


10. **Oblique lines** change their distance, and would meet, if produced on the side of the least distance.

11. One line is **Perpendicular** to another, when it inclines not more on the one side than the



the other, or when the angles on both sides of it are equal.

12. A line or circle is **Tangential**, or a **Tangent** to a circle, or other curve, when it touches it, without cutting, when both are produced.



13. An **Angle** is the inclination or opening of two lines, having different directions, and meeting in a point.




14. Angles are **Right** or **Oblique**, **Acute** or **Obtuse**.

15. A **Right Angle** is that which is made by one line perpendicular to another. Or when the angles on each side are equal to one another, they are right angles.



16. An **Oblique Angle** is that which is made by two oblique lines; and is either less or greater than a right angle.



17. An **Acute Angle** is less than a right angle.

18. An **Obtuse Angle** is greater than a right angle.



19. **Superficies** are either **Plane** or **Curved**.

20. A **Plane Superficies**, or a **Plane**, is that with which a right line may, every way coincide. Or, if the line touch the plane in two points, it will touch it in every point. But, if not, it is curved.

21. **Plane figures** are bounded either by right lines or curves.

22. **Plane figures** that are bounded by right lines have names according to the number of their sides, or of their angles; for they have as many sides as angles; the least number being three.

23. A figure of three sides and angles is called a **Triangle**. And it receives particular denominations from the relations of its sides and angles.

24. An **Equilateral Triangle** is that whose three sides are all equal.



25. An **Isosceles Triangle** is that which has two sides equal.



26. A


26. A Scalene Triangle is that whose three sides are all unequal.

27. A Right-angled Triangle is that which has one right-angle.

28. Other triangles are Oblique-angled, and are either Obtuse or Acute

29. An Obtuse-angled Triangle has one obtuse angle.

30. An Acute-angled Triangle has all its three angles acute.



31. A figure of Four sides and angles is called a Quadrangle, or a Quadrilateral.

32. A Parallelogram is a quadrilateral which has both its pairs of opposite sides parallel. And it takes the following particular names, viz. Rectangle, Square, Rhombus, Rhomboid.

33. A Rectangle is a parallelogram having a right angle.

34. A square is an equilateral rectangle ; having its length and breadth equal.


35. A Rhomboid is an oblique-angled parallelogram.

36. A Rhombus is an equilateral rhomboid ; having all its sides equal, but its angles oblique.

37. A Trapezium is a quadrilateral which hath not its opposite sides parallel.

38. A Trapezoid has only one pair of opposite sides parallel.

39. A Diagonal is a line joining any two opposite angles of a quadrilateral.



40. Plane figures that have more than four sides, are, in general, called Polygons : and they receive other particular names, according to the number of their sides or angles, Thus,

41. A Pentagon is a polygon of five sides ; a Hexagon, of six sides ; a Heptagon, seven ; an Octagon, eight ; a Nonagon, nine ; a Decagon, ten ; an Undecagon, eleven ; and a Dodecagon, twelve sides.

42. A Regular Polygon has all its sides and all its angles equal.—If they are not both equal, the polygon is Irregular.

43. An Equilateral Triangle is also a Regular Figure of three sides, and the Square is one of four, the former being also called a Trigon, and the latter a Tetragon.

44. Any figure is equilateral, when all its sides are equal; and it is equiangular when all its angles are equal. When both these are equal, it is a regular figure.

45. A Circle is a plain figure bounded by a curve line, called the Circumference, which is every where equidistant from a certain point within, called its Centre.

The circumference itself is often called a circle, and also the Periphery.

46. The Radius of a circle is a line drawn from the centre to the circumference.

47. The Diameter of a circle is a line drawn through the centre, and terminating at the circumference on both sides.

48. An Arc of a circle is any part of the circumference.

49. A Chord is a right line joining the extremities of an arc.

50. A Segment is any part of a circle bounded by an arc and its chord.

51. A semicircle is half the circle, or a segment cut off by a diameter.

The half circumference is sometimes called the Semicircle.


52. A Sector is any part of a circle which is bounded by an arc, and two radii drawn to its extremities.

53. A Quadrant, or Quarter of a circle, is a sector having a quarter of the circumference for its arc, and its two radii are perpendicular to each other. A quarter of the circumference is sometimes called a Quadrant.




54. The

54. The Height or Altitude of a figure is a perpendicular let fall from an angle, or its vertex, to the opposite side, called the base.



55. In a right-angled triangle, the side opposite the right angle is called the Hypotenuse; and the other two sides are called the Legs, and sometimes the Base and Perpendicular.

56. When an angle is denoted by three letters, of which one stands at the angular point, and the other two on the two sides, that which stands at the angular point is read in the middle. Thus the angle contained by the lines BA and AD is called the angle BAD or DAB .



57. The circumference of every circle is supposed to be divided into 360 equal parts, called Degrees: and each degree into 60 Minutes, each minutes into 60 Seconds, and so on. Hence a semicircle contains 180 degrees, and a quadrant 90 degrees.

58. The Measure of an angle, is an arc of any circle contained between the two lines which form that angle, the angular point being the centre; and it is estimated by the number of degrees contained in that arc.



59. Lines, or chords, are said to be Equidistant from the centre of a circle, when perpendiculars drawn to them from the centre are equal.



60. And the right line on which the Greater Perpendicular falls, is said to be farther from the centre.

61. An Angle in a segment is that which is contained by two lines, drawn from any point in the arc of the segment, to the two extremities of that arc.




62. An Angle On a segment, or an arc, is that which is contained by two lines, drawn from any point in the opposite or supplemental part of the circumference, to the extremities of the arc, and containing the arc between them.

63. An angle at the circumference, is that whose angular point is any where in the circumference. And an angle at the centre, is that whose angular point is at the centre.




64. A

64. A right-lined figure is **Inscribed** in a circle, or the circle **Circumscribes** it, when all the angular points of the figure are in the circumference of the circle.



65. A right-lined figure **Circumscribes** a circle, or the circle is **Inscribed** in it, when all the sides of the figure touch the circumference of the circle.



66. One right-lined figure is **Inscribed** in another, or the latter **Circumscribes** the former, when all the angular points of the former are placed in the sides of the latter.



67. A **Secant** is a line that cuts a circle, lying partly within, and partly without it.



68. Two triangles, or other right-lined figures, are said to be **mutually equilateral**, when all the sides of the one are equal to the corresponding sides of the other, each to each: and they are said to be **mutually equiangular**, when the angles of the one are respectively equal to those of the other.

69. **Identical figures**, are such as are both mutually equilateral and equiangular; or that have all the sides and all the angles of the one, respectively equal to all the sides and all the angles of the other, each to each; so that if the one figure were applied to, or laid upon the other, all the sides of the one would exactly fall upon and cover all the sides of the other; the two becoming as it were but one and the same figure.

70. **Similar figures**, are those that have all the angles of the one equal to all the angles of the other, each to each, and the sides about the equal angles proportional.

71. The **Perimeter** of a figure, is the sum of all its sides taken together.

72. A **Proposition**, is something which is either proposed to be done, or to be demonstrated, and is either a problem or a theorem.

73. A **Problem**, is something proposed to be done.

74. A **Theorem**, is something proposed to be demonstrated.

75. A **Lemma**, is something which is premised, or demonstrated, in order to render what follows more easy.

76. A **Corollary**, is a consequent truth, gained immediately from some preceding truth or demonstration.

77. A **Scholium**, is a remark or observation made upon something going before it.

AXIOMS.


A X I O M S.

1. THINGS which are equal to the same thing are equal to each other.
2. When equals are added to equals, the wholes are equal.
3. When equals are taken from equals, the remainders are equal.
4. When equals are added to unequals, the wholes are unequal.
5. When equals are taken from unequals ; the remainders are unequal.
6. Things which are double of the same thing, or equal things, are equal to each other.
7. Things which are halves of the same thing, are equal.
8. Every whole is equal to all its parts taken together.
9. Things which coincide, or fill the same space, are identical, or mutually equal in all their parts.
10. All right angles are equal to one another.
11. Angles that have equal measures, or arcs to the same radius, are equal.

THEOREM I.

IF two Triangles have Two Sides and the Included Angle in the one, equal to Two Sides and the Included Angle in the other, the Triangles will be Identical, or equal in all respects.

In the two triangles ABC , DEF , if the side AC be equal to the side DF , and the side BC equal to the side EF , and the angle C equal to the angle F ; then will the two triangles be identical, or equal in all respects.



For conceive the triangle ABC to be applied to, or placed on, the triangle DEF , in such a manner that the point c may coincide


coincide with the point F , and the side AC with the side DF , which is equal to it.

Then, since the angle F is equal to the angle c (by hyp.), the side BC will fall on the side EF . Also, because AC is equal to DF , and BC equal to EF (by hyp.), the point A will coincide with the point D , and the point B with the point E ; consequently the side AB will coincide with the side DE . Therefore the two triangles are identical, and have all their other corresponding parts equal (ax. 9), namely the side AB equal to the side DE , the angle A to the angle D , and the angle B to the angle E . Q. E. D.

THEOREM II.

WHEN Two Triangles have Two Angles and the included Side in the one, equal to Two Angles and the included Side in the other, the Triangles are Identical, or have their other sides and angle equal.

Let the two triangles ABC , DEE , have the angle A equal to the angle D , the angle B equal to the angle E , and the side AB equal to the side DE ; then these two triangles will be identical.




For, conceive the triangle ABC to be placed on the triangle DEF , in such manner that the side AB may fall exactly on the equal side DE . Then, since the angle A is equal to the angle D (by hyp.), the side AC must fall on the side DF ; and, in like manner, because the angle B is equal to the angle E , the side BC must fall on the side EF . Thus the three sides of the triangle ABC will be exactly placed on the three sides of the triangle DEF : consequently the two triangles are identical (ax. 9), having the other two sides AC , BC , equal to the two DF , EF , and the remaining angle c equal to the remaining angle F . Q. E. D.

THEOREM III.

IN an Isosceles triangle, the Angles at the Base are equal. Or, if a Triangle have Two Sides equal, their Opposite Angles will also be equal.

If the triangle ABC have the side AC equal to the side BC : then will the angle B be equal to the angle A .

For, conceive the angle c to be bisected, or divided into two equal parts by the line CD , making the angle ACD equal to the angle BCD .



Then, the two triangles ACD , BCD , have two sides and the contained angle of the one, equal to two sides and the contained angle of the other, viz. the side AC equal to BC , the angle ACD equal to BCD , and the side CD common; therefore these two triangles are identical, or equal in all respects (th. 1); and consequently the angle A equal to the angle B . Q. E. D.

Corol. 1. Hence the line which bisects the verticle angle of an isosceles triangle, bisects the base, and is also perpendicular to it.


Corol. 2. Hence too it appears, that every equilateral triangle, is also equiangular, or has all its angles equal.

THEOREM IV.

WHEN a triangle has Two of its Angles equal, the Sides Opposite to them are also equal.

If the triangle ABC , have the angle A equal to the angle B , it will also have the side AC equal to the side BC .

For, conceive the side AB to be bisected in the point D , making AD equal to DB ; and join DC , dividing the whole triangle into the two triangles ACD , BCD . Also conceive the triangle ACD to be turned over upon the triangle BCD , so that AD may fall on BD .




Then, because the line AD is equal to the line DB (by hyp.), the point A coincides with the point B , and the point D with the point D . Also, because the angle A is equal to the angle B by (hyp) the line AC will fall on the line BC , and the extremity C of the side AC will coincide with the extremity C of the side BC , because DC is common to both; consequently the side AC is equal to BC . Q. E. D.*

Corol. Hence every equiangular triangle is also equilateral.

THEOREM V.

WHEN Two Triangles have all the Three Sides in the one, equal to all the Three Sides in the other, the Triangles are Identical, or have also their Three Angles equal, each to each.

Let the two triangles ABC , AED , have their three sides respectively equal, viz. the side AB equal to AB , AC to AD , and BC to BD ; then shall the two triangles be identical, or have their angles equal, viz. those angles




* This demonstration of Theorem iv. does not appear to me to be conclusive.
EDITOR. *had the the*

that

that are opposite to the equal sides ; namely, the angle BAC to the angle BAD , the angle ABC to the angle ABD , and the angle C to the angle D .

For, conceive the two triangles to be joined together by their longest equal sides, and draw the line CD .



Then, in the triangle ACD , because the side AC is equal to AD (by hyp.), the angle ACD is equal to the angle ADC (th. 3). In like manner, in the triangle BCD , the angle BCD is equal to the angle BDC , because the side BC is equal to BD . Hence then, the angle ACD being equal to the angle ADC , and the angle BCD to the angle BDC , by equal additions the sum of the two angles ACD, BCD , is equal to the sum of the two ADC, BDC , (ax. 2), that is, the whole angle ACB equal to the whole angle ADB .

Since then, the two sides AC, CB , are equal to the two sides AD, DB , each to each, (by hyp.), and their contained angles ACB, ABD , also equal, the two triangles ABC, ABD , are identical (th. 1), and have the other angles equal, viz. the angle BAC to the angle BAD , and the angle ABC to the angle ABD . Q. E. D.

THEOREM VI.

WHEN one Line meets another, the Angles which it makes on the Same Side of the other, are together equal to Two Right Angles.

Let the line AB meet the line CD : then will the two angles ABC, ABD , taken together, be equal to two right angles.

For, first, when the two angles ABC, ABD , are equal to each other, they are both of them right angles (def. 15.)



But when the angles are unequal, suppose BE drawn perpendicular to CD . Then, since the two angles EBC, EBD , are right angles (def. 15), and the angle EBD is equal to the two angles EBA, ABD , together (ax. 3), the three angles, EBC, EBA , and ABD , are equal to two right Angles.

But the two angles EBC, EBA , are together equal to the angle ABC (ax. 8). Consequently the two angles ABC, ABD , are also equal to two right angles. Q. E. D.


Corol. 1. Hence also, conversely, if the two angles ABC, ABD , on both sides of the line AB , make up together two right angles, then CB and BD form one continued right line CD .

Corol.

Corol. 2. Hence, all the angles which can be made, at any point B, by any number of lines, on the same side of the right line CD, are, when taken all together, equal to two right angles.

Corol. 3. And, as all the angles that can be made on the other side of the line CD are also equal to two right angles; therefore all the angles that can be made quite round a point B, by any number of lines, are equal to four right angles.

Corol. 4. Hence also the whole circumference of a circle, being the sum of the measures of all the angles that can be made about the centre F (def. 57), is the measure of four right angles. Consequently, a semicircle, or 180 degrees, is the measure of two right angles; and a quadrant, or 90 degrees, the measure of one right angle.



THEOREM VII.

WHEN two Lines Intersect each other, the Opposite Angles are equal.

Let the two lines AB, CD, intersect in the point E; then will the angle AEC be equal to the angle BED, and the angle AED equal to the angle CEB.


For since the line CE meets the line AB, the two angles AEC, BEC, taken together, are equal to two right angles (th. 6).

In like manner, the line BE, meeting the line CD, makes the two angles BEC, BED, equal to two right angles.

Therefore the sum of the two angles AEC, BEC, is equal to the sum of the two BEC, BED (ax. 1).

And if the angle BEC, which is common, be taken away from both these, the remaining angle AEC will be equal to the remaining angle BED (ax. 3).

And in like manner it may be shown, that the angle AED is equal to the opposite angle BEC.




THEOREM VIII.

WHEN One Side of a triangle is produced, the Outward Angle is Greater than either of the two Inward Opposite Angles.

Let

Let ABC be a triangle, having the side AB produced to D ; then will the outward angle CBD be greater than either of the inward opposite angles A or C .

For, conceive the side BC to be bisected in the point E , and draw the line AE producing it till EF be equal to AE : and join BF .



Then, since the two triangles AEC , BEF , have the side $AE =$ the side EF , and the side $CE =$ the side BE (by suppos.) and the included or opposite angles at E also equal (th. 7), therefore those two triangles are equal in all respects (th. 1), and have the angle $c =$ the corresponding angle EBF . But the angle CBD is greater than the angle EBF ; consequently the said outward angle CBD is also greater than the angle c .

In like manner, if CB be produced to G , and AB be bisected, it may be shown that the outward angle ABG , or its equal CBD , is greater than the other angle A .

THEOREM IX.

THE Greater Side, of every Triangle, is opposite to the Greater Angle; and the Greater Angle opposite to the Greater Side.

Let ABC be a triangle, having the side AB greater than the side AC ; then will the angle ACB , opposite the greater side AB , be greater than the angle B , opposite the less side AC .



For, on the greater side AB , take the part AD equal to the less side AC , and join CD . Then, since BCD is a triangle, the outward angle ADC is greater than the inward opposite angle B (th. 8). But the angle ACD is equal to the said outward angle ADC , because AD is equal to AC (th. 3). Consequently the angle ACD also is greater than the angle B . And since the angle ACD is only a part of ACB , much more must the whole angle ACB be greater than the angle B . Q. E. D.

Again conversely, if the angle c be greater than the angle B , then will the side AB , opposite the former, be greater than the side AC , opposite the latter.

For, if AB be not greater than AC , it must be either equal to it, or less than it. But it cannot be equal, for then


then the angle c would be equal to the angle B (th. 3), which it is not, by the supposition. Neither can it be less, for then the angle c would be less than the angle B , by the former part of this; which is also contrary to the supposition. The side AB , then, being neither equal to AC , nor less than it, must necessarily be greater. Q. E. D.

THEOREM X.

THE Sum of any Two Sides of a Triangle is Greater than the Third Side.

Let ABC be a triangle; then will the sum of any two of its sides be greater than the third side, as for instances, $AC + CB$ greater than AB .

For, produce AC till CD be equal to CB , or AD equal to the sum of the two $AC + CB$; and join BD :—Then, because CD is equal to CB (by constr.), the angle D is equal to the angle CBD (th. 3). But the angle ABD is greater than the angle CBD , consequently it must also be greater than the angle D . And, since the greater side of any triangle is opposite to the greater angle (th. 9), the side AD (of the triangle ABD) is greater than the side AB . But AD is equal to AC and CD , or AC and CB , taken together (by constr.); therefore $AC + CB$ is also greater than AB . Q. E. D.




Corol. The shortest distance between two points, is a single right line drawn from the one point to the other.

THEOREM XI.

THE Difference of any Two Sides of a Triangle, is Less than the Third Side.

Let ABC be a triangle; then will the difference of any two sides, as $AB - AC$, be less than the third side BC .

For, produce the less side AC to D , till AD be equal to the greater side AB , so that CD may be the difference of the two sides $AB - AC$; and join BD . Then, because AD is equal to AB (by constr.), the opposite angles D and ABD are equal (th. 3). But the angle CBD is less than the angle ABD , and consequently also less than the equal angle D . And since the greater side of any triangle is opposite to the greater



greater angle (th. 9), the side CD (of the triangle BCD) is less than the side BC . Q. E. D.

THEOREM XII.

WHEN a Line Intersects two Parallel Lines, it makes the Alternate Angles Equal to each other.

Let the line EF cut the two parallel lines AB , CD ; then will the angle AEF be equal to the alternate angle EFD .

For if they are not equal, one of them must be greater than the other; let it be EFD for instance which is the greater if possible; and conceive the line FB to be drawn; cutting off the part of angle EFD equal to the angle AEF ; and meeting the line AB in the point B .

Then, since the outward angle AEF , of the triangle BEF , is greater than the inward opposite angle EFB (th. 8); and since these two angles also are equal (by the constr.) it follows, that these angles are both equal and unequal at the same time: which is impossible. Therefore the angle EFD is not unequal to the alternate angle AEF , that is, they are equal to each other. Q. E. D.

Corol. Right lines which are perpendicular to one, of two parallel lines, are also perpendicular to the other.

THEOREM XIII.



WHEN a line, cutting Two other Lines, makes the Alternate Angles Equal to each other, those two Lines are Parallel.

Let the line EF , cutting the two lines AB , CD , make the alternate angles AEF , DFE , equal to each other; then will AB be parallel to CD .

For if they be not parallel, let some other line, as FG , be parallel to AB . Then, because of these parallels, the angle AEF is equal to the alternate angle EFG (th. 12). But the angle AEF is equal to the angle EFD (by hyp.). Therefore the angle EFD is equal to the angle EFG (ax. 1); that is, a part is equal to the whole, which is impossible. Therefore no line but CD can be parallel to AB . Q. E. D.

Corol. Those lines which are perpendicular to the same line, are parallel to each other.


THEOREM



THEOREM XIV.

WHEN a Line cuts two Parallel Lines, the Outward Angle is Equal to the Inward Opposite one, on the Same Side; and the two Inward Angles, on the Same Side, equal to two Right Angles.

Let the line EF cut the two parallel lines AB, CD ; then will the outward angle EGB be equal to the inward opposite angle GHD , on the same side of the line EF ; and the two inward angles BGH, GHD , taken together, will be equal to two right angles.



For, since the two lines AB, CD , are parallel, the angle AGH is equal to the alternate angle GHD , (th. 12). But the angle AGH is equal to the opposite angle EGB (th. 7). Therefore the angle EGB is also equal to the angle GHD (ax. 1). Q. E. D.

Again, because the two adjacent angles EGB, BGH , are together equal to two right angles (th. 6); of which the angle EGB has been shown to be equal to the angle GHD ; therefore the two angles BGH, GHD , taken together, are also equal to two right angles.


Corol. 1. And, conversely, if one line meeting two other lines, make the angles on the same side of it equal, those two lines are parallels.

Corol. 2. If a line cutting two other lines, make the sum of the two inward angles, on the same side, less than two right angles, those two lines will not be parallel, but will meet each other when produced.

THEOREM XV.

THOSE Lines which are Parallel to the Same Line, are Parallel to each other.

Let the Lines AB, CD , be each of them parallel to the line EF ; then shall the lines AB, CD , be parallel to each other.




For, let the line GI be perpendicular to EF . Then will this line be also perpendicular to both the lines AB, CD , (corol. th. 12), and consequently the two lines AB, CD , are parallels (corol. th. 13). Q. E. D.

THEOREM XVI.

WHEN one Side of a triangle is produced, the Outward Angle is equal to both the Inward Opposite Angles taken together.

Let the side, AB , of the triangle ABC , be produced to D ; then will the outward angle CBD be equal to the sum of the two inward opposite angles A and C .




For, conceive BE to be drawn parallel to the side AC of the triangle. Then BC , meeting the two parallels AC , BE , makes the alternate angles C and CBE equal (th. 12). And AD , cutting the same two parallels AC , BE , makes the inward and outward angles on the same side, A and EBD , equal to each other (th. 14). Therefore, by equal additions, the sum of the two angles A and C , is equal to the sum of the two CBE and EBD , that is, to the whole angle CBD (by ax. 2). Q. E. D.

THEOREM XVII.

IN any Triangle, the sum of all the Three Angles is equal to Two Right Angles.

Let ABC be any plane triangle; then the sum of the three angles $A + B + C$ is equal to two right angles.

For, let the side AB be produced to D . Then the outward angle CBD is equal to the sum of the two inward opposite angles $A + C$ (th. 16). To each of these equals add the inward angle B , then will the sum of the three inward angles $A + B + C$ be equal to the sum of the two adjacent angles $ABC + CBD$ (ax. 2). But the sum of these two last adjacent angles is equal to two right angles (th. 6). Therefore also the sum of the three angles of the triangle $A + B + C$ is equal to two right angles (ax. 1). Q. E. D.



Corol. 1. If two angles in one triangle, be equal to two angles in another triangle, the third angles will also be equal (ax. 3), and the two triangles equiangular.

Corol. 2. If one angle in one triangle be equal to one angle in another, the sums of the remaining angles will also be equal (ax. 3).

Corol.

Corol. 3. If one angle of a triangle be right, the sum of the other two will also be equal to a right angle, and each of them singly will be acute, or less than a right angle.


Corol. 4. The two least angles of every triangle are acute, or each less than a right angle.

THEOREM XVIII.

IN any Quadrangle, the sum of all the Four Inward Angles, is equal to Four Right Angles.

Let $ABCD$ be a quadrangle; then the sum of the four inward angles, $A + B + C + D$ is equal to four right angles.

Let the diagonal AC be drawn, dividing the quadrangle into two triangles, ABC , ADC . Then, because the sum of the three angles of each of these triangles is equal to two right angles (th. 17); it follows, that the sum of all the angles of both triangles, which make up the four angles of the quadrangle, must be equal to four right angles (ax. 2). Q. E. D.



Corol. 1. Hence, if three of the angles be right ones, the fourth will also be a right angle.


Corol. 2. And, if the sum of two of the four angles be equal to two right angles, the sum of the remaining two will also be equal to two right angles.

THEOREM XIX.

IN any figure whatever, the Sum of all the Inward Angles, taken together, is equal to Twice as many Right Angles, wanting four, as the Figure has Sides.

Let $ABCDE$ be any figure; then the sum of all its inward angles, $A + B + C + D + E$, is equal to twice as many right angles, wanting four, as the figure has sides.

For, from any point P , within it, draw lines PA , PB , PC , &c. to all the angles, dividing the polygon into as many triangles as it has sides. Now the sum of the three angles of each of these triangles, is equal to two right angles (th. 17); therefore the sum of the angles of all the triangles is equal to twice as many right angles as the figure has sides. But the sum of all the angles about the point P , which are so many




many of the angles of the triangles, but no part of the inward angles of the polygon, is equal to four right angles (corol. 3, th. 6), and must be deducted out of the former sum. Hence it follows that the sum of all the inward angles of the polygon alone $A+B+C+D+E$, is equal to twice as many right angles as the figure has sides, wanting the said four right angles.* Q. E. D.

THEOREM XX.

WHEN every Side of any Figure is produced out, the Sum of all the Outward Angles thereby made, is equal to Four Right Angles.

Let $A, B, C, \&c.$ be the outward angles of any polygon, made by producing all the sides; then will the sum $A+B+C+D+E$, of all those outward angles, be equal to four right angles.

For every one of these outward angles, together with its adjacent inward angle, make up two right angles, as $A+a$ equal to two right angles, being the two angles made by one line meeting another (th. 6). And there being as many outward, or inward angles, as the figure has sides: therefore the sum of all the inward and outward angles, is equal to twice as many right angles as the figure has sides. But the sum of all the inward angles, with four right angles, is equal to twice as many right angles as the figure has sides (th. 19). Therefore the sum of all the inward and all the outward angles, is equal to the sum of all the inward angles and four right angles (by ax. 1). From each of these take away all the inward angles, and there remain all the outward angles equal to four right angles (by ax. 3).




THEOREM XXI.

A PERPENDICULAR is the Shortest Line that can be drawn from a Given Point to an Indefinite Line. And, of any other Lines drawn from the same Point, those that are Nearest the Perpendicular, are less than those More Remote.

If $AB, AC, AD, \&c.$ be lines drawn from the given point A , to the indefinite line DE , of which AB is perpendicular. Then shall the perpendicular AB be less than AC , and AC , less than $AD, \&c.$

For the angle B being a right one, the



* This demonstration does not apply to all rectilineal figures. Ed.

angle c is acute (by cor. 3, th. 17), and therefore less than the angle B . But the less angle of a triangle is subtended by the less side (th. 9). Therefore the side AB is less than the side AC .


Again, the angle ACB being acute, as before, the adjacent angle ACD will be obtuse (by th. 6); consequently the angle D is acute (corol. 3, th. 17), and therefore is less than the angle c . And since the less side is opposite to the less angle, therefore the side AC is less than the side AD . Q. E. D.

Corol. A perpendicular is the least distant of a given point from a line.

THEOREM XXII.

THE Opposite Sides and Angles of any Parallelogram are equal to each other; and the Diagonal divides it into two Equal Triangles.

Let $ABCD$ be a parallelogram, of which the diagonal is BD ; then will its opposite sides and angles be equal to each other, and the diagonal BD will divide it into two equal parts, or triangles.



For, since the sides AB and DC are parallel, as also the sides AD and BC (defin. 32), and the line BD meets them; therefore the alternate angles are equal (th. 12), namely the angle ABD to the angle CDB , and the angle ADB to the angle CBD . Hence the two triangles, having two angles in the one equal to two angles in the other, have also their third angles equal (cor. 1, th. 17), namely, the angle A equal to the angle C , which are two of the opposite angles of the parallelogram.

Also, if to the equal angles ABD , CDB , be added the equal angles CBD , ADB , the wholes will be equal (ax. 2), namely, the whole angle ABC to the whole ADC , which are the other two opposite angles of the parallelogram. Q. E. D.

Again, since the two triangles are mutually equiangular, and have a side in each equal, viz. the common side BD ; therefore the two triangles are identical (th. 2), or equal in all respects, namely, the side AB equal to the opposite side DC , and AD equal to the opposite side BC , and the whole triangle ABD equal to the whole triangle BCD . Q. E. D.

Corol.


Corol. 1. Hence, if one angle of a parallelogram be a right angle, all the other three will also be right angles, and the parallelogram a rectangle.

Corol. 2. Hence also, the sum of any two adjacent angles of a parallelogram is equal to two right angles.

THEOREM XXIII.

EVERY Quadrilateral, whose Opposite Sides are equal, is a Parallelogram, or has its Opposite Sides Parallel.

Let $ABCD$ be a quadrangle, having the opposite sides equal, namely, the side AB equal to DC , and AD equal to BC ; then shall these equal sides be also parallel, and the figure a parallelogram.



For, let the diagonal BD be drawn. Then, the triangles, ABD , CBD , being mutually equilateral (by hyp.), they are also mutually equiangular (th. 5), or have their corresponding angles equal; consequently the opposite sides are parallel (th. 13); viz. the side AB parallel to DC , and AD parallel to BC , and the figure is a parallelogram.

Q. E. D.

THEOREM XXIV.

THOSE Lines which join the Corresponding Extremes of two Equal and Parallel Lines, are themselves Equal and Parallel.

Let AB , DC , be two equal and parallel lines; then will the lines AD , BC , which join their extremes, be also equal and parallel. [See the fig. above.]


For, draw the diagonal BD . Then, because AB and DC are parallel (by hyp.), the angle ABD is equal to the alternate angle BDC (th. 12). Hence then, the two triangles having two sides and the contained angles equal, viz. the side AB equal to the side DC , and the side BD common, and the contained angle ABD equal to the contained angle BDC , they have the remaining sides and angles also respectively equal (th. 1); consequently AD is equal to BC , and also parallel to it (th. 12). Q. E. D.

THEOREM XXV.

PARALLELOGRAMS, as also Triangles, standing on the Same Base, and between the Same Parallels, are equal to each other.

Let

Let $ABCD$, $ABEF$, be two parallelograms, and ABC , ABF , two triangles, standing on the same base AB , and between the same parallels AB , DE ; then will the parallelogram $ABCD$, be equal to the parallelogram $ABEF$, and the triangle ABC equal to the triangle ABF .



For, since the line DE cuts the two parallels AF , BE , and the two AD , BC , it makes the angle E equal to the angle AFD , and the angle D equal to the angle BCE (th. 14); the two triangles ADF , BCF , are therefore equiangular (cor. 1, th. 17); and having the two corresponding sides AD , BC , equal (th. 22), being opposite sides of a parallelogram, these two triangles are identical, or equal in all respects (th. 2). If each of these equal triangles then be taken from the whole space $ABED$, there will remain the parallelogram $ABEF$ in the one case, equal to the parallelogram $ABCD$ in the other (by ax. 3).

Also the triangles ABC , ABF , on the same base AB , and between the same parallels, are equal, being the halves of the said equal parallelograms (th. 22). Q. E. D.


Corol. 1. Parallelograms, or triangles, having the same base and altitude, are equal. For the altitude is the same as the perpendicular or distance between the two parallels, which is every where equal, by the definition of parallels.

Corol. 2. Parallelograms, or triangles, having equal bases and altitudes, are equal. For, if the one figure be applied with its base on the other, the bases will coincide or be the same, because they are equal: and so the two figures, having the same base and altitude, are equal.

THEOREM XXVI.

IF a Parallelogram and a Triangle stand on the Same Base, and between the Same Parallels, the Parallelogram will be Double the Triangle, or the Triangle Half the Parallelogram.

Let $ABCD$ be a parallelogram, and ABE , a triangle, on the same base AB , and between the same parallels AB , DE ; then will the parallelogram $ABCD$ be double the triangle ABE , or the triangle half the parallelogram.



For, draw the diagonal ac of the parallelogram, dividing it into two equal parts (th. 22). Then because the triangles ABC ,

ABE ,

ABE , on the same base, and between the same parallels, are equal (th. 25); and because the one triangle ABC is half the parallelogram $ABCD$ (th. 22). the other equal triangle ABE is also equal to half the same parallelogram $ABCD$. Q. E. D.

Corol. 1. A triangle is equal to half a parallelogram of the same base and altitude, because the altitude is the perpendicular distance between the parallels which is every where equal, by the definition of parallels.

Corol. 2. If the base of a parallelogram be half that of a triangle, of the same altitude, or the base of the triangle be double that of the parallelogram, the two figures will be equal to each other.

THEOREM XXVII.

RECTANGLES that are contained by Equal Lines, are Equal to each other.

Let BD , FH , be two rectangles, having the sides AB , BC , equal to the sides EF , FG , each to each; then will the rectangle BD be equal to the rectangle FH .




For, draw the two diagonals AC , EG , dividing the two parallelograms each into two equal parts. Then the two triangles ABC , EFG , are equal to each other (th. 1), because they have the two sides AB , BC , and the contained angle B , equal to the two sides EF , FG , and the contained angle F (by hyp). But these equal triangles are the halves of the respective rectangles. And because the halves, or the triangles, are equal, the wholes, or the rectangles, DB , HF , are also equal (by ax. 6). Q. E. D.

Corol. The squares on equal lines are also equal; for every square is a species of rectangle.

THEOREM XXVIII.

THE Complements of the Parallelograms, which are about the diagonal of any Parallelogram, are equal to each other.

Let AC be a parallelogram, BD a diagonal, EIF parallel to AB or DC , and GIH parallel to AD or BC , making AI , IC complements to the parallelograms EG , HF , which are about the diagonal DB : then will the complement AI be equal to the complement IC .



For,

For, since the diagonal DE bisects the three parallelograms AC , EG , HF , (th. 22); therefore, the whole triangle DAB being equal to the whole triangle DCB , and the parts DEI , IHB , respectively equal to the parts DGI , IFB , the remaining parts AI , IC , must also be equal (by ax. 3). Q. E. D.

THEOREM XXIX.

A TRAPEZOID, or Trapezium having two Sides Parallel, is equal to Half a Parallelogram, whose Base is the Sum of those two Sides, and its Altitude the Perpendicular Distance between them.

Let $ABCD$ be the trapezoid, having its two sides AB , DC , parallel; and in AB produced take BE equal to DC , so that AE may be the sum of the two parallel sides; produce DC also, and let EF , GC , BH , be all three parallel to AD . Then is AE a parallelogram of the same altitude with the trapezoid $ABCD$, having its base AE equal to the sum of the parallel sides of the trapezoid; and it is to be proved that the trapezoid $ABCD$ is equal to half the parallelogram AE .




Now, since triangles, or parallelograms, of equal bases and altitude, are equal (corol. 2, th. 25), the parallelogram DG is equal to the parallelogram HE , and the triangle CGB equal to the triangle CHB ; consequently the line BE bisects, or equally divides, the parallelogram AE , and $ABCD$ is the half of it.

Q. E. D.

THEOREM XXX.


THE Sum of all the Rectangles contained under one Whole Line, and the several Parts of another Line, any way divided, is Equal to the Rectangle contained under the Two Whole Lines.

Let AD be the one line, and AB the other, divided into the parts AE , EF , FB ; then will the rectangle contained by AD and AB , be equal to the sum of the rectangles of AD and AE , and AD and EF , and AD and FB : thus expressed, $AD \cdot AB = AD \cdot AE + AD \cdot EF + AD \cdot FB$.



For, make the rectangle AC of the two whole lines AD , AB ; and draw EG , FH , perpendicular to AB , or parallel to AD , to which they are equal (th. 22). Then the whole rectangle AC is made up of all the other rectangles AG , FH , FC . But

these rectangles are contained by AD and AE, EG and EF, FH and FB; which are equal to the rectangles of AD and AE, AD and EF, AD and FB, because AD is equal to each of the two, EG, FH. Therefore the rectangle AD . AB is equal to the sum of all the other rectangles AD . AE, AD . EF, AD . FB. Q. E. D.




Corol. If a right line be divided into any two parts; the square on the whole line, is equal to both the rectangles of the whole line and each of the parts.

THEOREM XXXI.

THE SQUARE of the Sum of two Lines is greater than the Sum of their Squares, by Twice the Rectangle of the said Lines. Or, the Square of a whole Line, is equal to the Squares of its two Parts, together with Twice the Rectangle of those Parts.

Let the line AB be the sum of any two lines AC, CB: then will the square of AB be equal to the squares of AC, CB, together with twice the rectangle of AC . CB. That is, $AB^2 = AC^2 + CB^2 + 2AC . CB$.



For, let ABDE be the square on the sum of whole line AB, and ACFG the square on the part AC. Produce CF and GF to the other side at H and I.

From the lines CH, GI, which are equal, being each equal to the sides of the square AB or ED (th. 22), take the parts CF, GF, which are also equal, being the sides of the square AF, and there remains FH equal to FI, which are also equal to DH, DI, being the opposite sides of a parallelogram. Hence the figure HI is equilateral: and it has all its angles right ones (corol. 1, th. 22); it is therefore a square on the line FI, or the square of its equal CB. Also the figures EF, FB, are equal to two rectangles under AC and CB, because GF is equal to AC, and FH or FI equal to CB. But the whole square AD is made up of the four figures, viz. the two squares AF, FD, and the two equal rectangles EF, FB. That is, the square of AB is equal to the squares of AC, CB, together with twice the rectangle of AC, CB. Q. E. D.


Corol. Hence, if a line be divided into two equal parts; the square of the whole line, will be equal to four times the square of half the line.

THEOREM

THEOREM XXXII.

THE Square of the Difference of two Lines, is less than the Sum of their Squares, by Twice the Rectangle of the said Lines.

Let AC, BC, be any two lines, and AB their difference: then will the square of AB be less than the squares of AC, BC, by twice the rectangle of AC and BC. Or, $AB^2 = AC^2 + BC^2 - 2AC \cdot BC$.



For let ABDE be the square on the difference AB, and ACFG the square on the line AC. Produce ED to H; also produce DB and HC, and draw KI, making BK the square of the other line BC.

Now it is visible that the square AD is less than the two squares AF, BK, by the two rectangles EF, DI. But EF is equal to the one line AC, and GE or FH is equal to the other line BC; consequently the rectangle EF, contained under EG and GF, is equal to the rectangle of AC and BC.

Again, FH being equal to CI or BC or DH, by adding the common part HC, the whole HI will be equal to the whole FC, or equal to AC; and consequently the figure DI is equal to the rectangle contained by AC and BC.

Hence the two figures EF, DI, are two rectangles of the two lines AC, BC; and consequently the square of AB is less than the squares of AC, BC, by twice the rectangle AC . BC. Q. E. D.


THEOREM XXXIII.

THE Rectangle under the Sum and Difference of two Lines, is equal to the Difference of the Squares of those Lines.

Let AB, AC, be any two unequal lines; then will the difference of the squares of AB, AC, be equal to a rectangle under their sum and difference. That is,

$$AB^2 - AC^2 = \overbrace{AB+AC} \cdot \overbrace{AB-AC}$$

For, let ABDE be the square of AB, and ACFG the square of AC. Produce DB till BH be equal to AC; draw HI parallel to AB or ED, and produce FC both ways to I and K.



Then the difference of the two squares AD, AF, is evidently


dently the two rectangles EF, KB . But the rectangles EF, BI , are equal, being contained under equal lines; for EK and BH are each equal to AC , and GE is equal to CB , being each equal to the difference between AB and AC , or their equals AE and AG . Therefore the two EF, KB , are equal to the two KB, BI , or to the whole KH ; and consequently KH is equal to the difference of the squares AD, AF . But KH is a rectangle contained by DH , or the sum of AB and AC , and by KD , or the difference of AB and AC . Therefore the difference of the squares of AB, AC , is equal to the rectangle under their sum and difference. Q. E. D.

THEOREM XXXIV.

In any Right-angled Triangle, the square of the Hypotenuse, is equal to the Sum of the Squares of the other two Sides.

Let ABC be a right-angled triangle, having the right angle c ; then will the square of the hypotenuse AB , be equal to the sum of the squares of the other two sides AC, CB . Or $AB^2 = AC^2 + BC^2$.

For, on AB describe the square AE , and on AC, CB , the squares AG, BH ; then draw CK parallel to AD or BE ; and join AI, BF, CD, CE .



Now, because the line AC meets the two CG, CB , so as to make two right angles, these two form one straight line GB (corol. 1, th. 6). And because the angle FAC is equal to the angle DAB , being each a right angle, or the angle of a square; to each of these equals add the common angle BAC , so will the whole angle or sum FAB , be equal to the whole angle or sum CAD . But the line FA is equal to the line AC , and the line AB to the line AD , being sides of the same square; so that the two sides FA, AB , and their included angle FAB , are equal to the two sides CA, AD , and the contained angle CAD each to each; therefore the whole triangle AFB is equal to the whole triangle ACD (th. 1).

But the square AG is double the triangle AFB , on the same base FA , and between the same parallels FA, GB (th. 26); in like manner, the parallelogram AK is double the triangle ACD , on the same base AD , and between the same parallels AD, CK . And since the doubles of equal things are equal, (by ax. 6); therefore the square AG is equal to the parallelogram AK .

In like manner, the other square BH is proved equal to the other parallelogram BK . Consequently the two squares AG and BH together, are equal to the two parallelograms AK and BK together, or to the whole square AE . That is, the sum of the two squares on the two less sides, is equal to the square on the greatest side. Q. E. D.


Corol. 1. Hence, the square of either of the two less sides, is equal to the difference of the squares of the hypotenuse and the other side (ax. 3); or, equal to the rectangle contained by the sum and difference of the said hypotenuse and other side (th. 33).

Corol. 2. Hence also, if two right-angled triangles have two sides of the one equal to two corresponding sides of the other; their third sides will also be equal, and the triangles identical.

THEOREM. XXXV.

In any Triangle, the Difference of the Squares of the two Sides, is Equal to the Difference of the Squares of the Segments of the Base, or of the two Lines, or Distances, included between the Extremes of the Base and the Perpendicular.

Let ABC be any triangle, having CD perpendicular to AB ; then will the difference of the squares of AC , BC , be equal to the difference of the squares of AD , BD ; that is, $AC^2 - BC^2 = AD^2 - BD^2$



For, since AC^2 is equal to $AD^2 + CD^2$ } (by th. 34);
and BC^2 is equal to $BD^2 + CD^2$. }

Theref. the difference between AC^2 and BC^2 ,
is equal to the difference between $AD^2 + CD^2$
and $BD^2 + CD^2$,
or equal to the difference between AD^2 and BD^2 ,
by taking away the common square CD^2

Q. E. D.

Corol. The rectangle of the sum and difference of the two sides of any triangle, is equal to the rectangle of the sum and difference of the distances between the perpendicular and the two extremes of the base, or equal to the rectangle of the base and the difference or sum of the segments, according as the perpendicular falls within or without the triangle.

That

That is, $\overline{AC + BC} \cdot \overline{AC - BC} = \overline{AD + BD} \cdot \overline{AD - BD}$

Or, $\overline{AC + BC} \cdot \overline{AC - BC} = \overline{AB} \cdot \overline{AD - BD}$ in the 2d figure.

And $\overline{AC + BC} \cdot \overline{AC - BC} = \overline{AB} \cdot \overline{AD + BD}$ in the 1st figure.

THEOREM XXXVI.

IN any Obtuse-angled Triangle, the Square of the Side subtending the Obtuse Angle, is Greater than the Sum of the Squares of the other two Sides, by Twice the Rectangle of the Base and the Distance of the Perpendicular from the Obtuse Angle.

Let ABC be a triangle, obtuse angled at B, and CD perpendicular to AB; then will the square of AC be greater than the squares of AB, BC, by twice the rectangle of AB, BD. That is, $AC^2 = AB^2 + BC^2 + 2AB \cdot BD$. See the 1st fig. above, or below.


For, since the square of the whole line AD is equal to the squares of the parts AB, BD, with twice the rectangle of the same parts AB, BD (th. 31); if to each of these equals there be added the square of CD, then the squares of AD, CD, will be equal to the squares of AB, BD, CD, with twice the rectangle of AB, BD (by ax. 2).

But the squares of AD, CD, are equal to the square of AC; and the squares of BD, CD, equal to the square of BC (th. 34); therefore the square of AC is equal to the squares of AB, BC, together with twice the rectangle of AB, BD. Q. E. D.

THEOREM XXXVII.

IN any Triangle, the Square of the Side subtending an Acute Angle, is Less than the Squares of the Base and the other Side, by Twice the Rectangle of the Base and the Distance of the Perpendicular from the Acute Angle.

Let ABC be a triangle, having the angle A acute, and CD perpendicular to AB; then will the square of BC, be less than the squares of AB, AC, by twice the rectangle of AB, AD. That is, $BC^2 = AB^2 + AC^2 - 2AB \cdot AD$.



For.

For, in fig. 1, AC^2 is $= BC^2 + AB^2 + 2AB \cdot BD$ (th. 36).

To each of these equals add the square of AB ;
then is $AB^2 + AC^2 = BC^2 + 2AB^2 + 2AB \cdot BD$ (ax. 2),
or $= BC^2 + 2AB \cdot AD$ (th. 30).

Q. E. D.

Again, in fig. 2, AC^2 is $= AD^2 + DC^2$ (th. 34).

And $AB^2 = AD^2 + DB^2 + 2AD \cdot DB$ (th. 31).

Therefore $AB^2 + AC^2 = BD^2 + DC^2 + 2AD^2 + 2AD \cdot DB$ (ax. 2),

or $= BC^2 + 2AD^2 + 2AD \cdot DB$ (th. 34),


or $= BC^2 + 2AB \cdot AD$ (th. 30).

Q. E. D.

THEOREM XXXVIII.

In any Triangle, the Double of the Square of a Line drawn from the Vertex to the Middle of the Base, together with Double the Square of the Half Base, is equal to the Sum of the Squares of the other Two Sides.

Let ADC be a triangle, and CD the line drawn from the vertex to the middle of the base AB , dividing it into two equal parts AD, DB ; then will the sum of the squares of AC, CB , be equal to twice the sum of the squares of CD, BD ; or $AC^2 + CB^2 = 2CD^2 + 2DB^2$.




For, let CE be perpendicular to the base AB . Then, since (by th. 36) AC^2 exceeds the sum of the two squares AD^2 and CD^2 (or BD^2 and CD^2) by the double rectangle $2AD \cdot DE$ (or $2BD \cdot DE$); and since (by th. 37) BC^2 is less than the same sum by the said double rectangle; it is manifest that both AC^2 and BC^2 together must be equal to that sum twice taken; the excess on the one part making up the defect on the other.

Q. E. D.

THEOREM XXXIX.

In an Isosceles Triangle, the Square of a Line drawn from the Vertex to any Point in the Base, together with the Rectangle of the Segments of the Base, is equal to the Square of one of the Equal Sides of the Triangle.

Let ABC be the isosceles triangle, and CD a line drawn from the vertex to any point D in the base: then will the square of AC , be equal to the square of CD , together with the rectangle of AD and DB . That is, $AC^2 = CD^2 + AD \cdot DB$.



Q. E. D.

For, let CE bisect the vertical angle; then will it also bisect the base AB perpendicularly making $AE=EB$ (cor. 1, th. 3).


But, in the triangle ACD , obtuse angled at D , the square AC^2 is $= CD^2 + AD^2 + 2AD \cdot DE$ (th. 36),

$$\text{or } = CD^2 + AD \cdot \overline{AD + 2DE} \text{ (th. 30),}$$

$$\text{or } = CD^2 + AD \cdot \overline{AE + DE},$$

$$\text{or } = CD^2 + AD \cdot \overline{BE + DE},$$

$$\text{or } = CD^2 + AD \cdot \overline{DB}.$$




Q. E. D.

THEOREM XL.

In any Parallelogram, the two Diagonals Bisect each other; and the Sum of their Squares is equal to the Sum of the Squares of all the Four Sides of the Parallelogram.

Let $ABCD$ be a parallelogram, whose diagonals intersect each other in E ; then will AE be equal to EC , and BE to ED ; and the sum of the squares of AC , BD , will be equal to the sum of the squares of AB , BC , CD , DA . That is,



$$AE=EC, \text{ and } BE=ED,$$

$$\text{and } AC^2 + BD^2 = AB^2 + BC^2 + CD^2 + DA^2.$$

For, the triangles AEB , DEC , are equiangular, because they have the opposite angles at E equal (th. 7), and the two lines AC , BD , meeting the parallels AB , DC , make the angle BAE equal to the angle DCE , and the angle ABE equal to the angle CDE , and the side AB equal to the side DC (th. 22); therefore these two triangles are identical, and have their corresponding sides equal (th. 2), viz. $AE=EC$, and $BE=ED$.

Again, since AC is bisected in E , the sum of the squares $AD^2 + DC^2 = 2AE^2 + 2DE^2$ (th. 38).

$$\text{In like manner, } AB^2 + BC^2 = 2AE^2 + 2BE^2 \text{ or } 2DE^2.$$

$$\text{Theref. } AB^2 + BC^2 + CD^2 + DA^2 = 4AE^2 + 4DE^2 \text{ (ax. 2).}$$

But, because the square of a whole line is equal to 4 times the square of half the line (cor. th. 31), that is, $AC^2 = 4AE^2$, and $BD^2 = 4DE^2$.

$$\text{Theref. } AB^2 + BC^2 + CD^2 + DA^2 = AC^2 + BD^2 \text{ (ax. 1).}$$

Q. E. D.


THEOREM

THEOREM XLII.

IF a Line, drawn through or from the Centre of a Circle, Bisect a Chord, it will be Perpendicular to it ; or if it be Perpendicular to the Chord, it will Bisect both the Chord and the arc of the Chord.

Let AB be any chord in a circle, and CD a line drawn from the centre c to the chord. Then, if the chord be bisected in the point D , CD will be perpendicular to AB .

For, draw the two radii CA , CB . Then, the two triangles ACD , BCD , having CA equal to CB (def. 45), and CD common, also AD equal to BD (by hyp.) ; they have all the three sides of the one, equal to all the three sides of the other, and so have their angles also equal (th. 5). Hence then. the angle ADC being equal to the angle BDC , these angles are right angles, and the line CD is perpendicular to AB (def. 11).



Again, if CD be perpendicular to AB , then will the chord AB be bisected at the point D , or have AD equal to DB ; and the arc AEB bisected in the point E , or have AE equal EB .

For, having drawn CA , CB , as before. Then, in the triangle ABC , because the side CA is equal to the side CB , their opposite angles A and B are also equal (th. 3). Hence then, in the two triangles ACD , BCD , the angle A is equal to the angle B , and the angles at D are equal (def. 11) ; therefore their third angles are also equal (corol. 1, th. 17). And having the side CD common, they have also the side AD equal to the side DB (th. 2).

Also, since the angle ACE is equal to the angle BCE , the arc AE , which measures the former (def. 57), is equal to the arc BE , which measures the latter, since equal angles must have equal measures.


Corol. Hence a line bisecting any chord at right angles, passes through the centre of the circle.

THEOREM XLIII.

IF More than Two Equal Lines can be drawn from any Point within a Circle to the Circumference, that Point will be the centre.

Let ABC be a circle, and D a point within it: then if any three lines, DA , DB , DC , drawn from the point D to the circumference, be equal to each other, the point D will be the centre.

For, draw the chords AB , BC , which let be bisected in the point E , F , and join DE , DF .




Then, the two triangles, DAE , DBE , have the side DA equal to the side DB by supposition, and the side AE equal to the side EB by hypothesis, also the side DE common: therefore these two triangles are identical, and have the angles at E equal to each other (th. 5); consequently DE is perpendicular to the middle of the chord AB (def. 11), and therefore passes through the centre of the circle (corol. th. 41).

In like manner, it may be shown that DF passes through the centre. Consequently the point D is the centre of the circle, and the three equal lines, DA , DB , DC , are radii. Q. E. D.

THEOREM XLIII.

If two Circles touch one another Internally, the Centres of the Circles, and the Point of Contact will be all in the Same Right Line.

Let the two circles ABC , ADE , touch one another internally in the point A ; then will the point A and the centres of those circles be all in the same right line.



For, let F be the centre of the circle ABC , through which draw the diameter AFC . Then, if the centre of the other circle can be out of this line AC , let it be supposed in some other point as G ; through which draw the line FG cutting the two circles in B and D .

Now in the triangle AFG , the sum of the two sides FG , GA , is greater than the third side AF (th. 10), or greater than its equal radius FB . From each of these take away the common part FG , and the remainder GA will be greater than the remainder GB . But the point G being supposed the centre of the inner circle, its two radii, GA , GD , are equal to each other; consequently GD will also be greater than GB . But ADE being the inner circle, GD is necessarily less

less than GB . So that GD is both greater and less than GB ; which is absurd. Consequently the centre G cannot be out of the line AFC . Q. E. D.

THEOREM XLIV.

IF two Circles Touch one another Externally, the Centres of the Circles and the Point of Contact will be all in the Same Right Line.

Let the two circles ABC , ADE , touch one another externally at the point A ; then will the point of contact A and the centres of the two circles be all in the same right line.

For, let F be the centre of the circle ABC , through which draw the diameter AFC , and produce it to the other circle at E . Then, if the centre of the other circle ADE can be out of the line FE , let it, if possible, be supposed in some other point as G ; and draw the lines AG , FB , DG , cutting the two circles in B and D .

Then, in the triangle AFG , the sum of the two sides AF , AG , is greater than the third side FG (th. 10). But, F and G being the centres of the two circles, the two radii GA , GD , are equal, as are also the two radii AF , FB . Hence the sum of GA , AF , is equal to the sum of GD , BF ; and therefore this latter sum also. GD , BF , is greater than GF , which is absurd. Consequently the centre G cannot be out of the line EF .

Q. E. D.




THEOREM XLV.

ANY Chords in a Circle, which are Equally Distant from the Centre, are Equal to each other; or if they be Equal to each other, they will be Equally Distant from the Centre.


Let AB , CD , be any two chords at equal distances from the centre G ; then will these two chords AB , CD , be equal to each other.

For, draw the two radii GA , GC , and the two perpendiculars GE , GF , which are the equal distances from the centre G . Then, the two right-angled triangles, GAE , GCF , having the side GA equal the side GC , and the side GE equal the side GF , and the angle at E equal to the angle



at

at F , therefore the two triangles, GAE , GCF , are identical (cor. 2, th. 34), and have the line AE equal the line CF . But AB is the double of AE , and CD is the double of CF (th. 41); therefore AB is equal to CD (by ax. 6). Q. E. D.



Again, if the chord AB be equal to the chord CD : then will their distances from the centre, GE , GF , also be equal to each other.


For, since AB is equal CD by supposition, the half AE is equal the half CF . Also the radii GA , GC , being equal, as well as the right angles E and F , therefore the third sides are equal (cor. 2, th. 34), or the distance GE equal the distance GF .

Q. E. D.

THEOREM XLVI.

A Line Perpendicular to the Extremity of a Radius, is a Tangent to the Circle.

Let the line ADB be perpendicular to the radius CD of a circle; then shall AB touch the circle in the point D only.



For, from any other point E in the line AB draw CFE to the centre, cutting the circle in F .

Then, because the angle D , of the triangle CDE , is a right angle, the angle at E is acute (th. 17, cor. 3), and consequently less than the angle D . But the greater side is always opposite to the greater angle (th. 9); therefore the side CE is greater than the side CD , or greater than its equal CF . Hence the point E is without the circle; and the same for every other point in the line AB . Consequently the whole line is without the circle, and meets it in the point D only.

THEOREM

THEOREM XLVII.

WHEN a Line is a Tangent to a Circle, a Radius drawn to the Point of Contact is Perpendicular to the Tangent.

Let the line AB touch the circumference of a circle at the point D ; then will the radius CD be perpendicular to the tangent AB . [See the last figure.]

For, the line AB being wholly without the circumference except at the point D , every other, line, as CE drawn from the centre C to the line AB , must pass out of the circle to arrive at this line. The line CD is therefore the shortest that can be drawn from the point C to the line AB , and consequently (th. 21) it is perpendicular to that line.

Corol. Hence, conversely, a line drawn perpendicular to a tangent, at the point of contact, passes through the centre of the circle.

THEOREM XLVIII.


THE Angle formed by a Tangent and Chord is Measured by Half the Arc of that Chord.

Let AB be a tangent to a circle, and CD a chord drawn from the point of contact C ; then is the angle BCD measured by half the arc CFD , and the angle ACD measured by half the arc CGD .

For, draw the radius EC to the point of contact, and the radius EF perpendicular to the chord at H .

Then, the radius EF , being perpendicular to the chord CD , bisects the arc CFD (th. 41). Therefore CF is half the arc CFD .

In the triangle CEH , the angle H being a right one, the sum of the two remaining angles E and c is equal to a right angle (corol. 3, th. 17), which is equal to the angle BCE , because the radius CE is perpendicular to the tangent. From each of these equals take away the common part or angle c , and there remains the angle E equal to the angle BCD . But the angle E is measured by the arc CF (def. 57), which is the half of CFD ; therefore the equal angle BCD must also have the same measure, namely, half the arc CFD of the chord CD .



Again.

Again, the line GEF , being perpendicular to the chord CD , bisects the arc CGD , (th. 41). Therefore CG is half the arc CGD . Now, since the line CE , meeting FG , makes the sum of the two angles at E equal to two right angles (th. 6), and the line CD makes with AB the sum of the two angles at c equal to two right angles; if from these two equal sums there be taken away the parts or angles CEH and BCH which have been proved equal, there remains the angle CEG equal to the angle ACH . But the former of these, CEG , being an angle at the centre, is measured by the arc CG (def. 57); consequently the equal angle ACD must also have the same measure CG , which is half the arc CGD of the chord CD . Q. E. D.



Corol. 1. The sum of two right angles is measured by half the circumference. For the two angles BCD , ACD , which make up two right angles, are measured by the arcs, CF , CG , which make up half the circumference, FG being a diameter.

Corol. 2. Hence also one right angle must have for its measure a quarter of the circumference, or 90 degrees.

THEOREM XLIX.

An Angle at the Circumference of a Circle, is measured by Half the Arc that subtends it.

Let BAC be an angle at the circumference; it has for its measure, half the arc BC which subtends it.

For, suppose the tangent DE passing through the point of contact A . Then, the angle DAC being measured by half the arc ABC , and the angle DAB by half the arc AB (th. 48); it follows, by equal subtraction, that the difference, or angle BAC , must be measured by half the arc BC , which it stands upon. Q. E. D.




THEOREM L.

ALL Angles in the Same Segment of a Circle, or Standing on the Same Arc, are equal to each other.

Let c and d be two angles in the same segment $ACDB$, or, which is the same thing, standing on the supplemental arc AEB ; then will the angle c be equal to the angle d .

For each of these angles is measured by half the arc AEB ; and thus, having equal measures, they are equal to each other (ax. 11).




THEOREM LI.

AN Angle at the Centre of a Circle is double the Angle at the Circumference, when both stand on the Same Arc.

Let c be an angle at the centre c , and d an angle at the circumference, both standing on the same arc or same chord AB : then will the angle c be double of the angle d , or the angle d equal to half the angle c .

For, the angle at the centre c is measured by the whole arc AEB (def. 57), and the angle at the circumference d is measured by half the same arc AEB (th. 49); therefore the angle d is only half the angle c , or the angle c double the angle d .




THEOREM LII.

AN Angle in a Semicircle, is a Right Angle.

If ABC or ADC be a semicircle; then any angle d in that semicircle, is a right angle.

For, the angle d , at the circumference, is measured by half the arc ABC (th. 49), that is, by a quadrant of the circumference. But a quadrant is the measure of a right angle (corol. 4, th. 6; or corol. 2, th. 48),



Therefore the angle d is a right angle.

THEOREM LIII.

THE Angle formed by a Tangent to a Circle, and a Chord drawn from the Point of Contact, is Equal to the Angle in the Alternate Segment.

If AB be a tangent, and AC a chord, and D any angle in the alternate segment ADC ; then will the angle D be equal to the angle BAC made by the tangent and chord, of the arc AEC .




For the angle D , at the circumference, is measured by half the arc AEC (th. 49); and the angle BAC , made by the tangent and chord, is also measured by the same half arc AEC (th. 48); therefore these two angles are equal (ax. 11).

THEOREM LIV.

THE Sum of any Two Opposite Angles of a Quadrangle Inscribed in a Circle, is Equal to Two Right Angles.

Let $ABCD$ be any quadrilateral inscribed in a circle; then shall the sum of the two opposite angles A and C , or B and D , be equal to two right angles.




For the angle A is measured by half the arc DCB , which it stands on, and the angle C by half the arc DAB (th. 49); therefore the sum of the two angles A and C is measured by half the sum of these two arcs, that is, by half the circumference. But half the circumference is the measure of two right angles (corol. 4, th. 6); therefore the sum of the two opposite angles A and C is equal to two right angles. In like manner it is shown, that the sum of the other two opposite angles, D and B , is equal to two right angles.

Q. E. D.

THEOREM LV.

If any Side of a Quadrangle, Inscribed in a Circle, be Produced out, the Outward Angle will be Equal to the Inward Opposite Angle.

If the side AB , of the quadrilateral $ABCD$, inscribed in a circle, be produced to E ; the outward angle DAE will be equal to the inward opposite angle C .




For,

For, the sum of the two adjacent angles DAE and DAB is equal to two right angles (th. 6); and the sum of the two opposite angles c and DAB is also equal to two right angles (th. 54); therefore the former sum, of the two angles DAB and DAB , is equal to the latter sum, of the two c and DAE (ax. 1). From each of these equals taking away the common angle DAB , there remains the angle DAE equal the angle c .
Q. E. D.

THEOREM LVI.

Any Two Parallel Chords Intercept equal Arcs.

LET the two chords AB , CD , be parallel : then will the arcs AC , BD , be equal ; or $AC = BD$.




For, draw the line BC . Then, because the lines AB , CD , are parallel, the alternate angles B and c are equal (th. 12). But the angle at the circumference B , is measured by half the arc AC (th. 49); and the other equal angle at the circumference c is measured by half the arc BD : therefore the halves of the arcs AC , BD , and consequently the arcs themselves, are also equal. Q. E. D.

THEOREM LVII.

When a Tangent and Chord are Parallel to each other, they Intercept Equal Arcs.

LET the tangent ABC be parallel to the chord DF ; then are the arcs BD , BF , equal ; that is, $BD = BF$.




For, draw the chord BD . Then, because the lines AB , DF , are parallel, the alternate angles D and B are equal (th. 12). But the angle B , formed by a tangent and chord, is measured by half the arc BD (th. 48); and the other angle at the circumference D is measured by half the arc BF (th. 49); therefore the arcs BD , BF , are equal. Q. E. D.

THEOREM LVIII.

The Angle formed, Within a Circle, by the Intersection of two Chords, is Measured by Half the Sum of the Two Intercepted Arcs.

LET the two chords AB , CD , intersect at the point E : then the angle AEC , or DEB , is measured by half the sum of two arcs AC , DB .




For, draw the chord AF parallel to CD . Then, because the lines AF , CD , are parallel, and AB cuts them, the angles on the same side A and DEB are equal (th. 14). But the angle at the circumference A is measured by half the arc BF , or of the sum of FD and DB (th. 49); therefore the angle E is also measured by half the sum of FD and DB .

Again, because the chords AF , CD , are parallel, the arcs AC , FD , are equal (th. 56); therefore the sum of the two arcs AC , DB , is equal to the sum of the two FD , DB ; and consequently the angle E , which is measured by half the latter sum, is also measured by half the former. Q. E. D.

THEOREM LIX.

The Angle formed, Without a Circle, by two Secants, is Measured by Half the Difference of the Intercepted Arcs.

LET the angle E be formed by two secants EAB and ECD ; this angle is measured by half the difference of the two arcs AC , DB , intercepted by the two secants.



Draw the chord AF parallel to CD . Then, because the line AF , CD , are parallel, and AB cuts them, the angles on the same side A and BED are equal (th. 14). But the angle A , at the circumference, is measured by half the arc BF (th. 49), or of the difference of DF and DB : therefore the equal angle E is also measured by half the difference of DF , DB .

Again, because the chords AF , CD , are parallel, the arcs AC , FD , are equal (th. 56); therefore the difference of the two


two arcs AC, DB , is equal to the difference of the two DE, BE . Consequently the angle E , which is measured by half the latter difference, is also measured by half the former.

Q. E. D.

THEOREM LX.


The Angle formed by Two Tangents, is measured by Half the Difference of the two Intercepted Arcs.

Let EB, ED , be two tangents to a circle at the points A, C ; then the angle E is measured by half the difference of the two arcs, CFA, CGA .



For, draw the chord AF parallel to ED . Then, because the lines AF, ED , are parallel, and EB meets them, the angles on the same side A and E are equal (th. 14). But the angle A , formed by the chord AF and the tangent AB , is measured by half the arc AF (th. 48); therefore the equal angle E is also measured by half the same arc AF , or half the difference of the arcs CFA and CF , or CGA (th. 57).

Corol. In like manner it is proved, that the angle E formed by a tangent EDC , and a secant EAB , is measured by half the difference of the two intercepted arcs CA and CFB .



THEOREM LXI.

When two Lines, meeting a Circle each in two Points, Cut one another, either Within it or Without it; the Rectangle of the Parts of the one, is Equal to the Rectangle of the Parts of the other; the Parts of each being measured from the point of meeting to the two intersections with the circumference.

LET

LET the two lines, AB, CD, meet each other in E; then the rectangle of AE, EB, will be equal to the rectangle of CE, ED. Or, $AE \cdot EB = CE \cdot ED$.


For, through the point E draw the diameter FG; also, from the centre H draw the radius DH, and draw HI perpendicular to CD.

Then, since DEH is a triangle, and the perp. HI bisects the chord CD (th. 41), the line CE is equal to the difference of the segments DI, EI, the sum of them being DE. Also, because H is the centre of the circle and the radii DH, FH, GH, are all equal, the line EG is equal to the sum of the sides DH, HE; and EF is equal to their difference.

But the rectangle of the sum and difference of the two sides of a triangle, is equal to the rectangle of the sum and difference of the segments of the base (th. 35); therefore the rectangle of FE, EG, is equal to the rectangle of CE, ED. In like manner it is proved, that the same rectangle of FE, EG, is equal to the rectangle of AE, EB. Consequently the rectangle of AE, EB, is also equal to the rectangle of CE, ED (ax. 1). Q. E. D.

Corol. 1. When one of the lines in the second case, as DE, by revolving about the point E, comes into the position of the tangent EC or ED, the two points C and D running into one; then the rectangle of CE, ED, becomes the square of CE, because CE and DE are then equal. Consequently the rectangle of the parts of the secant, AE . EB, is equal to the square of the tangent CE².

Corol. 2. Hence both the tangents EC, EF, drawn from the same point E, are equal; since the square of each is equal to the same rectangle or quantity AE . EB.




THEOREM LXII.

In Equiangular Triangles, the Rectangles of the Corresponding or Like Sides, taken alternately, are equal.

LET

Let ABC , DEF , be two equiangular triangles, having the angle $A =$ the angle D , the angle $B =$ the angle E , and the angle $C =$ the angle F ; also the like sides AB , DE , and AC , DF , being those opposite the equal angles: then will the rectangle of AB , DF , be equal to the rectangle of AC , DE .



In BA produced take AG equal to DF ; and through the three points B , C , G , conceive a circle $BCGH$ to be described, meeting CA produced at H , and join GH .


Then the angle G is equal to the angle C on the same arc BH , and the angle H equal to the angle B on the same arc CG (th. 50); also the opposite angles at A are equal (th. 7): therefore the triangle AGH is equiangular to the triangle ACB , and consequently to the triangle DFE also. But the two like sides AG , DF , are also equal by supposition; consequently the two triangles AGH , DFE , are identical (th. 2.), having the two sides AG , AH , equal to the two DF , DE , each to each.

But the rectangle $CA \cdot AB$ is equal to the rectangle $HA \cdot AC$ (th. 61): consequently the rectangle $DF \cdot AB$ is equal the rectangle $DE \cdot AC$. Q. E. D.

THEOREM LXIII.

The Rectangle of the two Sides of any Triangle, is Equal to the Rectangle of the Perpendicular on the third Side and the Diameter of the Circumscribing Circle.

Let CD be the Perpendicular, and CE the diameter of the circle about the triangle ABC ; then the rectangle $CA \cdot CB$ is = the rectangle $CD \cdot CE$.




For, join BE : then in the two triangles ACD , ECB , the angles A and E are equal, standing on the same arc BC (th. 50): also the right angle D is equal to the angle B , which is also a right angle, being in a semicircle (th. 52): therefore these two triangles have also their third angles equal, and are equiangular. Hence, AC , CE , and CD , CB , being like sides, subtending the equal angles, the rectangle $AC \cdot CB$, of the first and last of them, is equal to the rectangle $CE \cdot CD$, of the other two (th. 62).

THEOREM

THEOREM LXIV.

The Square of a line bisecting any Angle of a Triangle, together with the Rectangle of the Two Segments of the opposite Side, is Equal to the Rectangle of the two other Sides including the Bisected Angle.

LET CD bisect the angle c of the triangle ABC ; then the square CD^2 + the rectangle $AD \cdot DB$ is = the rectangle $AC \cdot CB$.



For, let CD be produced to meet the circumscribing circle at E , and join AE .

Then the two triangles ACE , BCD , are equiangular: for the angles at c are equal by supposition, and the angles B and E are equal, standing on the same arc AC (th. 50); consequently the third angles at A and D are equal (corol. 1, th. 17); also AC , CD , and CE , CB , are like or corresponding sides, being opposite to equal angles: therefore the rectangle $AC \cdot CB$ is = the rectangle $CD \cdot CE$ (th. 62). But the latter rectangle $CD \cdot CE$ is = CD^2 + the rectangle $CD \cdot DE$ (th. 30); therefore also the former rectangle $AC \cdot CB$ is also = CD^2 + $CD \cdot DE$, or equal to CD^2 + $AD \cdot DB$, since $CD \cdot DE$ is = $AD \cdot DB$ (th. 61).

Q. E. D.

THEOREM LXV.

The Rectangle of the two Diagonals of any Quadrangle Inscribed in a Circle, is equal to the sum of the two Rectangles of the Opposite Sides.

LET $ABCD$ be any quadrilateral inscribed in a circle, and AC , BD , its two diagonals: then the rectangle $AC \cdot BD$ is = the rectangle $AB \cdot DC$ + the rectangle $AD \cdot BC$.



For, let CE be drawn, making the angle BCE equal to the angle DCA . Then the two triangles ACD , BCE , are equiangular; for the angles A and B are equal, standing on the same arc DC ; and the angles DCA , BCE , are equal by supposition; consequently the third angles ADC , BEC are also equal: also, AC , BC , and AD , BE , are like or corresponding sides, being opposite to the equal angles: therefore the rectangle $AC \cdot BE$ is = the rectangle $AD \cdot BC$ (th. 62).

Again,

Again, the two triangles ABC , DEC , are equiangular : for the angles BAC , BDC , are equal, standing on the same arc BC ; and the angle DCE is equal to the angle BCA , by adding the common angle ACE to the two equal angles DCA , BCE ; therefore the third angles E and ABC are also equal : but AC , DC , and AB , DE , are the like sides : therefore the rectangle $AC \cdot DE$ is = the rectangle $AB \cdot DC$ (th. 62).

Hence, by equal additions, the sum of the rectangles $AC \cdot BE + AC \cdot DE$ is = $AD \cdot BC + AB \cdot DC$. But the former sum of the rectangles $AC \cdot BE + AC \cdot DE$ is = the rectangle $AC \cdot BD$ (th. 30) : therefore the same rectangle $AC \cdot BD$ is equal to the latter sum, the rect. $AD \cdot BC +$ the rect. $AB \cdot DC$ (ax. 1).

Q. E. D.

OF RATIOS AND PROPORTIONS.

DEFINITIONS.

DEF. 76. **RATIO** is the relation which one magnitude bears to another magnitude of the same kind with respect to quantity.

The quantity or measure of a ratio is expressed by dividing the leading quantity or antecedent by the following or consequent. Thus the ratio of 6 to 2 is $\frac{6}{2} = 3$, the ratio of 20 to 8 is $\frac{20}{8} = \frac{5}{2} = 2\frac{1}{2}$, the ratio of 2 to 6 is $\frac{2}{6} = \frac{1}{3}$, and the ratio of 8 to 20 is $\frac{8}{20} = \frac{2}{5}$.

77. Proportion is an equality of ratios. Thus,

78. Three quantities are said to be Proportional, when the ratio of the first to the second, is equal to the ratio of the second to the third. As of the three quantities A (2), B (4), C (8), where $\frac{2}{4} = \frac{4}{8} = \frac{1}{2}$, both the same ratio.

79. Four quantities are said to be Proportional, when the ratio of the first to the second, is the same as the ratio of the third to the fourth. As of the four, A (2), B (4), C (5), D (10), where $\frac{2}{4} = \frac{5}{10} = \frac{1}{2}$, both the same ratio.

Note.

Note. To denote that four quantities, A, B, C, D , are proportional, they are usually stated or placed thus, $A : B :: C : D$; and read thus, A is to B as C is to D . But when three quantities are proportional, the middle one is repeated, and they are written thus, $A : B :: B : C$.

80. Of three proportional quantities, the middle one is said to be a Mean Proportional between the other two; and the last, a Third Proportional to the first and second.

81. Of four proportional quantities, the last is said to be a Fourth Proportional to the other three, taken in order.

82. Quantities are said to be Continually Proportional, or in Continued Proportion, when the ratio is the same between every two adjacent terms, viz. when the first is to the second, as the second to the third, as the third to the fourth, as the fourth to the fifth, and so on, all in the same common ratio.

As in the quantities 1, 2, 4, 8, 16, &c.; where the common ratio is equal to 2.

83. Of any number of quantities, A, B, C, D , the ratio of the first, A , to the last D , is said to be Compounded of the ratios of the first to the second, of the second to the third, and so on to the last.

84. Inverse ratio is, when the antecedent is made the consequent, and the consequent the antecedent.—Thus, if $1 : 2 :: 3 : 6$; then inversely, $2 : 1 :: 6 : 3$.

85. Alternate proportion is, when antecedent is compared with antecedent, and consequent with consequent.—As, if $1 : 2 :: 3 : 6$; then, by alternation, or permutation, it will be $1 : 3 :: 2 : 6$.

86. Compounded ratio, is when the sum of the antecedent and consequent is compared, either with the consequent, or with the antecedent.—Thus, if $1 : 2 :: 3 : 6$, then by composition, $1+2 : 1 :: 3+6 : 3$, and $1+2 : 2 :: 3+6 : 6$.

87. Divided ratio, is when the difference of the antecedent and consequent is compared, either with the antecedent, or with the consequent.—Thus, if $1 : 2 :: 3 : 6$, then, by division, $2-1 : 1 :: 6-3 : 3$, and $2-1 : 2 :: 6-3 : 6$.

Note. The term Divided, or Division, here means subtracting, or parting; being used in the sense opposed to compounding, or adding, in def. 86.

THEOREM LXVI.

Equimultiples of any two Quantities have the same Ratio as the Quantities themselves.

LET A and B be any two quantities, and m_A, m_B , any equimultiples of them, m being any number whatever: then will m_A and m_B have the same ratio as A and B , or $A : B :: m_A : m_B$.

For $\frac{m_A}{m_B} = \frac{A}{B}$, the same ratio.

Corol. Hence, like parts of quantities have the same ratio as the wholes; because the wholes are equimultiples of the like parts, or A and B are like parts of m_A and m_B .

THEOREM LXVII.

If Four Quantities, of the Same Kind, be Proportionals; they will be in Proportion by Alternation or Permutation, or the Antecedents will have the Same Ratio as the Consequents.

LET $A : B :: m_A : m_B$; then will $A : m_A :: B : m_B$.

For $\frac{A}{m_A} = \frac{1}{m}$, and $\frac{B}{m_B} = \frac{1}{m}$, both the same ratio.

THEOREM LXVIII.

If Four Quantities be Proportional; they will be in Proportion by Inversion, or Inversely.

LET $A : B :: m_A : m_B$; then will $B : A :: m_B : m_A$.

For $\frac{m_B}{m_A} = \frac{B}{A}$, both the same ratio.

THEOREM LXIX.

If Four Quantities be Proportional; they will be in Proportion by Composition and Division.

LET $A : B :: m_A : m_B$;

Then will $B \pm A : A :: m_B \pm m_A : m_A$,

and $B \pm A : B :: m_B \pm m_A : m_B$,

For $\frac{m_B \pm m_A}{m_A} = \frac{B \pm A}{A}$; and $\frac{m_B \pm m_A}{m_B} = \frac{B \pm A}{B}$.

Corol. It appears from hence, that the Sum of the Greatest and Least of four proportional quantities, of the same kind, exceeds the Sum of the Two Means. For, since $A : A + B :: m_A : m_A + m_B$, where A is the least, and $m_A + m_B$ the greatest; then $\frac{m+1}{m+1} \cdot A + mB$, the sum of the greatest and least exceeds $\frac{m+1}{m+1} \cdot A + B$, the sum of the two means.

THEOREM LXX.

If, of Four Proportional Quantities, there be taken any Equimultiples whatever of the two Antecedents, and any Equimultiples whatever of the two Consequents; the quantities resulting will still be proportional.

LET $A : B :: m_A : m_B$; also, let p_A and $p m_A$ be any equimultiples of the two antecedents, and q_B and $q m_B$ any equimultiples of the two consequents; then will $p_A : q_B :: p m_A : q m_B$.

For $\frac{p m_A}{q m_B} = \frac{p_A}{q_B}$, both the same ratio.

THEOREM LXXI.

If there be Four Proportional Quantities, and the two Consequents be either Augmented or Diminished by Quantities that have the Same Ratio as the respective Antecedents; the Results and the Antecedents will still be Proportionals.

LET $A : B :: m_A : m_B$, and n_A and $n m_A$ any two quantities having the same ratio as the two antecedents; then will $A : B \pm n_A :: m_A : m_B \pm n m_A$.

For $\frac{m_A}{m_B \pm n m_A} = \frac{A}{B \pm n_A}$, both the same ratio.

THEOREM LXXII.

If any Number of Quantities be Proportional, then any one of the Antecedents will be to its Consequent, as the Sum of all the Antecedents is to the Sum of all the Consequents.

LET $A : B :: m_A : m_B :: n_A : n_B$, &c; then will $A : B :: A + m_A + n_A :: B + m_B + n_B$, &c.

For $\frac{A + m_A + n_A}{B + m_B + n_B} = \frac{A}{B}$, the same ratio.

THEOREM

THEOREM LXXIII.

If a Whole Magnitude be to a Whole, as a Part taken from the first, is to a Part taken from the other; then the Remainder will be to the Remainder, as the whole to the whole.

$$\text{Let } A : B :: \frac{m}{n} A : \frac{m}{n} B;$$

$$\text{then will } A : B :: A - \frac{m}{n} A : B - \frac{m}{n} B.$$

$$\text{For } \frac{A - \frac{m}{n} A}{B - \frac{m}{n} B} = \frac{A}{B}, \text{ both the same ratio.}$$

THEOREM LXXIV.

If any Quantities be Proportional; their Squares, or Cubes, or any Like Powers, or Roots, of them, will also be Proportional.

$$\text{LET } A : B :: m_A : m_B; \text{ then will } A^n : B^n :: m_A^n : m_B^n.$$

$$\text{For } \frac{m_A^n A^n}{m_B^n B^n} = \frac{A^n}{B^n}, \text{ both the same ratio.}$$

THEOREM LXXV.

If there be two Sets of Proportionals; then the Products or Rectangles of the Corresponding Terms will also be Proportional.

$$\text{LET } A : B :: m_A : m_B,$$

$$\text{and } C : D :: n_C : n_D;$$

$$\text{then will } AC : BD :: mnAC : mnBD.$$

$$\text{For } \frac{mnAC}{mnBD} = \frac{AC}{BD}, \text{ both the same ratio.}$$

THEOREM LXXVI.

If Four Quantities be Proportional; the Rectangle or Product of the two Extremes, will be Equal to the Rectangle or Product of the two Means. And the converse.

$$\text{LET } A : B :: m_A : m_B;$$

$$\text{then is } A \times m_B = B \times m_A = m_{AB}, \text{ as is evident.}$$

THEOREM LXXVII.

If Three Quantities be Continued Proportionals; the Rectangle or Product of the two Extremes, will be Equal to the Square of the Mean. And the converse.

LET A, mA, m^2A be three proportionals,

OR $A : mA :: mA : m^2A$;

then is $A \times m^2A = m^2A^2$, as is evident.

THEOREM LXXVIII.

If any Number of Quantities be Continued Proportionals; the Ratio of the First to the Third, will be duplicate or the Square of the Ratio of the First and Second; and the Ratio of the First and Fourth will be triplicate or the cube of that of the First and Second; and so on.

LET $A, mA, m^2A, m^3A, \&c.$ be proportionals;

that is $\frac{A}{mA} = \frac{1}{m}$, but $\frac{A}{m^2A} = \frac{1}{m^2}$, $\frac{A}{m^3A} = \frac{1}{m^3}$; &c.

THEOREM LXXIX.

Triangles, and also Parallelograms, having equal Altitudes, are to each other as their Bases.

LET the two triangles ADC, DEF , have the same altitude, or between the same parallels AE, CF ; then is the surface of the triangle ADC , to the surface of the triangle DEF , as the base AD is to the base DE . Or, $AD : DE ::$ the triangle ADC : the triangle DEF .



For, let the base AD be to the base DE , as any one number m (2), to any other number n (3); and divide the respective bases into those parts, AB, BD, DG, GH, HE , all equal to one another; and from the points of division draw the lines BC, CG, GH, HE , to the vertices C and F . Then will these lines divide the triangles ADC, DEF , into the same number of parts as their bases, each equal to the triangle ABC , because those triangular parts have equal bases and altitude (corol. 2. th. 25); namely, the triangle ABC , equal to each of the triangles EDC, DFG, GFH, HFE . So that the triangle ADC , is to the triangle DFE , as the number

number of parts m (2) of the former, to the number n (3) of the latter, that is, as the base AD to the base DE (def. 79.)

In like manner, the parallelogram $ADKI$ is to the parallelogram $DEFK$, as the base AD is to the base DE ; each of these having the same ratio as the number of their parts, m to n .


Q. E. D.

THEOREM LXXX.

Triangles, and also Parallelograms, having Equal Bases, are to each other as their Altitudes.

Let ABC , BEF , be two triangles having the equal bases AB , BE , and whose altitudes are the perpendiculars CG , FH ; then will the triangle ABC : the triangle BEF :: CG : FH :

For, let BK be perpendicular to AB , and equal to CG ; in which let there be taken $BL = FH$; drawing AK and AL .



Then, triangles of equal bases and heights being equal (corol. 2, th. 25), the triangle ABK is $= ABC$, and the triangle $ABL = BEF$. But considering now ABK , ABL , as two triangles on the bases BK , BL , and having the same altitude AB , these will be as their bases (th. 79), namely the triangle ABK : the triangle ABL :: BK : BL .

But the triangle $ABK = ABC$, and the triangle $ABL = BEF$, also $BK = CG$, and $BL = FH$.

Theref. the triangle ABC : triangle BEF :: CG : FH .

And since parallelograms are the doubles of these triangles, having the same bases and altitudes, they will likewise have to each other the same ratio as their altitudes. Q. E. D.

Corol. Since, by this theorem, triangles and parallelograms, when their bases are equal, are to each other as their altitudes; and by the foregoing one, when their altitudes are equal, they are to each other as their bases; therefore universally, when neither are equal, they are to each other in the compound ratio, or as the rectangle or product of their bases and altitudes.


THEOREM LXXXI.

If Four Lines be Proportional; the Rectangle of the Extremes will be equal to the Rectangle of the Means. And, conversely, if the Rectangle of the extremes, of four Lines, be equal to the Rectangle of the means, the four Lines, taken alternately, will be Proportional.

Let the four lines, A, B, C, D, be proportionals, or $A : B :: C : D$; then will the rectangle of A and D be equal to the rectangle of B and C; or the rectangle $A \cdot D = B \cdot C$.

For, let the four lines be placed with their four extremities meeting in a common point, forming at that

point four right angles; and draw lines parallel to them to complete the rectangles P, Q, R, where P is the rectangle of A and D, Q the rectangle of B and C, and R the rectangle of B and D.



Then the rectangles P and R, being between the same parallels, are to each other as their bases A and B (th. 79); and the rectangles Q and R, being between the same parallels, are to each other as their bases C and D. But the ratio of A to B, is the same as the ratio of C to D, by hypothesis; therefore the ratio of P to R, is the same as the ratio of Q to R; and consequently the rectangles P and Q are equal. Q. E. D.

Again, if the rectangle of A and D, be equal to the rectangle of B and C; these lines will be proportional, or $A : B :: C : D$.

For, the rectangles being placed the same as before: then, because parallelograms between the same parallels are to one another as their bases, the rectangle $P : R :: A : B$, and $Q : R :: C : D$. But as P and Q are equal, by supposition, they have the same ratio to R, that is, the ratio of A to B is equal to the ratio of C to D, or $A : B :: C : D$. Q. E. D.

Corol. 1. When the two means, namely, the second and third terms, are equal, their rectangle becomes a square of the second term, which supplies the place of both the second and third. And hence it follows, that when three lines are proportionals, the rectangle of the two extremes is equal to the

the square of the mean ; and, conversely, if the rectangle of the extremes be equal to the square of the mean, the three lines are proportionals.

Corol. 2. Since it appears, by the rules of proportion in Arithmetic and Algebra, that when four quantities are proportional, the product of the extremes is equal to the product of the two means ; and, by this theorem, the rectangle of the extremes is equal to the rectangle of the two means ; it follows, that the area or space of a rectangle is represented or expressed by the product of its length and breadth multiplied together. And, in general, a rectangle in geometry is similar to the product of the measures of its two dimensions of length and breadth, or base and height. Also, a square is similar to, or represented by, the measure of its side multiplied by itself. So that, what is shown of such products, is to be understood of the squares and rectangles.

Corol. 3. Since the same reasoning, as in this theorem, holds for any parallelograms whatever, as well as for the rectangles, the same property belongs to all kinds of parallelograms, having equal angles, and also to triangles, which are the halves of parallelograms ; namely, that if the sides about the equal angles of parallelograms or triangles, be reciprocally proportional, the parallelograms or triangles will be equal ; and, conversely, if the parallelograms or triangles be equal, their sides about the equal angles will be reciprocally proportional.


Corol. 4. Parallelograms, or triangles, having an angle in each equal, are in proportion to each other as the rectangles of the sides which are about these equal angles.

THEOREM LXXXII.

If a Line be drawn in a Triangle Parallel to one of its sides, it will cut the other Sides Proportionally.

Let DE be parallel to the side BC of the triangle ABC ; then will $AD : DB :: AE : EC$.


For draw BE and CD . Then the triangles DBE , DCE , are equal to each other, because they have the same base DE , and are between the same parallels DE , BC (th. 25). But the two triangles ADE , BDE , on the bases AD , DB , have the same alti-



tude ;

tude ; and the two triangles ADE, CDE, on the bases AE, EC, have also the same altitude ; and because triangles of the same altitude are to each other as their bases, therefore

the triangle ADE : BDE :: AD : DB,
and triangle ADE : CDE :: AE : EC.



But BDE is = CDE ; and equals must have to equals the same ratio ; therefore AD : DB :: AE : EC. Q. E. D.


Corol. Hence, also, the whole lines AB, AC, are proportional to their corresponding proportional segments (corol. th. 66).

VIZ. AB : AC :: AD : AE,
and AB : AC :: BD : CE.

THEOREM LXXXIII.

A Line which Bisects any Angle of a Triangle, divides the opposite Side into Two Segments, which are Proportional to the two other Adjacent Sides.

Let the angle ACB, of the triangle ABC, be bisected by the line CD, making the angle r equal to the angle s ; then will the segment AD be to the segment DB, as the side AC is to the side CB. Or, -----
AD : DB :: AC : CB.



For, let BE be parallel to CD, meeting AC produced at E. Then, because the line BC cuts the two parallels CD, BE, it makes the angle CBE equal to the alternate angle s (th. 12), and therefore also equal to the angle r , which is equal to s by the supposition. Again, because the line AE cuts the two parallels DC, BE, it makes the angle E equal to the angle r on the same side of it (th. 14). Hence, in the triangle BCE, the angles B and E, being each equal to the angle r , are equal to each other, and consequently their opposite sides CB, CE, are also equal (th. 3).


But now, in the triangle ABE, the line CD, being drawn parallel to the side BE, cuts the two other sides AB, AE proportionally (th. 82), making AD to DB, as is AC to CE or to its equal CB. Q. E. D.

THEOREM

THEOREM LXXXIV.

Equiangular Triangles are similar, or have their Like Sides Proportional.

LET ABC , DEF , be two equiangular triangles, having the angle A equal to the angle D , the angle B to the angle E , and consequently the angle C to the angle F ; then will $AB : AC :: DE : DF$.



For, make $DG = AB$, and $DH = AC$, and join GH . Then the two triangles ABC , DGH , having the two sides AB , AC , equal to the two DG , DH , and the contained angles A and D also equal, are identical, or equal in all respects (th. 1) namely the angles B and C are equal to the angles G and H . But the angles B and C are equal to the angles E and F by the hypothesis; therefore also the angles G and H are equal to the angles E and F (ax. 1) and consequently the line GH is parallel to the side EF (cor. 1, th. 14).


Hence then, in the triangle DEF , the line GH , being parallel to the side EF , divides the two other sides proportionally, making $DG : DH :: DE : DF$ (cor. th. 82.) But DG and DH are equal to AB and AC ; therefore also $AB : AC :: DE : DF$.

Q. E. D.

THEOREM LXXXV.

Triangles which have their Sides Proportional, are Equiangular.

In the two triangles ABC , DEF , if $AB :: DE :: AC : DF :: BC : EF$; the two triangles will have their corresponding angles equal.



For, if the triangle ABC be not equiangular with the triangle DEF , suppose some other triangle, as DEG , to be equiangular with ABC . But this is impossible: for if the two triangles ABC , DEG , were equiangular, their sides would be proportional (th. 84). So that, AB being to DE as AC to DG , and AB to DE as BC to EG , it follows that DG and EG being fourth proportionals to the same three quantities as well as the two DF , EF , the

former DG , EG , would be equal to the latter, DF , EF . Thus then, the two triangles, DEF , DEG , having their three sides equal, would be identical (th. 5); which is absurd, since their angles are unequal.

THEOREM LXXXVI.

Triangles, which have an Angle in the one Equal to an Angle in the other, and the Sides about these angles Proportional, are Equiangular.

LET ABC , DEF , be two triangles, having the angle $A =$ the angle D , and the sides AB , AC , proportional to the sides DE , DF : then will the triangle ABC be equiangular with the triangle DEF .

For, make $DG = AB$, and $DH = AC$, and join GH .

Then, the two triangles ABC , DGH , having two sides equal, and the contained angles A and D equal, are identical and equiangular (th. 1), having the angles G and H equal to the angles B and C . But, since the sides DG , DH , are proportional to the sides DE , DF , the line GH is parallel to EF (th. 82); hence the angles E and F are equal to the angles G and H (th. 14), and consequently to their equals B and C .



Q. E. D.

THEOREM LXXXVII.

In a Right-Angled Triangle, a Perpendicular from the Right Angle, is a Mean Proportional between the Segments of the Hypotenuse; and each of the Sides, about the Right Angle, is a Mean Proportional between the Hypotenuse and the adjacent segment.


Let ABC be a right-angled triangle, and CD a perpendicular from the right angle C to the hypotenuse AB ; then will

CD be a mean proportional between AD and DB ;

AC a mean proportional between AB and AD ;

BC a mean proportional between AB and BD .

Or, $AD : CD :: CD : DB$; and $AB : BC :: BC : BD$; and $AB : AC :: AC :$



For

For, the two triangles ABC , ADC , having the right angles at c and d equal, and the angle A common, have their third angles equal, and are equiangular (corol. 1, th. 17). In like manner, the two triangles ABC , BDC , having the right angles at c and d equal, and the angle B common, have the third angles equal, and are equiangular.

Hence then, all the three triangles ABC , ADC , BDC , being equiangular will have their like sides proportional (th. 84).

$$\begin{aligned} \text{viz. } AD : CD &:: CD : DB ; \\ \text{and } AB : AC &:: AC : AD ; \\ \text{and } AB : BC &:: BC : BD. \end{aligned}$$

Q. E. D.

Corol. Because the angle in a semicircle is a right angle (th. 52); it follows, that if, from any point c in the periphery of the semicircle, a perpendicular be drawn to the diameter AB ; and the two chords CA , CB , be drawn to the extremities of the diameter: then are AC , BC , CD , the mean proportionals as in this theorem, or (by th. 77), $CD^2 = AD \cdot DB$; $AC^2 = AB \cdot AD$; and $BC^2 = AB \cdot BD$.

THEOREM LXXXVIII.


Equiangular or Similar Triangles, are to each other as the Squares of their Like Sides.

Let ABC , DEF , be two equiangular triangles, AB and DE being two like sides; then will the triangle ABC be to the triangle DEF , as the square of AB is to the square of DE , or as AB^2 to DE^2 .

For, let AL and DN be the squares on AB and DE ; also draw their diagonals BK , EM , and the perpendiculars CG , FH , of the two triangles.

Then, since equiangular triangles have their like sides proportional (th. 84), in the two equiangular triangles ABC , DEF , the side $AC : DF :: AB : DE$; and in the two ACG , DFH , the side $AC : DF :: CG : FH$; therefore, by equality $CG : FH :: AB : DE$, or $CG : AB :: FH : DE$.

But because triangles on equal bases are to each other as their altitudes, the triangles ABC , ABK , on the same base AB , are to each other, as their altitudes CG , AK . or $AB :$
and



and the triangles DEF, DEM, on the same base DE, are as their altitudes FH, DM, or DE ;

that is, triangle ABC : triangle ABK :: CG : AB,

and triangle DEF : triangle DEM :: FH : DE.

But it has been shown that $CG : AB :: FH : DE$;

theref. of equality $\triangle ABC : \triangle ABK :: \triangle DEF : \triangle DEM$,

or alternately, as $\triangle ABC : \triangle DEF :: \triangle ABK : \triangle DEM$.

But the squares AL, DN, being the double of the triangles ABH, DEM, have the same ratio with them ;


therefore the $\triangle ABC : \triangle DEF :: \text{square AL} : \text{square DN}$.

Q. E. D.

THEOREM LXXXIX.

All Similar Figures are to each other, as the Squares of their Like Sides.

LET ABCDE, FGHIK, be any two similar figures, the like sides being AB, FG, and BC, GH, and so on in the same order : then will the figure ABCDE be to the figure FGHIK, as the square of AB to the square of FG, or as AB^2 to FG^2 .



For, draw BE, BD, GK, GI, dividing the figures into an equal number of triangles, by lines from two equal angles B and G.

The two figures being similar (by suppos.), they are equiangular, and have their like sides proportional (def. 70).

Then, since the angle A is = the angle F, and the sides AB, AE, proportional to the sides FG, FK, the triangles ABE, FGK, are equiangular (th 86). In like manner, the two triangles BCD, GHI, having the angle c = the angle H, and the sides BC, CD, proportional to the sides GH, HI, are also equiangular. Also, if from the equal angles AED, FKI, there be taken the equal angles AEB, FKG, there will remain the equals BED, GKI ; and if from the equal angles CDE, HIK, be taken away the equals CDE, HIG, there will remain the equals BDE, GIK ; so that the two triangles BDE, GIK, having two angles equal, are also equiangular. Hence each triangle of the one figure, is equiangular with each corresponding triangle of the other.

But equiangular triangles are similar, and are proportional to the squares of their like sides (th. 88).

Therefore


Therefore the $\triangle ABE : \triangle FGK :: AB^2 : FG^2$,
 and $\triangle BCD : \triangle GHI :: BC^2 : GH^2$;
 and $\triangle BDE : \triangle GIK :: DE^2 : IK^2$.

But as the two polygons are similar, their like sides are proportional. and consequently their squares also proportional: so that all the ratios, AB^2 to FG^2 , and BC^2 to GH^2 , and DE^2 to IK^2 , are equal among themselves, and consequently the corresponding triangles also, ABE to FGK , and BCD to GHI , and BDE to GIK , have all the same ratio, viz. that of AB^2 to FG^2 : and hence all the antecedents, or the figure $ABCDE$, have to all the consequents, or the figure $FGHIK$, still the same ratio, viz. that of AB^2 to FG^2 (th. 72). Q. E. D.

THEOREM XC.

Similar Figures Inscribed in Circles, have their Like Sides, and also their Whole Perimeters, in the Same Ratio as the Diameters of the Circles in which they are Inscribed.

Let $ABCDE$, $FGHIK$, be two similar figures, inscribed in the circles whose diameters are AL and FM ; then will each side AB , BC , &c. of the one figure be to the like side GF , GH , &c, of the other figure, or the whole perimeter $AB+BC+ \&c$, of the one figure, to the whole perimeter $FG+GH+ \&c$, of the other figure, as the diameter AL to the diameter FM .



For, draw the two corresponding diagonals AC , FH , as also the lines BL , GM . Then since the polygons are similar, they are equiangular, and their like sides have the same ratio (def 70); therefore the two triangles ABC , FGH , have the angle $B =$ the angle G , and the sides AB , BC , proportional to the two sides FG , GH , consequently these two triangles are equiangular (th. 86), and have the angle $ACB = FHG$. But the angle $ACB = ALB$, standing on the same arc AB ; and the angle $FHG = FMG$, standing on the same arc FG ; therefore the angle $ALB = FMG$ (th. 1). And since the angle $ABL = FGM$, being both right angles, because in a semicircle; therefore the two triangles ABL , FGM . having two angles equal. are equiangular; and consequently their like sides are proportional


portional (th. 84); hence $AB : FG ::$ the diameter AL : the diameter FM .

In like manner, each side BC , CD , &c. has to each side GH , HI , &c. the same ratio of AL to FM ; and consequently the sums of them are still in the same ratio; viz $AB + BC + CD$, &c. : $FG + GH + HI$, &c. :: the diam. AL : the diam. FM (th. 72). Q. E. D.

THEOREM XCI.

Similar Figures Inscribed in Circles, are to each other as the Squares of the Diameters of those Circles.

Let $ABCDE$, $FGHIK$, be two similar figures inscribed in the circles whose diameters are AL and FM ; then the surface of the polygon $ABCDE$ will be to the surface of the polygon $FGHIK$, as AL^2 to FM^2 .



For, the figures being similar, are to each other as the squares of their like sides, AB^2 to FG^2 (th. 88). But, by the last theorem, the sides AB , FG , are as the diameters AL , FM ; and therefore the squares of the sides AB^2 to FG^2 , as the squares of the diameters AL^2 to FM^2 (th. 74). Consequently the polygons $ABCDE$, $FGHIK$, are also to each other as the squares of the diameters AL^2 to FM^2 (ax. 1). Q. E. D.

THEOREM XCII.

The Circumferences of all Circles are to each other as their Diameters.

LET D , d , denote the diameters of two circles, and c , c , their circumferences?

then will $D : d :: c : c$, or $D : c :: d : c$.

For, (by theor. 90), similar polygons inscribed in circles have their perimeters, in the same ratio as the diameters of those circles.

Now, as this property belongs to all polygons, whatever the number of the sides may be; conceive the number of the sides to be indefinitely great, and the length of each indefinitely small, till they coincide with the circumference of the

the circle, and be equal to it, indefinitely near. Then the perimeter of the polygon of an infinite number of sides, is the same thing as the circumference of the circle. Hence it appears that the circumferences of the circles, being the same as the perimeters of such polygons, are to each other in the same ratio as the diameters of the circles. Q. E. D.

THEOREM XCIII.

The Areas or Spaces of Circles, are to each other as the Squares of their Diameters, or of their Radii.

LET A a , denote the areas or spaces of two circles, and d d , their diameters; then $A : a :: d^2 : d^2$.

For (by theorem 91) similar polygons inscribed in circles are to each other as the squares of the diameters of the circles.


Hence, conceiving the number of the sides of the polygons to be increased more and more, or the length of the sides to become less and less, the polygon approaches nearer and nearer to the circle, till at length, by an infinite approach, coincide, and become in effect equal; and then it follows that the spaces of the circles, which are the same as of the polygons, will be to each other as the squares of the diameters of the circles. Q. E. D.

Corol. The spaces of circles are also to each other as the squares of the circumferences; since the circumferences are in the same ratio as the diameters (by theorem 92).

THEOREM XCIV.

The Area of any Circle, is Equal to the Rectangle of Half its Circumference and half its Diameter.

CONCEIVE a regular polygon to be inscribed in the circle: and radii drawn to all the angular points, dividing it into as many equal triangles as the polygon has sides, one of which ABC , of which the altitude is the perpendicular CD from the centre to the base AB .



Then the triangle ABC , being equal to a rectangle of half the base and equal altitude (th. 26, cor. 2), is equal to the rectangle of the half base AD and the altitude CD ;

conse-

consequently the whole polygon, or all the triangles added together which compose it, is equal to the rectangle of the common altitude CD , and the halves of all the sides, or the half perimeter of the polygon.



Now, conceive the number of sides of the polygon to be indefinitely increased; then will its perimeter coincide with the circumference of the circle, and consequently the altitude CD will become equal to the radius, and the whole polygon equal to the circle. Consequently the space of the circle, or of the polygon in that state, is equal to the rectangle of the radius and half the circumference. Q. E. D.

OF PLANES AND SOLIDS.

DEFINITIONS.

DEF. 88. The common Section of two Planes, is the line in which they meet, to cut each other.

89. A line is Perpendicular to a Plane, when it is perpendicular to every line in that plane which meets it.

90. One Plane is Perpendicular to Another, when every line of the one, which is perpendicular to the line of their common section, is perpendicular to the other.

91. The inclination of one Plane to another, or the angle they form between them, is the angle contained by two lines drawn from any point in the common section, and at right angles to the same, one of these lines in each plane.

92. Parallel Planes, are such as being produced ever so far both ways, will never meet, or which are every where at an equal perpendicular distance :

93. A Solid Angle, is that which is made by three or more plane angles, meeting each other in the same point.

94. Similar

94. Similar Solids, contained by plane figures, are such as have all their solid angles equal, each to each, and are bounded by the same number of similar planes, alike placed.

95. A Prism, is a solid whose ends are parallel, equal, and like plane figures, and its sides, connecting those ends, are parallelograms.

96. A prism takes particular names according to the figure of its base or ends, whether triangular, square, rectangular, pentagonal, hexagonal, &c.

97. A Right or Upright Prism, is that which has the planes of the sides perpendicular to the planes of the ends or base.

98. A Parallelopiped, or Parallelopipedon is a prism bounded by six parallelograms, every opposite two of which are equal, alike, and parallel.



99. A Rectangular Parallelopipedon, is that whose bounding planes are all rectangles, which are perpendicular to each other.

100. A Cube, is a square prism, being bounded by six equal square sides or faces, and are perpendicular to each other.



101. A Cylinder, is a round prism, having circles for its ends; and is conceived to be formed by the rotation of a right line about the circumferences of two equal and parallel circles, always parallel to the axis.



102. The Axis of a Cylinder, is the right line joining the centres of the two parallel circles, about which the figure is described.

103. A Pyramid, is a solid, whose base is any right-lined plane figure, and its sides triangles, having all their vertices meeting together in a point above the base, called the Vertex of the pyramid.



104. A pyramid, like the prism, takes particular names from the figure of the base.

105. A Cone, is a round pyramid, having a circular base, and is conceived to be generated by the rotation of a right line about the circumference of a circle, one end of which is fixed at a point above the plane of that circle.



106. The

106. The **Axis** of a cone, is the right line, joining the vertex, or fixed point, and the centre of the circle about which the figure is described.

107. **Similar Cones and Cylinders**, are such as have their altitudes and the diameters of their bases proportional.

108. A **Sphere**, is a solid bounded by one curve surface, which is every where equally distant from a certain point within called the **Centre**. It is conceived to be generated by the rotation of a semicircle about its diameter, which remains fixed.

109. The **Axis** of a Sphere, is the right line about which the semicircle revolves; and the centre is the same as that of the revolving semicircle.

110. The **Diameter** of a Sphere, is any right line passing through the centre, and terminated both ways by the surface.


111. The **Altitude** of a Solid, is the perpendicular drawn from the vertex to the opposite side or base.

THEOREM XCV.

A Perpendicular is the Shortest Line which can be drawn from any Point to a Plane.

LET AB be perpendicular to the plane DE ; then any other line, as AC , drawn from the same point A to the plane, will be longer than the line AB .

In the plane draw the line BC , joining the points B, C .



Then, because the line AB is perpendicular to the plane DE , the angle B is a right angle (def. 89), and consequently greater than the angle C ; therefore the line AB , opposite to the less angle, is less than any other line AC , opposite the greater angle (th. 21). Q. E. D.

THEOREM XCVI.

A Perpendicular Measures the Distance of any Point from a Plane.

THE distance of one point from another is measured by a right line joining them, because this is the shortest line which can be drawn from one point to another. So, also, the distance from a point to a line, is measured by a perpendicular, because this line is the shortest which can be drawn from


from the point to the line. In like manner, the distance from a point to a plane, must be measured by a perpendicular drawn from that point to the plane, because this is the shortest line which can be drawn from the point to the plane.

THEOREM XCVII.

The common Section of Two Planes, is a Right Line.

LET $ACBDA$, $AEBFA$, be two planes cutting each other, and A , B , two points in which the two planes meet : drawing the line AB , this line will be the common intersection of the two planes.

For because the right line AB touches the two planes in the points A and B , it touches them in all other points (def. 20) : this line is therefore common to the two planes. That is, the common intersection of the two planes is a right line. Q. E. D.




THEOREM XCVIII.

If a Line be Perpendicular to two other Lines, at their Common Point of Meeting ; it will be Perpendicular to the Plane of those Lines.

LET the line AB make right angles with the lines AC , AD ; then will it be perpendicular to the plane CDE which passes through these lines.

If the line AB were not perpendicular to the plane CDE , another plane might pass through the point A , to which the line AB would be perpendicular. But this is impossible ; for, since the angles BAC , BAD , are right angles, this other plane must pass through the points C , D . Hence, this plane passing through the two points A , C , of the line AC , and through the two points A , D , of the line AD , it will pass through both these two lines, and therefore be the same plane with the former.

Q. E. D.




THEOREM

THEOREM XCIX.

If Two Lines be Perpendicular to the Same Plane, they will be Parallel to each other.

LET the two lines AB , CD , be both perpendicular to the same plane $EBDF$; then will AB be parallel to CD .

FOR, join B , D , by the line BD in the plane. Then, because the lines AB , CD , are perpendicular to the plane EF , they are both perpendicular to the line BD (def. 89) in that plane: and consequently they are parallel to each other (corol. th. 13). Q. E. D.*




Corol. If two lines be parallel, and if one of them be perpendicular to any plane, the other will also be perpendicular to the same plane.

THEOREM C.

If Two planes Cut each other at Right Angles, and a Line be drawn in one of the Planes Perpendicular to their Common Intersection, it will be Perpendicular to the other Plane.

LET the two planes $ACBD$, $AEBF$, cut each other at right angles; and the line CG be perpendicular to their common section AB ; then will CG be also perpendicular to the other plane $AEBF$.

FOR, draw EG perpendicular to AB . Then because the two lines GC , GE , are perpendicular to the common intersection AB , the angle CGE is the angle of inclination of the two planes (def. 91). But since the two planes cut each other perpendicularly, the angle of inclination CGE , is a right angle. And since the line CG is perpendicular to the two lines GA , GE , in the plane $AEBF$, it is therefore perpendicular to that plane (th. 98). Q. E. D.



* This demonstration of Theorem XCIX. does not appear to me to be conclusive. EDITOR.

THEOREM CI.

If one Plane Meet another Plane, it will make Angles with that other Plane, which are together equal to two Right Angles.

LET the plane $ACBC$ meet the plane $AEBF$; these planes make with each other two angles whose sum is equal to two right angles.

For, through any point G , in the common section AB , draw CD , EF , perpendicular to AB . Then the line CG makes with EF two angles together equal to two right angles. But these two angles are (by def. 91) the angles of inclination of the two planes. Therefore the two planes make angles with each other, which are together equal to two right angles.

Corol. In like manner it may be demonstrated, that planes which intersect, have their vertical or opposite angles equal; also, that parallel planes have their alternate angles equal; and so on, as in parallel lines.

THEOREM CII.


If Two Planes be Parallel to each other; a Line which is Perpendicular to one of the planes, will also be Perpendicular to the other.

LET the two planes CD , EF , be parallel, and let the line AB be perpendicular to the plane CD ; then shall it also be perpendicular to the other plane EF .

For, from any point G in the plane EF , draw GH perpendicular to the plane CD , and draw AH , BG .

Then, because BA , GH , are both perpendicular to the plane CD , the angles A and H are both right angles. And because the planes CD , EF , are parallel, the perpendiculars BA , GH , are equal (def. 92). Hence it follows that the lines BG , AH , are parallel (def. 9). And the line AB being perpendicular to the line AH , is also perpendicular to the parallel line BG (cor. th. 12).

In like manner it is proved, that the line AB is perpendicular to all other lines which can be drawn from the point B in



in the plane EF . Therefore the line AB is perpendicular to the whole plane EF (def. 92). Q. E. D.


THEOREM CIII.

If Two Lines be Parallel to a Third Line, though not in the same Plane with it; they will be Parallel to each other.

LET the lines AB , CD , be each of them parallel to the third line EF , though not in the same plane with it; then will AB be parallel to CD .

For, from any point G in the line EF , let GH , GI , be each perpendicular to EF , in the planes EB , ED , of the proposed parallels.

Then, since the line EF is perpendicular to the two lines GH , GI , it is perpendicular to the plane GHI of those lines (th. 98). And because EF is perpendicular to the plane GHI , its parallel AB is also perpendicular to that plane (cor. th. 99). For the same reason, the line CD is perpendicular to the same plane GHI . Hence, because the two lines AB , CD , are perpendicular to the same plane, these two lines are parallel (th. 99). Q. E. D.




THEOREM CIV.

If Two Lines, that meet each other, be Parallel to Two other Lines that meet each other, though not in the same Plane with them; the Angles contained by those Lines will be equal.

LET the two lines AB , BC , be parallel to the two lines DE , EF ; then will the angle ABC be equal to the angle DEF .

For, make the lines AD , BE , CF , all equal to each other, and join AC , DF , AD , BE , CF .

Then, the lines AD , BE , joining the equal and parallel lines AB , DE , are equal and parallel (th. 24). For the same reason, CF , BE , are equal and parallel. Therefore AD , CF , are equal and parallel (th. 15); and consequently also AC , DF (th. 24). Hence, the two triangles ABC , DEF , having all their sides equal, each




each to each, have their angles also equal, and consequently the angle $\angle ABC = \text{the angle } \angle DEF$. Q. E. D.

THEOREM CV.

The Sections made by a Plane cutting two other Parallel Planes, are also Parallel to each other.

LET the two parallel planes AB, CD , be cut by the third plane $EFHG$, in the lines EF, GH : these two sections EF, GH , will be parallel.




Suppose EC, FH , be drawn parallel to each other in the plane $EFHG$; also let EI, FK , be perpendicular to the plane CD ; and let IG, KH , be joined.

Then EG, FH , being parallels, and EI, FK , being both perpendicular to the plane CD , are also parallel to each other (th. 99); consequently the angle $\angle HFK$ is equal to the angle $\angle GEI$ (th. 104). But the angle $\angle FKH$ is also equal to the angle $\angle IGE$, being both right angles; therefore the two triangles are equiangular (cor. 1, th. 17;) and the sides FK, EI , being the equal distances between the parallel planes (def. 92), it follows that the sides FH, EG , are also equal (th. 2). But these two lines are parallel (by suppos.), as well as equal; consequently the two lines EF, GH , joining those equal parallels, are also parallel (th. 24). Q. E. D.

THEOREM CVI.


If any Prism be cut by a Plane Parallel to its Base, the Section will be equal and Like to the Base.

LET AG be any prism. and IL a plane parallel to the base AC ; then will the plane IL be equal and like to the base AC , or the two planes will have all their sides and all their angles equal.



For the two planes AC, IL , being parallel, by hypothesis; and two parallel planes, cut by a third plane. having parallel sections (th. 105); therefore IK is parallel to AB , and KL to BC , and LM to CD , and IM to AD . But AI and BK are parallels (by def. 95). consequently AK is a parallelogram; and the opposite sides AB, IK , are equal (th. 22). In like manner,

manner, it is shown that KL is $=BC$, and $LM = CD$, and $IM = AD$, or the two planes AC, IL , are mutually equilateral. But these two planes, having their corresponding sides parallel, have the angles contained by them also equal (th. 104), namely, the angle $A =$ the angle I , the angle $B =$ the angle K , the angle $C =$ the angle L , and the angle $D =$ the angle M . So that the two planes AC, IL , have all their corresponding sides and angles equal, or they are equal and like. Q. E. D.




THEOREM CVII.

If a Cylinder be cut by a Plane Parallel to its Base, the Section will be a Circle, Equal to the Base.

LET AF be a cylinder, and GHI any section parallel to the base ABC ; then will GHI , be a circle equal to ABC .

For, let the planes KE, KF , pass through the axis of the cylinder MK , and meet the section GHI in the three points H, I, L ; and join the points as in the figure.



Then, since KL, CI , are parallel (by def. 101); and the plane KI , meeting the two parallel planes ABC, GHI , makes the two sections KC, LI , parallel (th. 105); the figure $KLIC$ is therefore a parallelogram, and consequently has the opposite sides LI, KC , equal, where KC is a radius of the circular base.


In like manner it is shown that LH is equal to the radius KB ; and that any other lines, drawn from the point L to the circumference of the section GHI , are all equal to radii of the base; consequently GHI is a circle, and equal to ABC . Q. E. D.

THEOREM CVIII.

All Prisms and Cylinders, of Equal Bases and Altitudes, are Equal to each other.

LET AC, DF , be two prisms, and a cylinder, on equal bases AB, DE , and having equal altitudes EC, FF ; then will the solids AC, DF , be equal.

For, let pq, rs , be any



any two sections parallel to the bases, and equidistant from them. Then, by the last two theorems, the section PQ is equal to the base AB , and the section RS equal to the base DE . But the bases AB , DE , are equal, by the hypothesis; therefore the sections PQ , RS , are equal also. In like manner, it may be shown, that any other corresponding sections are equal to one another.

Since then every section in the prism AC , is equal to its corresponding section in the prism or cylinder DE , the prisms and cylinder themselves, which are composed of an equal number or all those equal sections, must also be equal. Q. E. D.

Corol. Every prism, or cylinder, is equal to a rectangular parallelepipedon, of an equal base and altitude.

THEOREM CIX.

Rectangular Parallelepipedons, of Equal Altitudes, are to each other as their Bases.


LET AC , EG , be two rectangular parallelepipedons, having the equal altitudes AD , EH ; then will the solid AC be to the solid EG as the base AB is to the base EF .

For, let the proportion of the base AB to the base EF , be that of any one number m (3)

to any other number n (2). And conceive AB to be divided into m equal parts, or rectangles, AI , LK , MB , (by dividing AN into that number of equal parts, and drawing IL , KM , parallel to BN). And let EF be divided, in like manner, into n equal parts, or rectangles, EO , FP : all of these parts of both bases being mutually equal among themselves. And through the lines of division let the plane sections LR , MS , PV , pass parallel to AQ , ET .

Then, the parallelepipedons AR , LS , MC , EV , PG , are all equal, having equal bases and altitudes. Therefore the solid AC is to the solid EG , as the number of parts in the former, to the number of equal parts in the latter; or as the number of parts in AB to the number of equal parts in EF , that is, as the base AB to the base EF . Q. E. D.

Corol. From this theorem, and the corollary to the last, it appears, that all prisms and cylinders of equal altitudes, are



to each other as their bases ; every prism and cylinder being equal to a rectangular parallelopipedon of an equal base and altitude.

THEOREM CX.

Rectangular Parallelopipedons, of Equal Bases, are to each other as their Altitudes.

LET AB , CD , be two rectangular parallelopipedons, standing on the equal bases AE , CF ; then will the solid AB be to the solid CD , as the altitude EB is to the altitude FD .

For, let AG be a rectangular parallelopipedon on the base AE , and its altitude EG equal to the altitude FD of the solid CD .

Then AG and CD are equal, being prisms of equal bases and altitudes. But if HB , HG , be considered as bases, the solids AB , AG , of equal altitude AH , will be to each other as those bases HB , HG . But these bases HB , HG , being parallelograms of equal altitude HE , are to each other as their bases EB , EG ; therefore the two prisms AB , AG , are to each other as the lines EB , EG . But AG is equal to CD , and EG equal to FD ; consequently the prisms AG , CD , are to each other as their altitudes EB , FD ; that is, $AB : CD :: EB : FD$. Q. E. D.

Corol. 1. From this theorem, and the corollary to theorem 108, it appears, that all prisms and cylinders, of equal bases, are to one another as their altitudes.

Corol. 2. Because, by corollary 1, prisms and cylinders are as their altitudes, when their bases are equal. And, by the corollary to the last theorem, they are as their bases, when their altitudes are equal. Therefore, universally, when neither are equal, they are to one another as the product of their bases and altitudes. And hence also these products are the proper numeral measures of their quantities or magnitudes.


THEOREM CXI.

Similar Prisms and Cylinders are to each other, as the Cubes of their Altitudes, or of any other Like Linear Dimensions.

LET $ABGD$, $EFGH$, be two similar prisms ; then will the prism CD be to the prism GH , as AB^3 to EF^3 OR AD^3 to EH^3 .

For

For the solids are to each other as the product of their bases and altitudes (th. 110, cor. 2), that is, as $AC \cdot AD$ to $EG \cdot EH$. But the bases, being similar planes, are to each other as the squares of their like sides, that is, AC to EG as AB^2 to EF^2 , therefore the solid CD is to the solid GH , as $AB^2 \cdot AD$ to $EF^2 \cdot EH$. But BD and FH , being similar planes, have their like sides proportional, that is, $AB : EF :: AD : EH$, - - - or $AB^2 : EF^2 :: AD^2 : EH^2$; therefore $AB^2 \cdot AD : EF^2 \cdot EH :: AB^3 : EF^3$, or $CD : GH :: AD^3 : EH^3$; conseq. the solid $CD : \text{solid } GH :: AB^3 : EF^3 :: AD^3 : EH^3$. Q. E. D.




THEOREM CXII.

In any Pyramid, a Section Parallel to the Base is similar to the Base; and these two planes are to each other as the Squares of their Distances from the Vertex.


LET $ABCD$ be a pyramid, and EFG a section parallel to the base BCD , also AIH a line perpendicular to the two planes at H and I : then will BD, EG , be two similar planes, and the plane BD will be to the plane EG , as AH^2 to AI^2 .

For, join CH, FI . Then, because a plane cutting two parallel planes, makes parallel sections (th. 105), therefore the plane ABC , meeting the two parallel planes BD, EG , makes the sections BC, EF , parallel: In like manner, the plane ACD makes the sections CD, FG , parallel. Again, because two pair of parallel lines make equal angles (th. 104), the two EF, FG , which are parallel to BC, CD , make the angle EFG equal the angle BCD . And in like manner it is shown, that each angle in the plane EG is equal to each angle in the plane BD , and consequently those two planes are equiangular.

Again, the three lines AB, AC, AD , making with the parallels BC, EF , and CD, FG , equal angles (th. 14), and the angles at A being common, the two triangles ABC, AEF , are equiangular, as also the two triangles ACD, AFG , and have therefore their like sides proportional, namely, - - -



$AC : AF :: BC : EF :: CD : FG$. And in like manner it may be shown, that all the lines in the plane EG , are proportional to all the corresponding lines in the base BD . Hence these two planes, having their angles equal, and their sides proportional; are similar, by def. 63.




But, similar planes being to each other as the squares of their like sides, the plane $BD : EG :: BC^2 : EF^2$, or $AC^2 : AF^2$, by what is shown above. Also, the two triangles AHC , AIF , having the angles H and I right ones (th. 98), and the angle A common, are equiangular, and have therefore their like sides proportional, namely, $AC : AF :: AH : AI$, or $AC^2 : AF^2 :: AH^2 : AI^2$. Consequently the two planes BD , EG , which are as the former squares AC^2 , AF^2 , will be also as the latter squares AH^2 , AI^2 , that is, $BD : EG :: AH^2 : AI^2$. Q. E. D.

THEOREM CXIII.

If a Cone, any Section Parallel to the Base is a Circle; and this Section is to the Base, as the Squares of their Distances from the Vertex.

LET $ABCD$ be a cone, and GHI a section parallel to the base BCD ; then will GHI be a circle, and BCD , GHI , will be to each other, as the squares of their distances from the vertex.



For, draw ALF perpendicular to the two parallel planes; and let the planes ACE , ADE , pass through the axis of the cone AK , meeting the section in the three points H , I , K .

Then, since the section GHI is parallel to the base BCD , and the planes CK , DK , meet them, HK is parallel to CE , and IK to DE (th. 105). And because the triangles formed by these lines are equiangular, $KH : EC :: AK : AE :: KI : ED$. But EC is equal to ED , being radii of the same circle; therefore KI is also equal to KH . And the same may be shown of any other lines drawn from the point K to the perimeter of the section GHI , which is therefore a circle (def. 44).


Again, by similar triangles, $AL : AF :: AK : AE$, or $AL^2 : AF^2 :: AK^2 : AE^2$; hence $AL^2 : AF^2 :: KI^2 : ED^2$; but $KI^2 : AD^2 ::$ circle $GHI : circle BCD (th. 93); therefore $AL^2 : AF^2 ::$ circle $GHI : circle BCD . Q. E. D.$$

THEOREM

THEOREM CXIV.

All Pyramids, and Cones, of Equal Bases and Altitudes, are Equal to one another.

LET $ABC, DEF,$ be any pyramids and cone, of equal bases $BC, EF,$ and equal altitudes $AG, DH,$ then will the pyramids and cone ABC and $DEF,$ be equal.



For, parallel to the bases and at equal distances $AN, DO,$ from the vertices, suppose the planes $IK, LM,$ to be drawn.

Then, by the two preceding theorems, -----

$$DO^2 : DH^2 :: LM : EF, \text{ and}$$

$$AN^2 : AG^2 :: IK : BC.$$

But since $AN^2, AG^2,$ are equal to $DO^2, DH^2,$ therefore $IK : BC :: LM : EF.$ But BC is equal to $EF,$ by hypothesis; therefore IK is also equal to $LM.$

In like manner it is shown, that any other sections, at equal distance from the vertex, are equal to each other.


Since then, every section in the cone, is equal to the corresponding section in the pyramids, and the heights are equal, the solids $ABC, DEF,$ composed of all those sections, must be equal also. Q. E. D.

THEOREM CXV.

Every Pyramid is the Third Part of a Prism of the Same Base and Altitude.

LET $ABCDEF$ be a prism, and $BDEF$ a pyramid, on the same triangular base $DEF:$ then will the pyramid, $BDEF$ be a third part of the prism $ABCDEF.$

For, in the planes of the three sides of the prism, draw the diagonals $BF, BD, CD.$ Then the two planes $BDF, BCD,$ divide the whole prism into the three pyramids $BDEF, DABC, DCBF,$ which are proved to be all equal to one another, as follows.




Since the opposite ends of the prism are equal to each other, the pyramid whose base is ABC and vertex $B,$ is equal

to

to the pyramid whose base is DEF and vertex B (th. 114), being pyramids of equal base and altitude.

But the latter pyramid, whose base is DEF and vertex B , is the same solid as the pyramid whose base is BEF and vertex D , and this is equal to the third pyramid whose base is BCF and vertex D , being pyramids of the same altitude and equal bases BEF , BCF .



Consequently all the three pyramids, which compose the prism, are equal to each other, and each pyramid is the third part of the prism, or the prism is triple of the pyramid.

Q. E. D.

Hence also, every pyramid, whatever its figure may be, is the third part of a prism of the same base and altitude; since the base of the prism, whatever be its figure, may be divided into triangles, and the whole solid into triangular prisms and pyramids.

Corol. Any cone is the third part of a cylinder, or of a prism, of equal base and altitude; since it has been proved that a cylinder is equal to a prism, and a cone equal to a pyramid, of equal base and altitude.


Scholium. Whatever has been demonstrated of the proportionality of prisms, or cylinders, holds equally true of pyramids, or cones; the former being always triple the latter; viz. that similar pyramids or cones are as the cubes of their like linear sides, or diameters, or altitudes, &c. And the same for all similar solids whatever, viz. that they are in proportion to each other, as the cubes of their like linear dimensions, since they are composed of pyramids every way similar.

THEOREM CXVI.

If a Sphere be cut by a Plane, the Section will be a Circle.

LET the sphere $AEBF$ be cut by the plane ADB ; then will the section ADB be a circle.

Draw the chord AB , or diameter of the section; perpendicular to which, or to the section ADB , draw the axis of the sphere $ECGF$, through the centre C , which will bisect the chord AB in the point G (th. 41). Also, join CA , CB ;



and

and draw CD , GD , to any point D in the perimeter of the section ADB .

Then, because CG is perpendicular to the plane ADB , it is perpendicular both to GA and GD (def. 90). So that CGA , CGD are two right-angled triangles, having the perpendicular CG common, and the two hypotenuses CA , CD , equal, being both radii of the sphere; therefore the third sides GA , GD , are also equal (cor. 2, th. 34). In like manner it is shown, that any other line, drawn from the centre G to the circumference of the section ADB , is equal to GA or GB ; consequently that section is a circle.


Corol. The section through the centre, is a circle having the same centre and diameter as the sphere, and is called a great circle of the sphere; the other plane sections being little circles.

THEOREM CXVII.

Every Sphere is Two-Thirds of its Circumscribing Cylinder.

LET $ABCD$ be a cylinder, circumscribing the sphere $EFGH$; then will the sphere $EFGH$ be two thirds of the cylinder $ABCD$.


For, let the plane AC be a section of the sphere and cylinder through the centre I . Join AI , BI . Also, let FIH be parallel to AD or BC , and EIG and KL parallel to AB or DC , the base of the cylinder; the latter line KL meeting BI in M , and the circular section of the sphere in N .



Then if the whole plane $HFBC$ be conceived to revolve about the line HF as an axis, the square FG will describe a cylinder AG , and the quadrant IFG will describe a hemisphere EFG , and the triangle IFB will describe a cone IAB . Also, in the rotation, the three lines or parts KL , KN , KM , as radii, will describe corresponding circular sections of those solids, namely, KL a section of the cylinder, KN a section of the sphere, and KM a section of the cone.

Now, FB being equal to FI or IG , and KL parallel to FB , then by similar triangles IK is equal to HM (th. 82). And since, in the right-angled triangle IKN , IN^2 is equal to $IK^2 + KN^2$ (th. 34); and because KL is equal to the radius IC
or

or IN , and $KM=IK$, therefore KL^2 is equal to KM^2+KN^2 , or the square of the longest radius, of the said circular sections, is equal to the sum of the squares of the two others. And because circles are to each other as the squares of their diameters, or of their radii, therefore the circle described by KL is equal to both the circles described by KM and KN ; or the section of the cylinder, is equal to both the corresponding sections of the sphere and cone. And as this is always the case in every parallel position of KL , it follows, that the cylinder EB , which is composed of all the former sections, is equal to the hemisphere EFB and cone IFB , which are composed of all the latter sections.



But the cone IAB is a third part of the cylinder EB (cor. 2, th. 115); consequently the hemisphere EFB is equal to the remaining two-thirds; or the whole sphere $EFGB$ equal to two-thirds of the whole cylinder $ABCD$. Q. E. D.

Corol. 1. A cone, hemisphere, and cylinder of the same base and altitude, are to each other as the numbers 1, 2, 3.

Corol. 2. All spheres are to each other as the cubes of their diameters; all these being like parts of their circumscribing cylinders.


Corol. 3. From the foregoing demonstration it also appears, that the spherical zone or frustrum $EGNP$, is equal to the difference between the cylinder $EGLO$, and the cone IMQ , all of the same common height IK . And that the spherical segment PFN , is equal to the difference between the cylinder $ABLO$ and the conic frustrum $AQMB$, all of the same common altitude FK .

PROBLEMS.

PROBLEM I.

To Bisect a Line AB ; that is, to divide it into two Equal Parts.

FROM the two centres A and B , with any equal radii, describe arcs of circles, intersecting each other in C and D ; and draw the line CD , which will bisect the given line AB in the point E .




For, draw the radii AC , BC , AD , BD . Then, because all these four radii are equal, and the side CD common, the two triangles ACD , BCD , are mutually equilateral : consequently they are also mutually equiangular (th. 5), and have the angle ACE equal to the angle BCE .

Hence, the two triangles ACE , BCE , having the two sides AC , CE , equal to the two sides BC , CE , and their contained angles equal, are identical (th. 1), and therefore have the side AE equal to EB . Q. E. D.

PROBLEM II.

To Bisect an Angle BAC .

FROM the centre A , with any radius, describe an arc, cutting off the equal lines AD , AE ; and from the two centres D , E , with the same radius, describe arcs intersecting in F ; then draw AF , which will bisect the angle A as required.



For, join DF , EF . Then the two triangles ADF , AEF , having the two sides AD , DF , equal to the two AE , EF (being equal radii), and the side AF common, they are mutually equilateral ; consequently they are also mutually equiangular (th. 5), and have the angle BAF equal to the angle CAF .

Scholium. In the same manner is an arc of a circle bisected.

PROBLEM III.

At a Given Point c , in a Line AB , to Erect a Perpendicular.

From the given point c , with any radius, cut off any equal parts CD , CE , of the given line; and, from the two centres D and E , with any one radius, describe arcs intersecting in F ; then join CF , which will be perpendicular as required.




For, draw the two equal radii DF , EF . Then the two triangles CDF , CEF , having the two sides CD , DF , equal to the two CE , EF , and CF common, are mutually equilateral; consequently they are also mutually equiangular (th. 5), and have the two adjacent angles at c equal to each other; therefore the line CF is perpendicular to AB (def. 11).

Otherwise.

When the Given Point c is near the End of the line.

From any point D , assumed above the line, as a centre, through the given point c describe a circle, cutting the given line at E ; and through E and the centre D , draw the diameter EDF ; then join CF , which will be the perpendicular required.




For the angle at c , being an angle in a semicircle, is a right angle, and therefore the line CF is a perpendicular (by def. 15).

PROBLEM IV.

From a Given point A to let fall a Perpendicular on a given Line BC .

From the given point A as a centre, with any convenient radius, describe an arc, cutting the given line at the two points D and E ; and from the two centres D , E , with any radius, describe two arcs, intersecting at F ; then draw AGF , which will be perpendicular to BC as required.




For, draw the equal radii AD , AE , and DF , EF . Then the two triangles ADF , AEF , having the two sides AD , DF , equal to the two AE , EF , and AF common, are mutually

ally equilateral ; consequently they are also mutually equiangular (th. 5), and have the angle DAG equal the angle FAG . Hence then, the two triangles ADG , AEG , having the two sides AD , AG , equal to the two AE , AG , and their included angles equal, are therefore equiangular (th. 1), and have the angles at G equal ; consequently AG is perpendicular to BC (def. 11).

Otherwise.

When the Given Point is nearly Opposite the end of the Line.

FROM any point D , in the given line BC , as a centre, describe the arc of a circle through the given point A , cutting BC in E ; and from the centre E , with the radius EA , describe another arc, cutting the former in F ; then draw AGF , which will be perpendicular to BC as required.




For, draw the equal radii DA , DF , and EA , EF . Then the two triangles DAE , DFE , will be mutually equilateral ; consequently they are also mutually equiangular (th. 5), and have the angles at D equal. Hence, the two triangles DAG , DFG , having the two sides DA , DG , equal to the two DF , DG , and the included angles at D equal, have also the angles at G equal (th. 1) ; consequently those angles at G are right angles, and the line AG is perpendicular to BC .

PROBLEM V.

At a Given Point A , in a Line AB , to make an Angle Equal to a Given Angle c .

FROM the centres A and C with any one radius, describe the arcs DE , FG . Then, with radius DE , and centre F , describe an arc cutting FG in G . Through G draw the line AG , and it will form the angle required.




For, conceive the equal lines or radii, DE , FG , to be drawn. Then the two triangles CDE , AFG , being mutually equilateral, are mutually equiangular (th. 5), and have the angle at A equal to the angle c .

PROBLEM

PROBLEM VI.

Through a Given Point *A*, to draw a Line Parallel to a Given Line *BC*.

From the given point *A* draw a line *AD* to any point in the given line *BC*. Then draw the line *EAF* making the angle at *A* equal to the angle at *D* (by prob. 5); so shall *EF* be parallel to *BC* as required.




For, the angle *D* being equal to the alternate angle *A*, the lines *BC*, *EF*, are parallel, by th. 13.

PROBLEM VII.

To Divide a Line *AB* into any proposed Number of Equal Parts.

Draw any other line *AC*, forming any angle with the given line *AB*; on which set off as many of any equal parts, *AD*, *DE*, *EF*, *FC*, as the line *AB* is to be divided into. Join *BC*; parallel to which draw the other lines *FG*, *EH*, *DI*: then these will divide




AB in the manner as required.—For those parallel lines divide both the sides *AB*, *AC*, proportionally, by th. 82.

PROBLEM VIII.

To find a Third Proportional to Two given Lines *AB*, *AC*.

PLACE the two given lines *AB*, *AC*, forming any angle at *A*; and in *AB* take also *AD* equal to *AC*. Join *BC*, and draw *DE* parallel to it; so will *AE* be the third proportional sought.



For, because of the parallels *BC*, *DE*, the two lines *AB*, *AC*, are cut proportionally (th. 82); so that $AB : AC :: AD$ or $AC : AE$; therefore *AE* is the third proportional to *AB*, *AC*.


PROBLEM IX.

To find a Fourth Proportional to three Lines *AB*, *AC*, *AD*.

PLACE two of the given lines *AB*, *AC*, making any angle at *A*; also place *AD* on *AB*. Join *BC*; and parallel to it draw *DE* :

so shall AE be the fourth proportional as required.


For, because of the parallels BC, DE , the two sides AB, AC , are cut proportionally (th. 82); so that
 $AB : AC :: AD : AE$.



PROBLEM X.

To find a Mean Proportional between Two Lines AB, BC .


PLACE AB, BC , joined in one straight line AC : on which as a diameter, describe the semicircle ADC ; to meet which erect the perpendicular BD : and it will be the mean proportional sought, between AB and BC (by cor. th. 87).



PROBLEM XI.

To find the Centre of a Circle.


DRAW any chord AB ; and bisect it perpendicularly with the line CD , which will be a diameter (th. 41, cor.). Therefore, CD bisected into o , will give the centre, as required.



PROBLEM XII.

To describe the Circumference of a Circle through Three Given Points A, B, c .

FROM the middle point B draw chords BA, BC , to the two other points, and bisect these chords perpendicularly by lines meeting in o , which will be the centre. Then from the centre o , at the distance of any one of the points, as OA , describe a circle, and it will pass through the two other points B, c , as required.



For, the two right-angled triangles OAD, OBD , having the sides AD, DB , equal (by constr.), and OD common with the included right angles at D equal, have their third sides OA, OB , also equal (th. 1). And in like manner it is shown, that oc is equal to OB or OA . So that all the three OA, OB, oc , being equal, will be radii of the same circle.

PROBLEM



PROBLEM XIII.

To draw a Tangent to a Circle, through a Given Point *A*.

WHEN the given point *A* is in the circumference of the circle : Join *A* and the centre *o* ; perpendicular to which draw *BAC*, and it will be the tangent, by th. 46.

But when the given point *A* is out of the circle, draw *AO* to the centre *o* ; on which as a diameter describe a semi-circle, cutting the given circumference in *B* ; through which draw *BADc*, which will be the tangent as required.

For, join *DO*. Then the angle *ADO*, in a semicircle, is a right angle, and consequently *AD* is perpendicular to the radius, *DO*, or is a tangent to the circle (th. 46).




PROBLEM XIV.

On a Given Line *B* to describe a Segment of a Circle, to Contain a Given Angle *c*.

At the ends of the given line make angles *DAB*, *DBA*, each equal to the given angle *c*. Then draw *AE*, *BE*, perpendicular to *AD*, *BD* ; and with the centre *E*, and radius *EA* or *EB*, describe a circle ; so shall *AFB* be the segment required, as an angle *F* made in it will be equal to the given angle *c*.


For, the two lines *AD*, *BD*, being perpendicular to the radii *EA*, *EB* (by constr.), are tangents to the circle (th. 46) ; and the angle *A* or *B*, which is equal to the given angle *c* by construction, is equal to the angle *F* in the alternate segment *AFB* (th. 53).



PROBLEM XV.

To Cut off a Segment from a Circle, that shall Contain a Given Angle *c*.

DRAW any tangent *AB* to the given circle ; and a chord *AD* to make the angle *DAB* equal to the given angle *c* ; then *DEA* will be the segment required, an angle *E* made in it being equal to the given angle *c*.



For


For the angle A , made by the tangent and chord, which is equal to the given angle c by construction, is also equal to any angle E in the alternate segment (th. 53).

PROBLEM XVI.

To make an Equilateral Triangle on a Given Line AB .

FROM the centres A and B , with the distance AB , describe arcs, intersecting in C . Draw AC , BC , and ABC will be the equilateral triangle.

For the equal radii AC , BC , are, each of them, equal to AB .




PROBLEM XVII.

To make a Triangle with Three Given Lines AB , AC , BC .

WITH the centre A , and distance AC , describe an arc. With the centre B , and distance BC , describe another arc, cutting the former in C . Draw AC , BC , and ABC will be the triangle required.

For the radii, or sides of the triangle, AC , BC , are equal to the given lines AC , BC , by construction.




PROBLEM XVIII.

To make a Square on a Given Line AB .

RAISE AD , BC , each perpendicular and equal to AB ; and join DC ; so shall $ABCD$ be the square sought.

For all the three sides AB , AD , BC , are equal, by the construction, and DC is equal and parallel to AB (by th. 24); so that all the four sides are equal, and the opposite ones are parallel. Again, the angle A or B , of the parallelogram, being a right angle, the angles are all right ones (cor. 1, th. 22). Hence, then, the figure, having all its sides equal, and all its angles right, is a square (def. 34).



PROBLEM


PROBLEM XIX.

To make a Rectangle, or a Parallelogram, of a Given Length and Breadth, AB , BC .

ERECT AD , BC , perpendicular to AB , and each equal to BC ; then join DC , and it is done.

The demonstration is the same as the last problem.

And in the same manner is described any oblique parallelogram, only drawing AD and BC to make the given oblique angle with AB , instead of perpendicular to it.




PROBLEM XX.

To Inscribe a Circle in a Given Triangle ABC .

BISECT any two angles A and B , with the two lines AD , BD . From the intersection D , which will be the centre of the circle, draw the perpendiculars DE , DF , DG , and they will be the radii of the circle required.

For, since the angle DAE is equal to the angle DAG , and the angles at E , G , right angles (by constr.), the two triangles ADE , ADG , are equiangular; and, having also the side AD common, they are identical, and have the sides DE , DG , equal (th. 2). In like manner it is shown, that DF is equal to DE or DG .

Therefore, if with the centre D , and distance DE , a circle be described, it will pass through all the three points E , F , G , in which points also it will touch the three sides of the triangle (th. 46), because the radii DE , DF , DG , are perpendicular to them.




PROBLEM XXI.

To Describe a Circle about a Given Triangle ABC .

BISECT any two sides with two of the perpendiculars DE , DF , DG , and D will be the centre.

For, join DA , DB , DC . Then the two right-angled triangles DAE , DBE , have the two sides DE , EA , equal to the two DE , EB , and the included angles at E equal: those two triangles are therefore identical




(th.

(th. 1), and have the side DA equal to DB . In like manner it is shown, that DC is also equal to DA or DB . So that all the three DA , DB , DC , being equal, they are radii of a circle passing through A , B , and C .

PROBLEM XXII.

To Inscribe an Equilateral Triangle in a Given Circle.

THROUGH the centre c draw any diameter AB . From the point B as a centre, with the radius BC of the given circle describe an arc DCE . Join AD , AE , DE , and ADE is the equilateral triangle sought.




For, join DB , DC , EB , EC . Then DCE is an equilateral triangle, having each side equal to the radius of the given circle.

In like manner, BCE is an equilateral triangle. But the angle ADE is equal to the angle ABE or CBE , standing on the same arc AE ; also the angle AED is equal to the angle CBD , on the same arc AD ; hence the triangle DAE has two of its angles, ADE , AED , equal to the angles of an equilateral triangle, and therefore the third angle at A is also equal to the same; so that triangle is equiangular, and therefore equilateral.

PROBLEM XXIII.

To Inscribe a Square in a Given Circle.

DRAW two diameters AC , BD , crossing at right angles in the centre E . Then join the four extremities A , B , C , D , with right lines, and these will form the inscribed square $ABCD$.




For, the four right-angled triangles AEB , BEC , CED , DEA , are identical, because they have the sides EA , EB , EC , ED , all equal, being radii of the circle, and the four included angles at E all equal, being right angles, by the construction. Therefore all their third sides AB , BC , CD , DA , are equal to one another, and the figure $ABCD$ is equilateral. Also, all its four angles, A , B , C , D , are right ones, being angles in a semicircle. Consequently the figure is a square.

PROBLEM XXIV.

To Describe a Square about a Given Circle.

DRAW two diameters AC , BD , crossing at right angles in the centre E . Then through their four extremities draw FG , IH , parallel to AC , and FI , GH , parallel to BD , and they will form the square $FGHI$.



For, the opposite sides of parallelograms being equal, FG and IH are each equal to the diameter AC , and FI and GH each equal to the diameter BD ; so that the figure is equilateral. Again, because the opposite angles of parallelograms are equal, all the four angles F , G , H , I , are right angles, being equal to the opposite angles at E . So that the figure $FGHI$, having its sides equal, and its angles right ones, is a square, and its sides touch the circle at the four points A , B , C , D , being perpendicular to the radii drawn to those points.

PROBLEM XXV.

To Inscribe a Circle in a Given Square.

BISECT the two sides FG , FI , in the points A and B (last fig.). Then through these two points draw AC parallel to FG or IH , and BD parallel to FI or GH . Then the point of intersection E will be the centre, and the four lines EA , EB , EC , ED , radii of the inscribed circle.

For, because the four parallelograms EF , EG , EH , EI , have their opposite sides and angles equal, therefore all the four lines EA , EB , EC , ED , are equal, being each equal to half a side of the square. So that a circle described from the centre E , with the distance EA , will pass through all the points A , B , C , D , and will be inscribed in the square, or will touch its four sides in those points, because the angles there are right ones.

PROBLEM XXVI.

To Describe a Circle about a Given Square,
(see fig. Prob. xxiii.)

DRAW the diagonals AC , BD , and their intersection E will be the centre.

For the diagonals of a square bisect each other (th. 40), making EA , EB , EC , ED , all equal, and consequently these are radii of a circle passing through the four points A , B , C , D .


PROBLEM

PROBLEM XXVII.

To Cut a Given Line in Extreme and Mean Ratio.

Let AB be the given line to be divided in extreme and mean ratio, that is, so that the whole line may be to the greater part, as the greater part is to the less part.

Draw BC perpendicular to AB , and equal to half AB . Join AC ; and with centre C and distance CB , describe the circle BD ; then with centre A and distance AD , describe the arc DE ; so shall AB be divided in E in extreme and mean ratio, or so that $AB : AE :: AE : EB$.



For, produce AC to the circumference at F . Then, ADF being a secant, and AB a tangent, because B is a right angle: therefore the rectangle $AF \cdot AD$ is equal to AB^2 (cor. 1, th. 61); consequently the means and extremes of these are proportional (th. 77). viz. $AB : AF$ or $AD + DF :: AD : AB$. But AE is equal to AD by construction, and $AB = 2BC = DF$; therefore, $AB : AE + AB :: AE : AB$; and by division, $AB : AE :: AE : EB$.

PROBLEM XXVIII.

To Inscribe an Isosceles Triangle in a Given Circle, that shall have each of the Angles at the Base Double the Angle at the Vertex.

Draw any diameter AB of the given circle; and divide the radius CB , in the point D , in extreme and mean ratio, by the last problem. From the point B apply the chords BE, BF , each equal to the greater part CD . Then join AE, AF, EF ; and AEF will be the triangle required.



For, the chords BE, BF , being equal, their arcs are equal; therefore the supplemental arcs and chords AE, AF , are also equal; consequently the triangle AEF is isosceles, and has the angle E equal to the angle F ; also the angles at G are right angles.


Draw CF and DF . Then, $BC : CD :: CD : BD$, or $BC : BF :: BF : BD$ by constr. And $BA : BF :: BF : BG$ (by th. 87). But $BC = \frac{1}{2}BA$; therefore $BG = \frac{1}{2}BD = GD$; therefore the two triangles GBF, GDF , are identical (th. 1), and each equiangular to

to ABF and AGF (th. 87). Therefore their doubles, BFD , AFE , are isosceles and equiangular, as well as the triangle BCF ; having the two sides BC , CF , equal, and the angle B common with the triangle BFD . But CD is $= DF$ or BF ; therefore the angle $c =$ the angle DFC (th. 4); consequently the angle BDF , which is equal to the sum of these two equal angles (th. 16), is double of one of them c ; or the equal angle B or CFB double the angle c . So that CBF is an isosceles triangle, having each of its two equal angles double of the third angle c . Consequently the triangle AEF (which it has been shown is equiangular to the triangle CBF) has also each of its angles at the base double the angle A at the vertex.

PROBLEM XXIX.

To Inscribe a Regular Pentagon in a Given Circle.

INSCRIBE the isosceles triangle ABC having each of the angles ABC , ACB , double the angle BAC (prob. 23). Then bisect the two arcs ADB , AEC , in the points D , E ; and draw the chords AD , DB , AE , EC , so shall $ADBCE$ be the inscribed equilateral pentagon required.



For, because equal angles stand on equal arcs, and double angles on double arcs, also the angles ABC , ACB , being each double the angle BAC , therefore the arcs ADB , AEC , subtending the two former angles, each one double the arcs BC subtending the latter. And since the two former arcs are bisected in D and E , it follows that all the five arcs AD , DB , BC , CE , EA , are equal to each other, and consequently the chords also which subtend them, or the five sides of the pentagon, are all equal.


Note. In the construction, the points D and E are most easily found, by applying BD and CE each equal to BC .

PROBLEM XXX.

To Inscribe a Regular Hexagon in a Circle.

APPLY the radius AO of the given circle as a chord, AB , BC , CD , &c, quite round the circumference, and it will complete the regular hexagon $ABCDEF$.

For, draw the radii AO , BO , CO , DO , EO , FO , completing six equal triangles; of which any one, as ABO , being equilateral



(by


by constr.) its three angles are all equal (cor. 2, th. 3), and any one of them, as AOB , is one third of the whole, or of two right angles (th. 17), or one-sixth of four right angles. But the whole circumference is the measure of four right angles (cor. 4, th. 6). Therefore the arc AB is one-sixth of the circumference of the circle, and consequently its chord AB one side of an equilateral hexagon inscribed in the circle. And the same of the other chords.

Corol. The side of a regular hexagon is equal to the radius of the circumscribing circle, or to the chord of one-sixth part of the circumference.

PROBLEM XXXI.

To describe a Regular Pentagon or Hexagon about a Circle.

In the given circle inscribe a regular polygon of the same name or number of sides, as ABCDE , by one of the foregoing problems. Then to all its angular points draw tangents (by prob. 13) and these will form the circumscribing polygon required.



For, all the chords, or sides of the inscribing figure, AB , BC , &c. being equal, and all the radii OA , OB , &c. being equal, all the vertical angles about the point O are equal. But the angles OEF , OAF , OAG , OBG , made by the tangents and radii, are right angles; therefore $\text{OEF} + \text{OAF} = \text{two right angles}$, and $\text{OAG} + \text{OBG} = \text{two right angles}$; consequently, also, $\text{AOE} + \text{AFE} = \text{two right angles}$, and $\text{AOB} + \text{AGB} = \text{two right angles}$ (cor. 2, th. 18). Hence, then, the angles $\text{AOE} + \text{AFE}$ being $= \text{AOB} + \text{AGB}$, of which AOB is $= \text{AOE}$; consequently the remaining angles F and G are also equal. In the same manner it is shown, that all the angles F , G , H , I , K , are equal.

Again, the tangents from the same point FE , FA , are equal, as also the tangents AG , GB (cor. 2, th. 61); and the angles F and G of the isosceles triangles AFE , AGB , are equal, as well as their opposite sides AE , AB ; consequently those two triangles are identical (th. 1), and have their other sides EF , FA , AG , GB , all equal, and FG equal to the double of any one of them. In like manner it is shown, that all the other sides GH , HI , IK , KF , are equal to FG , or double of the tangents GB , BH , &c.

Hence, then, the circumscribed figure is both equilateral and equiangular, which was to be shown.

Corol.


Corol. The inscribed circle touches the middles of the sides of the polygon.

PROBLEM XXXII.

To inscribe a circle in a Regular Polygon.

BISECT any two sides of the polygon by the perpendiculars GO , FO , and their intersection O will be the centre of the inscribed circle, and OG or OF will be the radius.

For the perpendiculars to the tangents AF , AG , pass through the centre (cor. th. 47); and the inscribed circle touches the middle points F , G , by the last corollary. Also, the two sides AG , AO of the right-angled triangle AOG , being equal to the two sides AF , AO of the right-angled triangle AOF , the third sides OF , OG , will also be equal (cor. 2. th 34). Therefore the circle described with the centre O and radius OG , will pass through F , and will touch the sides in the points G and F . And the same for all the other sides of the figure.




PROBLEM XXXIII.

To Describe a Circle about a Regular Polygon.

BISECT any two of the angles, C and D , with the lines CO , DO ; then their intersection O will be the centre of the circumscribing circle; and OC , or OD , will be the radius.

For, draw OB , OA , OE , &c, to the angular points of the given polygon. Then the triangle OCD is isosceles, having the angles at C , and D equal, being the halves of the equal angles of the polygon BCD , CDE ; therefore their opposite sides CO , DO , are equal (th. 4). But the two triangles OCD , OCB , having the two sides OC , CD equal to the two OC , CB , and the included angles OCD , OCB also equal, will be identical (th. 1), and have their third sides BO , OD , equal. In like manner it is shown, that all the lines OA , OB , OC , OD OE , are equal. Consequently a circle described with the centre O and radius OA , will pass through all the other angular points, B , C , D , &c, and will circumscribe the polygon.




PROBLEM

PROBLEM XXXIV.

To make a Square Equal to the Sum of two or more Given Squares.

LET AB and AC be the sides of two given squares. Draw two indefinite lines AP, AQ, at right angles to each other; in which place the sides AB, AC, of the given squares; join BC; then a square described on BC will be equal to the sum of the two squares described on AB and AC (th. 34).



In the same manner, a square may be made equal to the sum of the three or more given squares. For, if AB, AC, AD, be taken as the sides of the given squares, then, making $AE = BC$, $AD = AD$, and drawing DE, it is evident that the square on DE will be equal to the sum of the three squares on AB, AC, AD. And so on for more squares.

PROBLEM XXXV.

To make a Square equal to the Difference of two Given Squares.

LET AB and AC, taken in the same straight line, be equal to the sides of the two given squares.—From the centre A, with the distance AB, describe a circle; and make CD perpendicular to AB, meeting the circumference in D: so shall a square described on CD be equal to $AD^2 - AC^2$, or $AB^2 - AC^2$, as required (cor. th. 34).




PROBLEM XXXVI.

To make a Triangle Equal to a Given Quadrangle ABCD.

DRAW the diagonal AC, and parallel to it DE, meeting BA produced at E, and join CE; then will the triangle CEB be equal to the given quadrilateral ABCD.

For, the two triangles ACE, ACD, being on the same base AC, and between the same parallels AC, DE, are equal (th. 25); therefore, if ABC be added to each, it will make EC equal to ABCD (ax. 2).




PROBLEM

PROBLEM XXXVII.

To make a Triangle Equal to a Given Pentagon $ABCDE$.

DRAW DA and DB , and also EF , CG , parallel to them, meeting AB produced at F and G ; then draw DF and DG ; so shall the triangle DFG be equal to the given pentagon $ABCDE$.




For the triangle $DFA = DEA$, and the triangle $DGB = DCB$ (th. 25); therefore, by adding DAB to the equals, the sums are equal (ax. 2), that is, $DAB + DAF + DBG = DAB + DAE + DBC$, or the triangle $DFG =$ to the pentagon $ABCDE$.

PROBLEM XXXVIII.

To make a Rectangle Equal to a Given Triangle ABC .


BISECT the base AB in D ; then raise DE and BF , perpendicular to AB , and meeting CF parallel to AB , at E and F : so shall DF be the rectangle equal to the given triangle ABC (by cor. 2, th. 26).



PROBLEM XXXIX.

To make a Square Equal to a Given Rectangle $ABCD$.

PRODUCE one side AB , till BE be equal to the other side BC . On AE as a diameter describe a circle, meeting BC produced at F : then will BF be the side of the square $BFGH$, equal to the given rectangle BD , as required; as appears by cor. th. 87, and th. 77.



APPLICATION

APPLICATION OF ALGEBRA

TO

GEOMETRY.

WHEN it is proposed to resolve a geometrical problem algebraically, or by algebra, it is proper, in the first place, to draw a figure that shall represent the several parts or conditions of the problem, and to suppose that figure to be the true one. Then, having considered attentively the nature of the problem, the figure is next to be prepared for a solution, if necessary, by producing or drawing such, lines in it as appear most conducive to that end. This done, the usual symbols or letters, for known and unknown quantities, are employed to denote the several parts of the figure, both the known and unknown parts, or as many of them as necessary, as also such unknown line or lines as may be easiest found, whether required or not. Then proceed to the operation, by observing the relations that the several parts of the figure have to each other; from which, and the proper theorems in the foregoing elements of geometry, make out as many equations independent of each other, as there are unknown quantities employed in them: the resolution of which equations, in the same manner as in arithmetical problems, will determine the unknown quantities, and resolve the problem proposed.

As no general rule can be given for drawing the lines, and selecting the fittest quantities to substitute for, so as always to bring out the most simple conclusion, because different problems require different modes of solution; the best way to gain experience, is to try the solution of the same problem in different ways, and then apply that which succeeds best, to other cases of the same kind, when they afterwards occur. The following particular directions, however, may be of some use.

1st. In preparing the figure, by drawing lines, let them be either parallel or perpendicular to other lines in the figure, or so as to form similar triangles. And if an angle be given, it will be proper to let the perpendicular be opposite to that angle, and to fall from one end of a given line, if possible.

2d, In selecting the quantities proper to substitute for, those are to be chosen, whether required or not, which lie nearest the known or given parts of the figure, and by means of which the next adjacent parts may be expressed by addition and subtraction only, without using surds.


3d, When two lines or quantities are alike related to other parts of the figure or problem, the best way is, not to make use of either of them separately, but to substitute for their sum, or difference, or rectangle, or the sum of their alternate quotients, or for some line or lines, in the figure, to which they have both the same relation.

4th, When the area, or the perimeter, of a figure, is given, or such parts of it as have only a remote relation to the parts required: it is sometimes of use to assume another figure similar to the proposed one, having one side equal to unity, or some other known quantity. For, hence the other parts of the figure may be found, by the known proportions of the like sides or parts, and so an equation be obtained. For examples, take the following problems.

PROBLEM I.

In a Right-angled Triangle, having given the Base (3), and the Sum of the Hypotenuse and Perpendicular (9); to find both these two Sides.

LET ABC represent the proposed triangle, right-angled at B. Put the base $AB = 3 = b$, and the sum $AC + BC$ of the hypotenuse and perpendicular $= 9 = s$; also, let x denote the hypotenuse AC , and y the perpendicular BC .



Then by the question - - - $x + y = s$,
and by theorems 34, - - - $x^2 = y^2 + b^2$.

By transpos. y in the 1st equ. gives $x = s - y$,

This value of x substi. in the 2d,

gives - - - - - $s^2 - 2sy + y^2 = y^2 + b^2$,

Taking away y^2 on both sides leaves $s^2 - 2sy = b^2$.

By transpos. $2sy$ and b^2 , gives $s^2 - b^2 = 2sy$,

And dividing by $2s$, gives - - $\frac{s^2 - b^2}{2s} = y = 4 = BC$.

Hence $x = s - y$ $5 = AC$.

N. B. In this solution, and the following ones, the notation is made by using as many unknown letters, x and y , as there

there are unknown sides of the triangle, a separate letter for each ; in preference to using only one unknown letter for one side, and expressing the other unknown side in terms of that letter and the given sum or difference of the sides ; though this latter way would render the solution shorter and sooner ; because the former way gives occasion for more and better practice in reducing equations, which is the very end and reason for which these problems are given at all.

PROBLEM II.

In a Right-angled Triangle, having given the Hypotenuse (5) ; and the sum of the Base and Perpendicular (7) ; to find both these two Sides.

LET $\triangle ABC$ represent the proposed triangle, right-angled at B . Put the given hypotenuse $AC=5=a$, and the sum $AB+BC$ of the base and perpendicular $=7=s$; also let x denote the base AB , and y the perpendicular BC .

Then by the question - - - $x+y=s$

and by theorem 34 - - - $x^2+y^2=a^2$

By transpos. y in the 1st, gives $x=s-y$

By substitu. this valu. for x , gives $s^2-2sy+2y^2=a^2$

By transposing s^2 , gives - - - $2y^2-2sy=a^2-s^2$

By dividing by 2, gives - - - $y^2-sy=\frac{1}{2}a^2-\frac{1}{2}s^2$

By completing the square, gives $y^2-sy+\frac{1}{4}s^2=\frac{1}{2}a^2-\frac{1}{4}s^2$

By extracting the root, gives - $y-\frac{1}{2}s=\sqrt{\frac{1}{2}a^2-\frac{1}{4}s^2}$


By transposing $\frac{1}{2}s$, gives - - - $y=\frac{1}{2}s\pm\sqrt{\frac{1}{2}a^2-\frac{1}{4}s^2}$

4 and 3, the values of x and y .

PROBLEM III.

In a Rectangle, having given the Diagonal (10), and the Perimeter, or Sum of all the Four Sides (28) ; to find each of the Sides severally.

LET $ABCD$ be the proposed rectangle ; and put the diagonal $AC=10=d$, and half the perimeter $AB+BC$ or $AD+DC=14=a$; also put one side $AB=x$, and the other side $BC=y$. Hence, by




right-angled

right-angled triangles, - - - - - $x^2 + y^2 = d^2$
 And by the question, - - - - - $x + y = a$
 Then by transposing y in the 2d, gives $x = a - y$
 This value substituted in the 1st, gives $a^2 - 2ay + 2y^2 = d^2$
 Transposing a^2 , gives - - $2y^2 - 2ay = d^2 - a^2$
 And dividing by 2, gives - $y^2 - ay = \frac{1}{2}d^2 - \frac{1}{2}a^2$
 By completing the square, it is $y^2 - ay + \frac{1}{4}a^2 = \frac{1}{2}d^2 - \frac{1}{4}a^2$
 And extracting the root, gives $y - \frac{1}{2}a = \sqrt{\frac{1}{2}d^2 - \frac{1}{4}a^2}$
 and transposing $\frac{1}{2}a$, gives $y = \frac{1}{2}a \pm \sqrt{\frac{1}{2}d^2 - \frac{1}{4}a^2} = 8$
 or 6, the values of x and y .

PROBLEM IV.

Having given the Base and Perpendicular of any Triangle; to find the Side of a Square Incribed in the Same.

LET ABC represent the given triangle, and EFGH its inscribed square. Put the base $AB = b$, the perpendicular $CD = a$, and the side of the square GF or $GH = DI = x$; then will $CI = CD - DI = a - x$.




Then, because the like lines in the similar triangles ABC, GFC, are proportional (by theor. 84, Geom.), $AB : CD :: GE : CI$, that is, $b : a :: x : a - x$. Hence, by multiplying extremes and means, $ab - bx = ax$, and transposing bx , gives $ab = ax + bx$; then dividing by $a + b$, gives $x = \frac{ab}{a + b} = GF$ or GH the side of the inscribed square: which therefore is of the same magnitude, whatever the species or the angles of the triangles may be.

PROBLEM V.

In an Equilateral Triangle, having given the lengths of the three Perpendiculars drawn from a certain Point within, on the three Sides; to determine the Sides.

LET ABC represent the equilateral triangle, and DE, DF, DG, the given perpendiculars from the point D. Draw the lines DA, DB, DC, to the three angular points; and let fall the perpendicular CH on the base AB. Put the three given perpendiculars $DE = a$, $DF = b$, $DG = c$, and put $x = AH$ or BH , half the side of



the

the equilateral triangle. Then is AC or $BC = 2x$, and by right angled triangles the perpendicular $CH = \sqrt{AC^2 - AH^2} = \sqrt{4x^2 - x^2} = \sqrt{3x^2} = x\sqrt{3}$.

Now, since the area or space of a rectangle, is expressed by the product of the base and height (cor. 2, th. 81 Geom.), and that a triangle is equal to half a rectangle of equal base and height (cor. 1, th. 26), it follows that

the whole triangle ABC is $= \frac{1}{2}AB \times CH = x \times x\sqrt{3} = x^2\sqrt{3}$,

the triangle $ABD = \frac{1}{2}AB \times DG = x \times c = cx$,

the triangle $BCD = \frac{1}{2}BC \times DE = x \times a = ax$,

the triangle $ACD = \frac{1}{2}AC \times DF = x \times b = bx$.

But the three last triangles make up, or are equal to, the whole former, or great triangle ;

that is, $x^2\sqrt{3} = ax + bx + cx$; hence, dividing by x , gives

$x\sqrt{3} = a + b + c$, and dividing by $\sqrt{3}$, gives

$x = \frac{a+b+c}{\sqrt{3}}$, half the side of the triangle sought.

Also, since the whole perpendicular CH is $= x\sqrt{3}$, it is therefore $= a + b + c$. That is, the whole perpendicular CH , is just equal to the sum of all the three smaller perpendiculars $DE + DF + DG$ taken together, wherever the point D is situated.

PROBLEM VI.

In a Right-angled Triangle, having given the Base (3), and the Difference between the Hypotenuse and Perpendicular (1); to find both these two Sides.

PROBLEM VII.

In a Right-angled Triangle, having given the Hypotenuse (5), and the Difference between the Base and Perpendicular (1); to determine both these two Sides.

PROBLEM VIII.

HAVING given the Area, or Measure of the Space, of a Rectangle, inscribed in a given Triangle; to determine the Sides of the Rectangle.

PROBLEM

PROBLEM IX.

IN a Triangle, having given the Ratio of the two Sides, together with both the Segments of the Base, made by a Perpendicular from the Vertical Angle ; to determine the Sides of the Triangle.

PROBLEM X.

IN a Triangle, having given the Base, the Sum of the other two Sides, and the Length of a Line drawn from the Vertical Angle to the Middle of the Base ; to find the Sides of the Triangle.

PROBLEM XI.

IN a Triangle, having given the two Sides about the Vertical Angle, with the Line bisecting that Angle, and terminating in the Base ; to find the Base.

PROBLEM XII.

To determine a Right-angled Triangle ; having given the Lengths of two Lines drawn from the acute angles, to the Middle of the opposite Sides.

PROBLEM XIII.

To determine a Right-angled Triangle ; having given the Perimeter, and the Radius of its Inscribed Circle.

PROBLEM XIV.

To determine a Triangle ; having given the Base, the Perpendicular, and the Ratio of the two Sides.

PROBLEM XV.

To determine a Right angled Triangle ; having given the Hypothenuse, and the Side of the Inscribed Square.

PROBLEM

PROBLEM XVI.

To determine the Radii of three Equal Circles, described in a given Circle, to touch each other and also the Circumference of the given Circle.

PROBLEM XVII.

In a Right-angled Triangle, having given the Perimeter or Sum of all the Sides, and the Perpendicular let fall from the Right Angle on the Hypothense; to determine the Triangle, that is, its Sides.

PROBLEM XVIII.

To determine a Right-angled Triangle; having given the Hypothense, and the Difference of two lines drawn from the two acute angles to the Centre of the Inscribed Circle.

PROBLEM XIX.

To determine a Triangle; having given the Base, the Perpendicular, and the Difference of the two other Sides.

PROBLEM XX.

To determine a Triangle; having given the Base, the Perpendicular, and the Rectangle or Product of the two Sides.

PROBLEM XXI.

To determine a Triangle; having given the Lengths of three Lines drawn from the three Angles, to the Middle of the opposite Sides.

PROBLEM XXII.

In a Triangle, having given all the three Sides; to find the Radius of the Inscribed Circle.

PROBLEM XXIII.

To determine a Right-angled Triangle; having given the Side of the Inscribed Square, and the Radius of the Inscribed Circle.

PROBLEM

PROBLEM XXIV.

To determine a Triangle, and the Radius of the Inscribed Circle ; having given the Lengths of three Lines drawn from the three Angles, to the Centre of that Circle.

PROBLEM XXV.

To determine a Right angled Triangle : having given the Hypotenuse, and the Radius of the Inscribed Circle.

PROBLEM XXVI.

To determine a Triangle ; having given the Base. the Line bisecting the Vertical Angle, and the Diameter, of the Circumscribing Circle.

PLANE TRIGONOMETRY.

DEFINITIONS.

1. **PLANE TRIGONOMETRY** treats of the relations and calculations of the sides and angles of plane triangles.

2. The circumference of every circle (as before observed in Geom. Def. 57) is supposed to be divided into 360 equal parts, called Degrees; also each degree into 60 Minutes, each minute into 60 Seconds, and so on. Hence a semicircle contains 180 degrees, and a quadrant 90 degrees.

3. The measure of an angle (Def. 58, Geom.) is an arc of any circle contained between the two lines which form that angle, the angular point being the centre; and it is estimated by the number of degrees contained in that arc.


Hence, a right angle, being measured by a quadrant, or quarter of the circle, is an angle of 90 degrees; and the sum of the three angles of every triangle, or two right angles, is equal to 180 degrees. Therefore, in a right-angled triangle, taking one of the acute angles from 90 degrees, leaves the other acute angle; and the sum of two angles, in any triangle, taken from 180 degrees, leaves the third angle; or one angle being taken from 180 degrees, leaves the sum of the other two angles.

4. Degrees are marked at the top of the figure with a small $^{\circ}$, minute with $'$, seconds with $''$, and so on. Thus $57^{\circ} 30' 12''$, denote 57 degrees 30 minutes and 12 seconds.

5. The Complement of an arc, is what it wants of a quadrant or 90° . Thus, if AD be a quadrant. then BD is the compliment of the arc AB ; and, reciprocally, AB is the compliment of BD . So that, if AE be an arc of 50° , then its complement BD will be 40° .

6. The Supplement of an arc, is what it wants of a semicircle, or 180° .

Thus, if ADE be a semicircle, then BDE is the supplement of the arc AB ; and, reciprocally, AB is the supplement of the arc BDE . So that, if AB be an arc of 50° , then its supplement BDE will be 130° .



7. The Sine, or Right Sine, of an arc, is the line drawn from one extremity of the arc, perpendicular to the diameter which passes through the other extremity. Thus, BF is the sine of the arc AB , or of the supplemental arc BDE . Hence the sine (BF) is half the chord (BG) of the double arc (BAG).

8. The Versed Sine of an arc, in the part of the diameter intercepted between the arc and its sine. So, AF is the versed sine of the arc AB , and EF the versed sine of the arc EDB .

9. The Tangent of an arc, is a line touching the circle in one extremity of that arc, continued from thence to meet a line drawn from the centre through the other extremity; which last line is called the Secant of the same arc. Thus, AH is the tangent, and CH the secant of the arc AB . Also, EF is the tangent, and CF the secant, of the supplemental arc BDE . And this latter tangent and secant are equal to the former, but are accounted negative, as being drawn in an opposite or contrary direction to the former.

10. The Cosine, Cotangent, and Cosecant, of an arc, are the sine, tangent, and secant of the complement of that arc, the *Co* being only a contraction of the word complement. Thus, the arcs AB , BD , being the complements of each other, the sine, tangent, or secant of the one of these, is the cosine, cotangent, or cosecant of the other. So, BF , the sine of AB , is the cosine of BD ; and BK , the sine of BD , is the cosine of AB : in like manner, AH , the tangent of AB , is the cotangent of BD ; and DL , the tangent of DB , is the cotangent of AB : also, CH , the secant of AB , is the cosecant of BD ; and CL , the secant of BD , is the cosecant of AB .

Corol. Hence several remarkable properties easily follow from these definitions; as,

1st, That an arc and its supplement have the same sine, tangent, and secant; but the two latter, the tangent and secant are accounted negative when the arc is greater than a quadrant or 90 degrees.

2d, When the arc is 0, or nothing, the sine and tangent are nothing, but the secant is then the radius CA , the least it can be. As the arc increases from 0, the sines, tangents, and secants, all proceed increasing, till the arc becomes a whole quadrant AD , and then the sine is the greatest it can be,
being


being the radius CD of the circle ; and both the tangent and secant are infinite.

3d, Of an arc AB , the versed sine AF , and cosine BK , or CF , together make up the radius CA of the circle.—The radius CA , the tangent AH , and the secant CH , form a right-angled triangle CAH . So also do the radius, sine, and cosine, form another right-angled triangle CBF or CBK . As also the radius, cotangent, and cosecant, another right-angled triangle CDL . And all these right-angled triangles are similar to each other.

11. The sine, tangent, or secant of an angle, is the sine, tangent, or secant of the arc by which the angle is measured, or of the degrees, &c. in the same arc or angle.

12. The method of constructing the scales of chords, sines, tangents, and secants, usually engraven on instruments, for practice, is exhibited in the annexed figure.

13. A Trigonometrical Canon, is a table showing the length of the sine, tangent, and secant, to every degree and minute of the quadrant, with respect to the radius, which is expressed by unity or 1, with any number of cyphers. The logarithms of these sines, tangents and secants, are also ranged in the tables ; and these are most commonly used, as they perform the calculations by only addition and subtraction, instead of the multiplication and division by the natural sines, &c. according to the nature of logarithms. Such a table of log. sines and tangents, as well as the logs. of common numbers, are placed at the end of the second volume, and the description and use of them are as follow ; viz. of the sines and tangents ; and the other table, of common logs. has been already explained.



Description of the Table of Log. Sines and Tangents.

In the first column of the table are contained all the arcs, or angles, for every minute in the quadrant, viz. from 1' to 45°, descending from top to bottom by the left-hand side, and then returning back by the right-hand side, ascending from bottom to top, from 45° to 90°; the degrees being set at top or bottom, and the minutes in the column. Then the sines, cosines, tangents, cotangents, of the degrees and minutes, are placed on the same lines with them, and in the annexed columns, according to their several respective names or titles, which are at the top of the columns for the degrees at the top, but at the bottom of the columns for the degrees at the bottom of the leaves. The secants and cosecants are omitted in this table, because they are so easily found from the sines and cosines; for, of every arc or angle, the sine and cosecant together make up 20 or double the radius, and the cosine and secant together make up the same 20 also. Therefore, if a secant is wanted, we have only to subtract the cosine from 20; or, to find the cosecant, take the sine from 20. And the best way to perform these subtractions, because it may be done at sight, is to begin at the left hand, and take every figure from 9, but the last or right hand figure from 10, prefixing 1, for 10, before the first figure of the remainder.

PROBLEM I.

To compute the Natural Sine and Cosine of a Given Arc.

THIS problem is resolved after various ways. One of these is as follows, viz. by means of the ratio between the diameter and circumference of a circle, together with the known series for the sine and cosine, hereafter demonstrated. Thus, the semicircumference of the circle, whose radius is 1, being 3.141592653589793, &c. the proportion will therefore be, as the number of degrees or minutes in the semicircle, is to the degrees or minutes in the proposed arc, so is 3.14159265, &c. to the length of the said arc.

This length of the arc being denoted by the letter a ; and

its

its sine and cosine by s and c ; then will these two be expressed by the two following series, viz.

$$\begin{aligned}
 s &= a - \frac{a^3}{2.3} + \frac{a^5}{2.3.4.5} - \frac{a^7}{2.3.4.5.6.7} + \&c. \\
 &= a - \frac{a^3}{6} + \frac{a^5}{120} - \frac{a^7}{5040} + \&c. \\
 c &= 1 - \frac{a^2}{2} + \frac{a^4}{2.3.4} - \frac{a^6}{2.3.4.5.6} + \&c. \\
 &= 1 - \frac{a^2}{2} + \frac{a^4}{24} - \frac{a^6}{720} + \&c.
 \end{aligned}$$

EXAM. 1. If it be required to find the sine and cosine of one minute. Then the number of minutes in 180° being 10800, it will be first, as $10800 : 1 :: 3.14159265 \&c. : .000290888208665 =$ the length of an arc of one minute. Therefore, in this case.

$$\begin{aligned}
 a &= .0002908882 \\
 \text{and } \frac{1}{6}a^3 &= .000000000004 \&c. \\
 \text{the dif. is } s &= .0002908882 \text{ the sine of 1 minute.} \\
 \text{Also, from } 1. \\
 \text{take } \frac{1}{2}a^2 &= 0.0000000423079 \&c. \\
 \text{leave } c &= .9999999577 \text{ the cosine of 1 minute.}
 \end{aligned}$$

EXAM. 2. For the sine and cosine of 5 degrees. Here, as $180^\circ : 5^\circ :: 3.14159265 \&c. : .08726646 = a$ the length of 5 degrees. Hence $a = .08726646$

$$\begin{aligned}
 -\frac{1}{6}a^3 &= - .00011076 \\
 +\frac{1}{120}a^5 &= .00000004
 \end{aligned}$$

these collected give $s = .08715574$ the sine of 5° .

And, for the cosine, $1=1$.

$$\begin{aligned}
 -\frac{1}{2}a^2 &= - .00380771 \\
 +\frac{1}{24}a^4 &= .00000241
 \end{aligned}$$

these collected give $c = .99619470$ the cosine of 5° .

After the same manner, the sine and cosine of any other arc may be computed. But the greater the arc is the slower the series will converge, in which case a greater number of terms must be taken, to bring out the conclusion to the same degree of exactness.

Or,

Or, having found the sine, the cosine will be found from it, by the property of the right-angled triangle CBF , viz: the cosine $CF = \sqrt{CB^2 - BF^2}$, or $c = \sqrt{1 - s^2}$.

There are also other methods of constructing the canon of sines and cosines, which, for brevity's sake, are here omitted.

PROBLEM II.

To compute the Tangents and Secants.

THE sines and cosines being known, or found by the foregoing problem; the tangents and secants will be easily found, from the principle of similar triangles, in the following manner:

In the first figure, where, of the arc AB , BF is the sine, CF or BK the cosine, AH the tangent, CH the secant, DL the cotangent, and CL the cosecant, the radius being CA or CB or CD ; the three similar triangles CFB , CAH , CDL , give the following proportions:

1st, $CF : FB :: CA : AH$; whence the tangent is known, being a fourth proportional to the cosine, sine, and radius.

2d, $CF : CB :: CA : CH$; whence the secant is known, being a third proportional to the cosine and radius.

3d, $BF : FC :: CD : DL$; whence the cotangent is known, being a fourth proportional to the sine, cosine, and radius.

4th, $BF : BC :: CD : CL$; whence the cosecant is known, being a third proportional to the sine and radius.

As for the log. sines, tangents, and secants, in the tables, they are only the logarithms of the natural sines, tangents, and secants, calculated as above.

HAVING given an idea of the calculation and use of sines, tangents, and secants, we may now proceed to resolve the several cases of Trigonometry; previous to which, however, it may be proper to add a few preparatory notes and observations, as below.

Note 1. There are usually three methods of resolving triangles, or the cases of trigonometry; namely, Geometrical Construction, Arithmetical Computation, and Instrumental Operation.

In the first Method, The triangle is constructed, by making the parts of the given magnitudes, namely the sides from a scale of equal parts, and the angles from a scale of chords,

or by some other instrument. Then measuring the unknown parts by the same scales or instruments, the solution will be obtained near the truth.

In the Second Method, Having stated the terms of the proportion according to the proper rule or theorem, resolve it like any other proportion, in which a fourth term is to be found from three given terms, by multiplying the second and third together, and dividing the product by the first, in working with the natural numbers; or, in working with the logarithms, add the logs. of the second and third terms together, and from the sum take the log. of the first term; then the natural number answering to the remainder in the fourth term sought.

In the Third Method, Or Instrumentally, as suppose by the log. lines on one side of the common two-foot scales; Extend the Compasses from the first term, to the second or third, which happens to be of the same kind with it; then that extent will reach from the other term to the fourth term, as required, taking both extents towards the same end of the scale.

Note 2. Every triangle has six parts, viz. three sides and three angles. And in every triangle, or case in trigonometry, there must be given three of these parts, to find the other three. Also, of the three parts that are given, one of them at least must be a side; because, with the same angles, the sides may be greater or less in any proportion.

Note 3. All the cases in trigonometry, may be comprised in three varieties only; viz.

1st, When a side and its opposite angle are given.

2d, When two sides and the contained angle are given.

3d, When the three sides are given.

For there cannot possibly be more than these three varieties of cases; for each of which it will therefore be proper to give a separate theorem, as follows:

THEOREM I.

When a Side and its Opposite Angle are two of the Given Parts.

THEN the unknown parts will be found by this theorem; viz. The sides of the triangle have the same proportion to each other, as the sines of their opposite angles have.

That is, As any one side,


Is to the sine of its opposite angle;

So is any other side,

To the sine of its opposite angle.

Demonstr.

Demonstr. For, let ABC be the proposed triangle, having AB the greatest side, and BC the least. Take $AD=BC$, considering it as a radius; and let fall the perpendiculars DE , CF , which will evidently be the sines of the angles A and B , to the radius AD or BC . Now the triangles ADE , ACF , are equiangular; they therefore have their like sides proportional, namely, $AC : CF :: AD \text{ or } BC : DE$; that is, the side AC is to the sine of its opposite angle B , as the side BC is to the sine of its opposite angle A .



Note 1. In practice, to find an angle, begin the proportion with a side opposite to a given angle. And to find a side, begin with an angle opposite to a given side.


Note 2. An angle found by this rule is ambiguous, or it is uncertain whether it be acute or obtuse, unless it be a right angle, or unless its magnitude be such as to prevent the ambiguity; because the sine answers to two angles, which are supplements to each other; and accordingly the geometrical construction forms two triangles with the same parts that are given, as in the example below; and when there is no restriction or limitation included in the question, either of them may be taken. The number of degrees in the table, answering to the sine, is the acute angle; but if the angle be obtuse, subtract those degrees from 180° , and the remainder will be the obtuse angle. When a given angle is obtuse, or a right one, there can be no ambiguity; for then neither of the other angles can be obtuse, and the geometrical construction will form only one triangle.

EXAMPLE I.

In the plane triangle ABC ,

Given $\begin{cases} AB \text{ 345 yards} \\ BC \text{ 232 yards} \\ \angle A \text{ } 37^\circ 20' \end{cases}$

Required the other parts.



1. Geometrically.

Draw an indefinite line; on which set off $AB=345$ from some convenient scale of equal parts.—Make the angle $A = 37^\circ \frac{2}{3}$.—With a radius of 232, taken from the same scale of equal parts, and centre B , cross AC in the two points C, C' .—Lastly, join BC, BC' , and the figure is constructed, which gives two triangles, and shows that the case is ambiguous.

Then,

Then, the sides AC measured by the scale of equal parts, and the angles B and c measured by the line of chords, or other instrument, will be found to be nearly as below ; viz.

$$\begin{array}{lll} \text{AC } 174 & \angle B \ 27^\circ & \angle C \ 115^\circ \frac{1}{2}. \\ \text{or } 374\frac{1}{2} & \text{or } 78\frac{1}{4} & \text{or } 64\frac{1}{2} \end{array}$$

2. Arithmetically.

First, to find the angles at c.

As side	BC	232	-	-	log.	2 365488
To sin. op.	$\angle A$	$37^\circ 20'$	-	-		9.782796
So side	AB	345	-	-		2.537819
So sin. op.	$\angle C$	$115^\circ 36'$ or $64^\circ 24'$				9.955127
	add $\angle A$	37 20				37 20
	the sum	152 56	or	101 44		
	taken from	180 00		180 00		
	leaves $\angle B$	27 04	or	78 16		

Then, to find the side AC.

As sine	$\angle A$	$37^\circ 20'$	-	-	log.	9.782796
To op. side	BC	232	-	-		2.365488
So sin.	$\angle B$	$\left\{ \begin{array}{l} 27^\circ 04' \\ 78 16 \end{array} \right.$	-	-		9.658037
			-	-		9.990829
To op. side	AC	174.07	-	-		2.240729
	or	374.56	-	-		2.573521

3. Instrumentally.

In the first proportion.—Extend the compasses from 232 to 345 on the line of numbers ; then that extent will reach, on the sines, from $37^\circ \frac{1}{3}$ to $64^\circ \frac{1}{2}$, the angle c.

In the second proportion.—Extend the compasses from $37^\circ \frac{1}{3}$ to 27° or $78^\circ \frac{1}{4}$, on the sines ; then that extent will reach, on the line of numbers, from 232, to 174 or $374\frac{1}{2}$, the two values of the side AC.

EXAMPLE II.

In the plane triangle ABC.

Given	$\left\{ \begin{array}{l} \text{AB } 365 \text{ poles} \\ \angle A \ 57^\circ \ 12' \\ \angle B \ 24 \ 45 \end{array} \right.$	Ans.	$\left\{ \begin{array}{l} \angle C \ 98^\circ \ 3' \\ \text{AC} \ 154 \ 33 \\ \text{BC} \ 309.86 \end{array} \right.$
-------	--	------	---

Required the other parts.

EXAMPLE III.

In the plane triangle ABC,

$$\text{Given } \left\{ \begin{array}{l} AC \quad 120 \text{ feet} \\ BC \quad 112 \text{ feet} \\ \angle A \quad 57^\circ 28' \end{array} \right.$$

Required the other parts.

$$\text{Ans. } \left\{ \begin{array}{l} \angle B \quad 64^\circ 35' \\ \text{or } 115 \quad 25 \\ \angle C \quad 57 \quad 57 \\ \text{or } 7 \quad 7 \\ AB \quad 112.6 \text{ feet.} \\ \text{or } 16.47 \text{ feet.} \end{array} \right.$$

THEOREM II.

When two Sides and their Contained Angle are given.

FIRST add the two given sides together, to get their sum, and subtract them, to get their difference. Next subtract the given angle from 180° , or two right angles, and the remainder will be the sum of the two other angles; then divide that by 2, which will give the half the sum of the said unknown angles. Then say,


As the sum of the two given sides,
Is to the difference of the same sides;
So is the tang. of half the sum of their op. angles,
To the tang. of half the diff. of the same angles.

Then add the half difference of the angles so found, to their half sum, and it will give the greater angle, and subtracting the same will leave the less angle; because the half sum of any two quantities, increased by their half difference, gives the greater, and diminished by it, gives the less.

Then all the angles being now known, the unknown side will be found by the former theorem.

Note. Instead of the tangent of the half sum of the unknown angles, in the third term of the proportion, may be used the cotangent of half the given angle, which is the same thing.

Demonst. Let ABC be the proposed triangle, having the two given sides AC, BC, including the given angle C. With the centre C, and radius CA, the less of these two sides, describe a semicircle, meeting the other side BC produced in D, E, and the unknown side AB, in A, G. Join AE, AD, CG, and draw DF parallel to AE.




Then BE is the sum of the two given sides AC, CB, or of EC, CB; and BD is the difference of the same two given sides AC,

AC, BC, or of CD, CB. Also, the external angle ACE, is equal to the given sum of the two internal angles CAB, CBA; but the angle ADE, at the circumference, is equal to half the angle ACE at the centre: therefore the same angle ADE is equal to half the given sum of the angles CAB, CBA. Also, the external angle AGC, of the triangle BCG, is equal to the sum of the two internal angles GCB, GBC, or the angle GCB is equal to the difference of the two angles AGC, GBC; but the angle CAB is equal to the said angle AGC, these being opposite to the equal sides AC, CG; and the angle DAB, at the circumference, is equal to half the angle DCG at the centre; therefore the angle DAB is equal to half the difference of the two angles CAB, CBA; of which it has been shown that ADE or CDA is the half sum.

Now the angle DAE, in a semicircle, is a right angle, or AE is perpendicular to AD; and DF, parallel to AE, is also perpendicular to AD: consequently AE is the tangent of CDA the half sum, and DF the tangent of DAB the half difference of the angles, to the same radius AD, by the definition of a tangent. But the tangents AE, DF, being parallel, it will be as BE : BD :: AE : DF; that is, as the sum of the sides is to the difference of the sides, so is the tangent of half the sum of the opposite angles, to the tangent of half their difference.

EXAMPLE I.

In the plane triangle ABC,
 Given $\left\{ \begin{array}{l} AB \text{ 345 yards} \\ AC \text{ 174.07 yards} \\ \angle A \text{ } 37^\circ 20' \end{array} \right.$
 Required the other parts.



1. Geometrically.

Draw $AB = 345$ from a scale of equal parts. Make the angle $A = 37^\circ 20'$. Set off $AC = 174$ by the scale of equal parts. Join BC, and it is done.

Then the other parts being measured, they are found to be nearly as follows; viz. the side BC 232 yards, the angle B 27° , and the angle C, $115^\circ \frac{1}{2}$.

2. Arithmetically.

The side AB	345	From	$180^\circ 00'$
the side AC	174.07	take $\angle A$	$37 \quad 20$
their sum	519.07	sum of c and B	142 40
their differ.	170.93	half sum of do.	71 20

As

As sum of sides AB, AC, - -	519.07	log.	2.715226
To diff. of sides AB, AC, - -	170.93	-	2.232818
So tang. half sum \angle S C and B	$71^\circ 20'$	-	10.471298
To tang. half diff. \angle S C and B	44 16	-	9.988890
these added give \angle C	115 36		
and subtr. give \angle B	27 4		
Then, by the former theorem,			
As sin. \angle C $115^\circ 36'$ or $64^\circ 24'$	-	log.	9.955126
To its op. side AB	345	-	2.537819
So sin. of \angle A $37^\circ 20'$	-	-	9.782796
To its op. side BC	232	-	2.365489

3. *Instrumentally.*

In the first proportion.—Extend the compasses from 519 to 171, on the line of numbers; then that extent will reach, on the tangents, from $71^\circ \frac{1}{3}$ (the contrary way, because the tangents are set back again from 45°) a little beyond 45, which being set so far back from 45, falls upon $44^\circ \frac{1}{4}$, the fourth term.

In the second proportion.—Extend from $64^\circ \frac{1}{3}$ to $37^\circ \frac{1}{3}$ on the sines; then that extent will reach on the numbers, from 345 to 232, the fourth term sought.

EXAMPLE II.

In the plane triangle ABC,

Given	{	AB 365 poles	Ans.	{	BC 309 86
		AC 154.33			\angle B $24^\circ 45'$
		\angle A $57^\circ 12'$			\angle C 98 3

Required the other parts.

EXAMPLE III.

In the plane triangle ABC,

Given	{	AC 120 yards	Ans.	{	AB 112.6
		BC 112 yards			\angle A $57^\circ 28'$
		\angle C $57^\circ 57'$			\angle B 64 35

Required the other parts.

THEOREM III.

When the Three Sides of a Triangle are given.

First, let fall a perpendicular from the greatest angle on the opposite sine, or base, dividing it into two segments, and the whole triangle into two right-angled triangles: then the proportion will be,

As

As the base, or sums of the segments,
 Is to the sum of the other two sides ;
 So is the difference of those sides,
 To the diff. of the segments of the base.

Then take half this difference of the segments, and add it to the half sum, or the half base; for the greater segment; and subtract the same for the less segment.


Hence, in each of the two right-angled triangles, there will be known two sides, and the right angle opposite to one of them: consequently the other angles will be found by the first theorem.

Demonstr. By theor. 35, Geom. the rectangle of the sum and difference of the two sides, is equal to the rectangle of the sum and difference of the two segments. Therefore, by forming the sides of these rectangles into a proportion by theor. 76, Geometry, it will appear that the sums and differences are proportional as in this theorem.

EXAMPLE I.

In the plane triangle ABC,

Given $\left\{ \begin{array}{l} AB \text{ 345 yards} \\ AC \text{ 232} \\ BC \text{ 174.07} \end{array} \right.$
 the sides



To find the angles.

1. Geometrically.

Draw the base $AB=345$ by a scale of equal parts. With radius 232, and centre A, describe an arc; and with radius 174, and centre B, describe another arc, cutting the former in C. Join AC, BC, and it is done.

Then, by measuring the angles, they will be found to be nearly as follows, viz.

$$\angle A \ 27^\circ, \ \angle B \ 37^\circ \frac{1}{3}, \ \text{and} \ \angle C \ 115^\circ \frac{1}{2}.$$

2. Arithmetically.

Having let fall the perpendicular CP, it will be,

As the base $AB : AC + BC :: AC - BC : AP - BP$,

that is, as $345 : 406.07 :: 57.93 : 68.18 = AP - BP$.

its half is	-	34.09
the half base is		172.50
the sum of these is		206.59 = AP
and their diff. is		138.41 = BP

Then,

Then, in the triangle, ΔPC , right-angled at P ,

As the side	AC	- -	232	-	log.	2.365488
To sin. op.	$\angle P$	- -	90°	- -		10.000000
So is the side	AP	- -	206.59	- -		2.315109
To sin. op.	$\angle ACP$	- -	$62^\circ 56'$	- -		9.949621
Which taken from		-	90 00			
			leaves the $\angle A$			27 04

Again, in the triangle ΔPC , right-angled at P ,

As the side	BC	- -	174.07	-	log.	2.240724
To sin. op.	$\angle P$	- -	90°	- -		10.000000
So is side	BP	- -	138.41	- -		2.141168
To sin. op.	$\angle BCP$	- -	$52^\circ 40'$	- -		9.900444
which taken from		-	90 00			
			leaves the $\angle B$			37 20

Also, the $\angle ACP$ $62^\circ 56'$
 added to $\angle BCP$ $52 40$
 gives the whole $\angle ACB$ $115 36$

So that all the three angles are as follow, viz.

the $\angle A$ $27^\circ 4'$; the $\angle B$ $37^\circ 20'$; the $\angle C$ $115^\circ 36'$.

3. Instrumentally.

In the first proportion.—Extend the compasses from 345 to 406, on the line of numbers; then that extent will reach, on the same line, from 58 to 68.2 nearly, which is the difference of the segments of the base.

In the second proportion.—Extend from 232 to $206\frac{1}{2}$, on the line of numbers; then that extent will reach, on the sines, from 90° to 63° .

In the third proportion.—Extend from 174 to $138\frac{1}{2}$; then that extent will reach from 90° to $52^\circ\frac{2}{3}$ on the sines.

EXAMPLE II.

In the plane triangle ΔABC ,

Given the sides	{	AB	365 poles	Ans. {	$\angle A$	$57^\circ 12'$
		AC	154 33		$\angle B$	24 45
		BC	309 86		$\angle C$	98 3
To find the angles.						

EXAMPLE

EXAMPLE III.

In the plane triangle ABC,

Given the sides $\left\{ \begin{array}{l} AB \ 120 \\ AC \ 112 \ 6 \\ BC \ 112 \end{array} \right.$

Ans. $\left\{ \begin{array}{l} \angle A \ 57^\circ \ 28' \\ \angle B \ 57 \ 57 \\ \angle C \ 64 \ 35 \end{array} \right.$

To find the angles.


The three foregoing theorems include all the cases of plane triangles, both right-angled and oblique. But there are other theorems suited to some particular forms of triangles, which are sometimes more expeditious in their use than the general ones; one of which, as the case for which it serves so frequently occurs, may be here taken, as follows :

THEOREM IV.

When a Triangle is Right-angled; any of the unknown parts may be found by the following proportions: viz.

As radius
Is to either leg of the triangle;
So is tang. of its adjacent angle,
To its opposite leg;
And so is secant of the same angle,
To the hypotenuse.

Demonstr. AB being the given leg, in the right-angled triangle ABC; with the centre A, and any assumed radius AD, describe an arc DE, and draw DF perpendicular to AB, or parallel to BC. Then it is evident, from the definitions, that DF is the tangent, and AF the secant of the arc DE, or of the angle A which is measured by that arc, to the radius AD. Then, because of the parallels BC, DF, it will be, - - - as AD : AB :: DF : BC and :: AF : AC, which is the same as the theorem is in words.



Note. The radius is equal, either to the sine of 90° , or the tangent of 45° ; and is expressed by 1, in a table of natural sines, or by 10 in the log. sines.

EXAMPLE I.

In the right-angled triangle ABC,

Given $\left\{ \begin{array}{l} \text{the leg } AB \ 162 \\ \angle A \ 53^\circ \ 7' \ 48'' \end{array} \right\}$ To find AC and BC.

1. Geometrically.

1. Geometrically.

Make $AB = 162$ equal parts, and the angle $A = 53^\circ 7' 48''$; then raise the perpendicular BC , meeting AC in c . So shall AC measure 270, and BC 216.

2. Arithmetically.

As radius	-	-	log.	10.000000
To leg. AB	-	162	-	2.209515
So tang. $\angle A$	-	$53^\circ 7' 48''$	-	10.124937
To leg. BC	-	216	-	2.334452
So secant $\angle A$	-	$53^\circ 7' 48''$	-	10.221848
To hyp. AC	-	270	-	2.431363

3. Instrumentally.

Extend the compasses from 45° to $53^\circ \frac{1}{8}$, on the tangents. Then that extent will reach from 162 to 216 on the line of numbers.

EXAMPLE II.

In the right-angled triangle ABC ,

$$\text{Given } \left\{ \begin{array}{l} \text{the leg } AB \ 180 \\ \text{the } \angle A \ 62^\circ 40' \end{array} \right. \quad \text{Ans. } \left\{ \begin{array}{l} AC \ 392.0146 \\ BC \ 348.2464 \end{array} \right.$$

To find the other two sides.

Note. There is sometimes given another method for right-angled triangles, which is this :


ABC being such a triangle, make one leg AB radius ; that is, with centre A , and distance AB , describe an arc BF . Then it is evident that the other leg BC represents the tangent, and the hypotenuse AC the secant, of the arc BF , or of the angle A .

In like manner, if the leg BC be made radius ; then the other leg AB will represent the tangent, and the hypotenuse AC the secant, of the arc, BG or angle C .

But if the hypotenuse be made radius ; then each leg will represent the sine of its opposite angle ; namely, the leg AB the sine of the arc AE or angle C , and the leg BC the sine of the arc CD or angle A .

Then the general rule for all these cases is this, namely, that the sides of the triangle bear to each other the same proportion as the parts which they represent.

And this is called, Making every side radius.



Note

Note 2. When there are given two sides of a right-angled triangle, to find the third side; this is to be found by the property of the squares of the sides, in theorem 34, Geom. viz. that the square of the hypotenuse, or longest side, is equal to both the squares of the two other sides together. Therefore, to find the longest side, add the squares of the two shorter sides together, and extract the square root of that sum; but to find one of the shorter sides, subtract the one square from the other, and extract the root of the remainder.

—◆—

OF HEIGHTS AND DISTANCES, &c.

BY the mensuration and protraction of lines and angles, are determined the lengths, heights, depths, and distances of bodies or objects.

Accessible lines are measured by applying to them some certain measure a number of times, as an inch, or foot, or yard. But inaccessible lines must be measured by taking angles, or by such-like method, drawn from the principles of geometry.


When instruments are used for taking the magnitude of the angles in degrees, the lines are then calculated by trigonometry: in the other methods, the lines are calculated from the principle of similar triangles, or some other geometrical property, without regard to the measure of the angles.

Angles of elevation, or of depression, are usually taken either with a theodolite, or with a quadrant, divided into degrees and minutes, and furnished with a plummet suspended from the centre, and two open sights fixed on one of the radii, or else with telescopic sights.

To take an Angle of Altitude and Depression with the Quadrant.

Let *A* be any object, as the sun, moon, or a star, or the top of a tower, or hill, or other eminence: and let it be required to find the measure of the angle *ABC*, which a line drawn from the object makes above the horizontal line *BC*.


Place the centre of the quadrant in the angular point, and move it



round

round there as a centre. till with one eye at D , the other being shut, you perceive the object A through the sights; then will the arc GH of the quadrant, cut off, by the plumb-line BH , be the measure of the angle ABC as required.

The angle ABC of depression of any object A , below the horizontal line BC , is taken in the same manner; except that here the eye is applied to the centre, and the measure of the angle is the arc GH , on the other side of the plumb-line.



The following examples are to be constructed and calculated by the foregoing methods, treated of in Trigonometry.

EXAMPLE I.


Having measured a distance of 200 feet, in a direct horizontal line, from the bottom of a steeple, the angle of elevation of its top, taken at that distance, was found to be $47^\circ 30'$; from hence it is required to find the height of the steeple.

Construction.

Draw an indefinite line; on which set off $AC = 200$ equal parts for the measured distance. Erect the indefinite perpendicular AB ; and draw CB so as to make the angle $C = 47^\circ 30'$; the angle of elevation; and it is done. Then AB , measured on the scale of equal parts, is nearly $218\frac{1}{2}$.

Calculation.

As radius	- -	10.000000
To AC 200	- -	2.301030
So tang. $\angle C$ $47^\circ 30'$		10.037948
To AB 218 26 required		2.338978



EXAMPLE II.

What was the perpendicular height of a cloud, or of a balloon, when its angles of elevation were 35° and 64° , as taken by two observers, at the same time, both on the same side of it, and in the same verticle plane; the distance between them being half a mile or 880 yards. And what was its distance from the said two observers?


Construction.

Construction.

Draw an indefinite ground line, on which set off the given distance $AB = 880$; then A and B are the places of the observers. Make the angle $A = 35^\circ$, and the angle $B = 64^\circ$; then the intersection of the lines at c will be the place of the balloon: whence the perpendicular cd , being let fall, will be its perpendicular height. Then by measurement are found the distances and height nearly as follow, viz. AC 1631, BC 1041, DC 936.

Calculation.

First, from $\angle B$ 64°
 take $\angle A$ 35
 leaves $\angle ACB$ 29



Then in the triangle ABC ,

As sin. $\angle ACB$	29°	-	-	-	9.685571
To op. side AB	880	-	-	-	2.944483
So sin. $\angle A$	35°	-	-	-	9.758591
To op. side BC	1041.125	-	-	-	3.017503

As sin. $\angle ACB$	29°	-	-	-	9.685571
To op. side AB	880	-	-	-	2.944483
So sin. $\angle B$ 116° or 64°		-	-	-	9.953660
To op. side AC	1631.442	-	-	-	3.212572

And in the triangle BCD ,

As sin. $\angle D$	90°	-	-	-	10.000000
To op. side BC	1041.125	-	-	-	3.017503
So sin. $\angle B$	64°	-	-	-	9.953660
To op. side CD	935.757	-	-	-	2.971163

EXAMPLE III.

Having to find the height of an obelisk standing on the top of a declivity, I first measured from its bottom a distance of 40 feet, and there found the angle, formed by the oblique plane and a line imagined to go to the top of the obelisk, 41° ; but after measuring on in the same direction 60 feet farther, the like angle was only $23^\circ 45'$. What then was the height of the obelisk?


Construction.

Construction.

Draw an indefinite line for the sloping plane or declivity, in which assume any point *A* for the bottom of the obelisk, from which set off the distance $AC = 40$, and again $CD = 60$ equal parts. Then make the angle $c = 41^\circ$, and the angle $D = 23^\circ 45'$; and the point *B* where the two lines meet will be the top of the obelisk. Therefore *AB*, joined, will be its height.

Calculation.

From the $\angle C$	$41^\circ 00'$
take the $\angle D$	$23 \ 45$
leaves the $\angle DBC$	$17 \ 15$



Then in the triangle *DBC*,

As sin. $\angle DBC$ $17^\circ 15'$	-	-	-	9.472086
To op. side <i>DC</i> 60	-	-	-	1.778151
So sin. $\angle D$ $23 \ 45$	-	-	-	9.605032
To op. side <i>CB</i> 81.488	-	-	-	1.911097

And in the triangle *ABC*,

As sum of sides <i>CB</i> , <i>CA</i> 121.488	-	-	-	2.084533
To diff. of sides <i>CB</i> , <i>CA</i> 41.488	-	-	-	1.617923
So tang. half sum $\angle SA, B$ $69^\circ 30'$	-	-	-	10.427262
To tang. half diff. $\angle SA, B$ $42 \ 24\frac{1}{2}$	-	-	-	9.960652

the diff. of these is $\angle CBA$ $27 \ 5\frac{1}{2}$

Lastly, as sin. $\angle CBA$ $27^\circ 5\frac{1}{2}$	-	-	-	9.658284
To op. side <i>CA</i> 40	-	-	-	1.602060
So sin. $\angle C$ $41^\circ 0'$	-	-	-	9.816943
To op. side <i>AB</i> 57.623	-	-	-	1.760719


EXAMPLE IV.

Wanting to know the distance between two inaccessible trees, or other objects, from the top of a tower 120 feet high, which lay in the same right line with the two objects, I took the angles formed by the perpendicular wall and lines conceived to be drawn from the top of the tower to the bottom of each tree, and found them to be 33° and 64° . What then may be the distance between the two objects?

Construction.

Construction.

Draw the indefinite ground line ED , and perpendicular to it $BA = 120$ equal parts. Then draw the two lines AC, AD , making the two angles BAC, BAD , equal to the given angles 33° and $64^\circ\frac{1}{2}$. So shall c and d be the places of the two objects.



Calculation.

First, in the right-angled triangle ABC ,

As radius	-	-	-	-	-	10·000000
To AB	-	120	-	-	-	2·079181
So tang. $\angle BAC$		33°	-	-	-	9·812517
To BC	-	77·929	-	-	-	1·891698

Then in the right-angled triangle ABD ,

As radius	-	-	-	-	-	10·000000
To AB	-	-	120	-	-	2·079181
So tang. $\angle BAD$		-	$64^\circ\frac{1}{2}$	-	-	10·321504
To BD	-	-	251·585	-	-	2·400685

From which take BC 77·929

leaves the dist. CD 173·656 as required.

EXAMPLE V.

Being on the side of a river, and wanting to know the distance to a house which was seen on the other side, I measured 200 yards in a strait line by the side of the river; and then, at each end of this line of distance, took the horizontal angle formed between the house and the other end of the line; which angles were, the one of them $68^\circ 2'$, and the other $73^\circ 15'$; What then were the distances from each end to the house?


Construction.

Draw the line $AB = 200$ equal parts. Then draw AC so as to make the angle $A = 68^\circ 2'$, and BC to make the angle $B = 73^\circ 15'$. So shall the point c be the place of the house required.

Calculation.

Calculation.

To the given $\angle A$	68°	$2'$	
add the given $\angle B$	73	15	
then their sum	141	17	
being taken from	180	0	
leaves the third $\angle C$	38	43	
Hence, As sin. $\angle C$			
To op. side AB	200	$43'$	- 9.796206
So sin. $\angle A$	68°	$2'$	- 2.301030
To op. side BC	296.54		- 2.472092
And. As sin. $\angle C$	38°	$43'$	- 9.796206
To op. side AB	200		- 2.301030
So sin. $\angle B$	73°	$15'$	- 9.981171
To op. side AC	306.19		- 2.485995



EXAM. VI. From the edge of a ditch, of 36 feet wide, surrounding a fort, having taken the angle of elevation of the top of the wall, it was found to be $62^\circ 40'$: required the height of the wall, and the length of a ladder to reach from my station to the top of it?

Ans. $\left\{ \begin{array}{l} \text{height of wall } 69.64, \\ \text{ladder, } 78.4 \text{ feet.} \end{array} \right.$

EXAM. VII. Required the length of a shoar, which being to strut 11 feet from the upright of a building, will support a jamb 23 feet 10 inches from the ground?

Ans. 26 feet 3 inches.

EXAM. VIII. A ladder 40 feet long, can be so planted, that it shall reach a window 33 feet from the ground, on one side of the street; and by turning it over, without moving the foot out of its place, it will do the same by a window 21 feet high, on the other side: required the breadth of the street?

Ans. 56.649 feet.

EXAM. IX. A maypole, whose top was broken off by a blast of wind, struck the ground at 15 feet distance from the foot of the pole: what was the height of the whole maypole, supposing the broken piece to measure 39 feet in length?

Ans. 75 feet.

EXAM. X. At 170 feet distance from the bottom of a tower, the angle of its elevation was found to be $52^\circ 30'$: required the altitude of the tower?

Ans. 221.55 feet.

EXAM. XI. From the top of a tower, by the sea-side of 143 feet high, it was observed that the angle of depression of a ship's bottom, then at anchor, measured 35° : what then was the ship's distance from the bottom of the wall?

Ans. 204.22 feet.

EXAM.

EXAM. XII. What is the perpendicular height of a hill, its angle of elevation, taken at the bottom of it, being 46° , and 200 yards farther off, on a level with the bottom, the angle was 31° ?

Ans. 286.28 yards.

EXAM. XIII. Wanting to know the height of an inaccessible tower ; at the least distance from it, on the same horizontal plane, I took its angle of elevation equal to 58° ; then going 300 feet directly from it, found the angle there to be only 32° : required its height, and my distance from it at the first station ?

Ans. $\left\{ \begin{array}{l} \text{height } 307.53 \\ \text{distance } 192.15 \end{array} \right.$

EXAM. XIV. Being on a horizontal plane, and wanting to know the height of a tower placed on the top of an inaccessible hill ; I took the angle of elevation of the top of the hill 40° , and of the top of the tower 51° ; then measuring in a line directly from it to the distance of 200 feet farther, I found the angle to the top of the towers to be $23^\circ 45'$. What then is the height of the tower ?

Ans. 93.33148 feet.

EXAM. XV. From a window near the bottom of a house, which seemed to be on a level with the bottom of a steeple, I took the angle of elevation of the top of the steeple equal 40° ; then from another window, 18 feet directly above the former, the like angle was $37^\circ 30'$: what then is the height and distance of the steeple ?

Ans. $\left\{ \begin{array}{l} \text{height } 210.44 \\ \text{distance } 250.79 \end{array} \right.$

EXAM. XVI. Wanting to know the height of, and my distance from, an object on the other side of a river, which seemed to be on a level with the place where I stood, close by the side of the river ; and not having room to measure backward, on the same plane, because of the immediate rise of the bank, I placed a mark where I stood, and measured in a direction from the object, up the ascending ground to the distance of 264 feet, where it was evident that I was above the level of the top of the object ; there the angles of depression were found to be, viz. of the mark left at the river's side 42° , of the bottom of the object, 27° , and of its top 19° . Required then the height of the object, and the distance of the mark from its bottom ?

Ans. $\left\{ \begin{array}{l} \text{height } 57.26 \\ \text{distance } 150.50 \end{array} \right.$

EXAM. XVII. If the height of the mountain called the Peak of Teneriffe be $2\frac{1}{2}$ miles, as it is nearly, and the angle taken

taken at the top of it, as formed between a plumb-line and a line conceived to touch the earth in the horizon, or farthest visible point, be $87^{\circ} 58'$; it is required from these to determine the magnitude of the whole earth, and the utmost distance that can be seen on its surface from the top of the mountain, supposing the form of the earth to be perfectly round?

$$\text{Ans. } \left\{ \begin{array}{l} \text{dist. } 140\cdot876 \\ \text{diam. } 7936 \end{array} \right\} \text{ miles.}$$

EXAM. XVIII. Two ships of war, intending to cannonade a fort, are by the shallowness of the water, kept so far from it, that they suspect their guns cannot reach it with effect. In order therefore to measure the distance, they separate from each other a quarter of a mile, or 440 yards; then each ship observes and measures the angle which the other ship and the fort subtends, which angles are $83^{\circ} 45'$ and $85^{\circ} 15'$. What then is the distance between each ship and the fort?

$$\text{Ans } \left\{ \begin{array}{l} 2292\cdot26 \text{ yards.} \\ 2298\cdot05 \end{array} \right.$$

EXAM. XIX. Being on the side of a river, and wanting to know the distance to a house which was seen at a distance on the other side; I measured out for a base 400 yards in a right line by the side of the river, and found that the two angles, one at each end of this line, subtended by the other end and the house, were $68^{\circ} 2'$ and $73^{\circ} 15'$. What then was the distance between each station and the house?

$$\text{Ans. } \left\{ \begin{array}{l} 593\cdot08 \\ 612\cdot38 \end{array} \right. \text{ yards.}$$

EXAM. XX. Wanting to know the breadth of a river, I measured a base of 500 yards in a straight line close by one side of it; and at each end of this line I found the angles subtended by the other end and a tree close to the bank on the other side of the river, to be 53° and $79^{\circ} 12'$. What then was the perpendicular breadth of the river?

$$\text{Ans. } 529\cdot48 \text{ yards.}$$

EXAM. XXI. Wanting to know the extent of a piece of water, or distance between two headlands; I measured from each of them to a certain point inland, and found the two distances to be 735 yards and 840 yards; also the horizontal angle subtended between these two lines was $55^{\circ} 40'$. What then was the distance required?

$$\text{Ans. } 741\cdot2 \text{ yards}$$

EXAM. XXII. A point of land was observed, by a ship at sea, to bear east-by-south; and after sailing north-east 12 miles, it was found to bear south-east-by-east. It is required

to

to determine the place of that headland, and the ship's distance from it at the last observation? . Ans. 26·0728 miles.

EXAM. XXIII. Wanting to know the distance between a house and a mill, which were seen at a distance on the other side of a river, I measured a base line along the side where I was, of 600 yards, and at each end of it took the angles subtended by the other end and the house and mill, which were as follow, viz. at one end the angles were $53^{\circ} 20'$ and $95^{\circ} 20'$, and at the other end the like angles were $53^{\circ} 30'$ and $98^{\circ} 45'$. What then was the distance between the house and mill?

Ans. 959·5866 yards.

EXAM. XXIV. Wanting to know my distance from an inaccessible object O, on the other side of a river; and having no instrument for taking angles, but only a chain or chord for measuring distances; from each of two stations, A and B, which were taken at 500 yards asunder, I measured in a direct line from the object O 100 yards, viz. AC and BD each equal to 100 yards; also the diagonal AD measured 550 yards, and the diagonal BC 560. What then was the distance of the object O from each station A and B?

Ans. $\left\{ \begin{array}{l} AO \ 536\cdot25 \\ BO \ 500\cdot09 \end{array} \right.$

EXAM. XXV. In a garrison besieged are three remarkable objects, A, B, C, the distances of which from each other are discovered by means of a map of the place, and are as follow, viz. AB $266\frac{1}{4}$, AC 530, BC $327\frac{1}{2}$ yards. Now, having to erect a battery against it, at a certain spot without the place, and being desirous to know whether the distances from the three objects be such, as that they may from thence be battered with effect, I took, with an instrument, the horizontal angles subtended by these objects from my station s, and found them to be as follow, viz. the angle ASB $13^{\circ} 30'$, and the angle BSC $29^{\circ} 50'$; required the three distances, SA, SB, SC; the object B being situated nearest to me, and between the two others A and C?

Ans. $\left\{ \begin{array}{l} SA \ 757\cdot14 \\ SB \ 537\cdot10 \\ SC \ 655\cdot30 \end{array} \right.$

EXAM. XXVI. Required the same as in the last example, when the object B is the farthest from my station, but still seen between the two others as to angular position, and those angles being thus, the angles ASB $33^{\circ} 45'$, and BSC $22^{\circ} 30'$, also the three distances, AB 600, AC 800, BC 400 yards?


Ans. $\left\{ \begin{array}{l} SA \ 709\frac{1}{3} \\ SB \ 1042\frac{2}{3} \\ SC \ 934 \end{array} \right.$

MENSURATION OF PLANES.

THE area of any plane figure, is the measure of the space contained within its extremes or bounds ; without any regard to thickness.

This area, or the content of the plane figure, is estimated by the number of little squares that may be contained in it ; the side of those little measuring squares being an inch, a foot, a yard, or any other fixed quantity. And hence the area or content is said to be so many square inches, or square feet, or square yards, &c.

Thus, if the figure to be measured be the rectangle $ABCD$, and the little square E , whose side is one inch, be the measuring unit proposed : then as often as the said little square is contained in the rectangle, so many square inches the rectangle is said to contain which in the present case is 12.



PROBLEM I.

To find the Area of any Parallelogram ; whether it be a Square, a Rectangle, a Rhombus, or a Rhomboid.

MULTIPLY the length by the perpendicular breadth, or height, and the product will be the area.*

EXAMPLES.

* The truth of this rule is proved in the Geom. theor. 81, cor. 2.

The same is otherwise proved thus : Let the foregoing rectangle be the figure proposed : and let the length and breadth be divided into several parts each equal to the linear measuring unit, being here 4 for the length, and 3 for the breadth ; and let the opposite points of division be connected by right lines.—Then it is evident that these lines divide the rectangle into a number of little squares, each equal to the square measuring unit E ; and further, that the number of these little squares, or the area of the figure, is equal to the number of linear measuring units

EXAMPLES.

Ex. 1. To find the area of a parallelogram, the length being 12·25, and height 8 5.

$$\begin{array}{r}
 12\cdot25 \text{ length} \\
 8\cdot5 \text{ breadth} \\
 \hline
 6125 \\
 9800 \\
 \hline
 104\cdot125 \text{ area} \\
 \hline
 \end{array}$$

Ex. 2. To find the area of a square, whose side is 35·25 chains. Ans. 124 acres, 1 rood, 1 perch.

Ex. 3. To find the area of a rectangular board, whose length is $12\frac{1}{2}$ feet, and breadth 9 inches. Ans. $9\frac{3}{8}$ feet.

Ex. 4. To find the content of a piece of land, in form of a rhombus, its length being 6·20 chains, and perpendicular height 5·45. Ans. 3 acres, 1 rood, 20 perches.

Ex 5. To find the number of square yards of painting in a rhomboid, whose length is 37 feet, and breadth 5 feet 3 inches. Ans. $21\frac{7}{2}$ square yards.

PROBLEM II.

To find the Area of a Triangle.

RULE I. MULTIPLY the base by the perpendicular height, and take half the product for the area.* Or, multiply the one of these dimensions by half the other.

in the length, repeated as often as there are linear measuring units in the breadth, or height; that is, equal to the length drawn into the height; which here is 4×3 or 12.

And it is proved. (Geom. theor. 25, cor. 2), that any oblique parallelogram is equal to a rectangle, of equal length and perpendicular breadth. Therefore the rule is general for all parallelograms whatever.

* The truth of this rule is evident, because any triangle is the half of a parallelogram of equal base and altitude, by Geom. theor. 26.

EXAMPLES.

EXAMPLES.

Ex. 1. To find the area of a triangle, whose base is 625, and perpendicular height 520 links ?

Here $625 \times 260 = 162500$ square links,
or equal 1 acre, 2 roods, 20 perches, the answer.

Ex. 2. How many square yards contains the triangle, whose base is 40, and perpendicular 30 feet ?

Ans. $66\frac{2}{3}$ square yards.

Ex. 3. To find the number of square yards in a triangle, whose base is 49 feet, and height $25\frac{1}{4}$ feet ?

Ans. $68\frac{5}{8}$, or 68.7361.

Ex. 4. To find the area of a triangle, whose base is 18 feet 4 inches, and height 11 feet 10 inches ?

Ans. 108 feet, $5\frac{2}{3}$ inches.

RULE II. When two sides and their contained angle are given: Multiply the two given sides together, and take half their product: Then say, as radius is to the sine of the given angle, so is that half product, to the area of the triangle.

Or, multiply that half product by the natural sine of the said angle, for the area.*

Ex. 1. What is the area of a triangle, whose two sides are 30 and 40, and their contained angle $28^{\circ} 57'$?

By Natural Numbers.

First $\frac{1}{2} \times 40 \times 30 = 600$,
then, $1 : 600 :: .484046 \sin. 28^{\circ} 57'$
600

By Logarithms.


log. 9.684887
2.778151

Answer 290.4276 the area answering 2.463038

Ex. 2. How many square yards contains the triangle, of which one angle is 45° , and its containing sides 25 and $21\frac{1}{4}$ feet ?

Ans. 20.86947.


* For, let AB, AC, be the two given sides, including the given angle A. Now $\frac{1}{2} AB \times CP$ is the area, by the first rule, CP being the perpendicular. But, by trigonometry; as $\sin \angle P$, or radius : AC :: $\sin \angle A$: CP, which is therefore $= AC \times \sin. \angle A$, taking radius = 1. Therefore the area $\frac{1}{2} AB \times CP$ is $= \frac{1}{2} AB \times AC \times \sin. \angle A$, to radius 1; or, as radius : $\sin. \angle A$:: $\frac{1}{2} AB \times AC$: the area.



RULE III.

RULE III. When the three sides are given: Add all the three sides together, and take half that sum. Next, subtract each side severally from the said half sum, obtaining three remainders. Then multiply the said half sum and those three remainders all together, and extract the square root of the last product, for the area of the triangle.*

* For let ABC be the given triangle. Draw the parallels AE, BD , meeting the two sides AC, CB , produced, in D and E , and making $CD = CB$, and $CE = CA$. Also draw CFG bisecting DB and AE perpendicularly in F and G ; and FHI parallel to the side AB , meeting AC in H , and AE produced in I . Lastly, with centre H , and radius HF , describe a circle meeting AC produced in K ; which will pass through G , because G is a right angle, and through I , because, by means of the parallels, $AI = FB = DF$, therefore $HD = HA$, and $HF = HI = \frac{1}{2} AB$.



Hence HA or HD is half the difference of the sides AC, CB , and $HC =$ half their sum or $= \frac{1}{2} AC + \frac{1}{2} CB$; also $HK = HI = \frac{1}{2} IF$ or $\frac{1}{2} AB$; conseq. $CK = \frac{1}{2} AC + \frac{1}{2} CE + \frac{1}{2} AB$ half the sum of all the three sides of the triangle ABC , or $CK = \frac{1}{2} s$, calling s the sum of those three sides. Again $HK = HI = \frac{1}{2} IF = \frac{1}{2} AB$ or $KL = AB$; theref. $CL = CK - KL = \frac{1}{2} s - AB$, and $AK = CK - CA = \frac{1}{2} s - AC$, and $AL = DK = CK - CD = \frac{1}{2} s - CB$.

Now, by the first rule, $AG \cdot CG =$ the ΔACE , and $AG \cdot FG =$ the ΔABE , theref. $AG \cdot CF = \Delta ACB$. Also by the parallels, $AG : CG :: DF$ or $IA : CF$, theref. $AG \cdot CF = (\Delta ACB) = CG \cdot IA = CG \cdot DF$, conseq. $AG \cdot CF : CG \cdot DF = \Delta^2 ACB$.

But $CG \cdot CF = CK \cdot CL = \frac{1}{2} s \cdot \frac{1}{2} s - AB$, and $AG \cdot DF = AK \cdot AL = \frac{1}{2} s - AC \cdot \frac{1}{2} s - BC$; theref. $AG \cdot CF \cdot CG \cdot DF = \Delta^2 ACB = \frac{1}{2} s \cdot \frac{1}{2} s - AB \cdot \frac{1}{2} s - AC \cdot \frac{1}{2} s - BC$ is the square of the area of the triangle ABC . $\Omega. E. D.$

Otherwise.

Because the rectangle $AG \cdot CF =$ the ΔABC , and since $CG : AG :: CF : DF$, drawing the first and second terms into CF , and the third and fourth into AG , the propor. becomes $CG \cdot CF : AG \cdot CF :: AG \cdot CF : AG \cdot DF$, or $CG \cdot CF : \Delta ABC :: \Delta ABC : AG \cdot DF$, that is, the ΔABC is a mean proportional between $CG \cdot CF$ and $AG \cdot DF$, or between $\frac{1}{2} s \cdot \frac{1}{2} s - AB$ and $\frac{1}{2} s - AC \cdot \frac{1}{2} s - BC$. $\Omega. E. D.$

Ex. 1.

Ex. 1. To find the area of the triangle whose three sides are 20, 30, 40.

20	45	45	45
30	20	30	40
40	—	—	—
2)90	25 1st rem.	15 2d rem.	5 3d rem.
45 half sum	—	—	—

Then $45 \times 25 \times 15 \times 5 = 84375$,
The root of which is 290.4737, the area.

Ex. 2. How many square yards of plastering are in a triangle, whose sides are 30, 40, 50, feet? Ans. $66\frac{2}{3}$.

Ex. 3. How many acres, &c. contains the triangle, whose sides are 2569, 4900, 5025 links?

Ans. 61 acres, 1 rood, 39 perches.

PROBLEM III.

To find the Area of a Trapezoid.

ADD together the two parallel sides; then multiply their sum by the perpendicular breadth, or the distance between them; and take half the product for the area. By Geom. theor. 29.

Ex. 1. In a trapezoid, the parallel sides are 750 and 1225, and the perpendicular distance between them 1540 links: to find the area.

1225
750

—————
 $1975 \times 770 = 152075$ square links = 15 acr. 33 perc.

Ex. 2. How many square feet are contained in the plank, whose length is 12 feet 6 inches, the breadth at the greater end 15 inches, and at the less end 11 inches?

Ans. $131\frac{3}{4}$ feet.

Ex. 3. In measuring along one side AB of a quadrangular field, that side, and the two perpendiculars let fall on it from the two opposite corners, measured as below, required the content.

AP = 110 links


AQ = 745

AB = 1110

CP = 352

DQ = 595

Ans. 4 acres, 1 rood, 5.792 perches



PROBLEM IV.

To find the Area of any Trapezium.

DIVIDE the trapezium into two triangles by a diagonal; then find the areas of these triangles, and add them together.

Or thus, let fall two perpendiculars on the diagonal from the other two opposite angles; then add these two perpendiculars together, and multiply that sum by the diagonal, taking half the product for the area of the trapezium.

Ex. 1. To find the area of the trapezium, whose diagonal, is 42, and the two perpendiculars on it 16 and 18.

Here $16 + 18 = 34$, its half is 17.

Then $42 \times 17 = 714$ the area.

Ex. 2. How many square yards of paving are in the trapezium, whose diagonal is 65 feet, and the two perpendiculars let fall on it 28 and $33\frac{1}{2}$ feet?

Ans. $222\frac{1}{2}$ yards.

Ex. 3. In the quadrangular field ABCD, on account of obstructions there could only be taken the following measures, viz. the two sides BC 265 and AD 220 yards, the diagonal AC 378, and the two distances of the perpendiculars from the ends of the diagonal, namely. AE 100, and CF 70 yards. Required the construction of the figure, and the area in acres, when 4840 square yards make an acre?

Ans. 17 acres, 2 roods, 21 perches.

PROBLEM V.

To find the Area of an Irregular Polygon.


DRAW diagonals dividing the proposed polygon into trapeziums and triangles. Then find the areas of all these separately, and add them together for the content of the whole polygon.

EXAMPLE

EXAMPLE. To find the content of the irregular figure ABCDEFGA, in which are given the following diagonals and perpendiculars namely,

AC 55
 FD 52
 GC 44
 gm 13
 BN 18
 GO 12
 Ep 8
 Dq 23

Ans. $1878\frac{1}{2}$



PROBLEM VI.

To find the Area of a Regular Polygon.

RULE I. MULTIPLY the perimeter of the polygon, or sum of its sides, by the perpendicular drawn from its centre on one of its sides, and take half the product for the area.*

Ex. I. To find the area of the regular pentagon, each side being 25 feet, and the perpendicular from the centre on each side is $17\cdot2047737$.

Here $25 \times 5 = 125$ is the perimeter.

And $17\cdot2047737 \times 125 = 2150\cdot5967125$.

Its half $1075\cdot298356$ is the area sought.

RULE II. Square the side of the polygon; then multiply that square by the tabular area, or multiplier set against its name in the following table, and the product will be the area.†

No.

* This is only in effect resolving the polygon into as many equal triangles as it has sides, by drawing lines from the centre to all the angles; then finding their areas, and adding them all together.

† This rule is founded on the property, that like polygons, being similar figures, are to one another as the squares of their like sides; which is proved in the Geom. theor. 89. Now, the multipliers in the table, are the areas of the respective polygons to the side 1. Whence the rule is manifest.

Note.

No. of Sides.	Names.	Areas, or Multipliers.
3	Trigon or triangle	0.4330127
4	Tetragon or square	1.0000000
5	Pentagon	1.7204774
6	Hexagon	2.5980762
7	Heptagon	3.6339124
8	Octagon	4.8284271
9	Nonagon	6.1818242
10	Decagon	7.6942088
11	Undecagon	9.3656399
12	Dodecagon	11.1961524

EXAM. Taking here the same example as before, namely, a pentagon, whose side is 25 feet.

Then 25^2 being = 625,

And the tabular area 1.7204774 ;

Theref. $1.7204774 \times 625 = 1075.293375$, as before,


Ex. 2. To find the area of the trigon, or equilateral triangle, whose side is 20. Ans. 173.20508.

Ex. 3. To find the area of the hexagon whose side is 20. Ans. 1039.23048.

Ex. 4. To find the area of an octagon whose side is 20. Ans. 1931.37084.

Ex. 5. To find the area of a decagon whose side is 20. Ans. 3077.68352.

Note. The areas in the table, to each side 1, may be computed in the following manner: From the centre c of the polygon draw lines to every angle, dividing the whole figure into as many equal triangles as the polygon has sides; and let abc be one of those triangles, the perpendicular of which is cd . Divide 360 degrees by the number of sides in the polygon, the quotient gives the angle at the centre acb . The half of this gives the angle acd ; and this taken from 90° , leaves the angle cad . Then it will be, as radius is to ad , so is tang. angle cad , to the perpendicular cd . This perpendicular, multiplied by the half base ad , gives the area of the triangle abc ; which being multiplied by the number of the triangles, or of the sides of the polygon, gives its whole area, as in the table, for every one of the figures.



PROBLEM VII.

To find the Diameter and Circumference of any Circle, the one from the other.

THIS may be done nearly by either of the two following proportions,

viz. As 7 is to 22, so is the diameter to the circumference.

Or, As 1 is to 3.1416, so is the diameter to the circumference.*


Ex 1. To find the circumference of the circle whose diameter is 20.

By the first rule, as $7 : 22 :: 20 : 62\frac{2}{7}$, the answer.

Ex. 2.

* For, let ABCD be any circle, whose centre is E, and let AB, BC be any two equal arcs. Draw the several chords as in the figure, and join BE; also draw the diameter DA, which produce to F, till BF be equal to the chord BD.

Then the two isosceles triangles DEB, DBF, are equiangular, because they have the angle at D common; consequently $DE : DB :: DB : DF$. But the two triangles AFB, DCB are identical, or equal in all respects, because they have the angle F = the angle BDC, being each equal to the angle ADB, these being subtended by the equal arcs AB, BC; also the exterior angle FAB of the quadrangle ABCD, is equal to the opposite interior angle at C; and the two triangles have also the side BF = the side BD; therefore the side AF is also equal to the side DC. Hence the proportion above, viz. $DE : DB :: DB : DF = DA + AF$, becomes $DE : DB :: DB : 2DE + DC$. Then, by taking the rectangles of the extremes and means, it is $DB^2 = 2DE^2 + DE \cdot DC$.



Now, if the radius DE be taken = 1, this expression becomes $DB^2 = 2 + DC$, and hence the root $DB = \sqrt{2 + DC}$. That is, If the measure of the supplemental chord of any arc be increased by the number 2, the square root of the sum will be the supplemental chord of half that arc.

Now, to apply this to the calculation of the circumference of the circle, let the arc AC be taken equal to $\frac{1}{6}$ of the circumference, and be successively bisected by the above theorem: thus, the chord AC of $\frac{1}{6}$ of the circumference, is the side of the inscribed regular hexagon, and is therefore equal to the radius AE or 1: hence, in the right-angled triangle ACD, it will be $DC = \sqrt{AD^2 - AC^2} = \sqrt{2^2 - 1^2} = \sqrt{3} = 1.7320508076$, the supplemental chord of $\frac{1}{6}$ of the periphery.

Then,

Ex. 2. If the circumference of the earth be 25000 miles, what is its diameter ?

By the 2d rule, as $3.1416 : 1 :: 25000 : 7957\frac{3}{8}$ nearly the diameter.

NOTE BY R. ADRAIN. Having applied my new theory of most probable values to the determination of the magnitude and figure of the earth. I found the true mean diameter of the earth, taken as a globe, to be 7918.7 English miles, and consequently its circumference 24877.4 E. miles, and a degree of a great circle equal to 69.1039 miles.

Then, by the foregoing theorem, by always bisecting the arcs, and adding 2 to the last square root, there will be found the supplemental chords of the 12th, the 24th, the 48th, the 96th, &c. parts of the periphery ; thus,

$\sqrt{3.7320508076} = 1.9318516525$	}	for the supplement-	tal chord of	}	$\frac{1}{12}$	}	the periphery.
$\sqrt{3.9318516525} = 1.9828897227$					$\frac{1}{24}$		
$\sqrt{3.9828897227} = 1.9957178465$					$\frac{1}{48}$		
$\sqrt{3.9957178465} = 1.9989291743$					$\frac{1}{96}$		
$\sqrt{3.9989291743} = 1.9997322757$					$\frac{1}{192}$		
$\sqrt{3.9997322757} = 1.9999330678$					$\frac{1}{384}$		
$\sqrt{3.9999330678} = 1.9999832669$					$\frac{1}{768}$		
$\sqrt{3.9999832669} = - - - - -$					$\frac{1}{1536}$		

Since then it is found that 3.9999832669 is the square of the supplemental chord of the 1536th part of the periphery, let this number be taken from 4, which is the square of the diameter, and the remainder 0.0000167331 will be the square of the chord of the said 1536th part of the periphery, and consequently the root $\sqrt{0.0000167331} = 0.0040906112$ is the length of that chord ; this number then being multiplied by 1536, gives 6.2831788 for the perimeter of a regular polygon of 1536 sides inscribed in the circle ; which, as the sides of the polygon nearly coincide with the circumference of the circle, must also express the length of the circumference itself, very nearly.

But now, to show how near this determination is to the truth, let $AQ = 0.0040906112$ represent one side of such a regular polygon of 1536 sides, and SRT a side of another similar polygon described about the circle ; and from the centre E let the perpendicular EQR be drawn, bisecting AP and ST in Q and R . Then since AQ is $= \frac{1}{2} AP = 0.0020453056$, and $EA = 1$, therefore $EQ^2 = EA^2 - AQ^2 = .9999958167$, and consequently its root gives $EQ = .9999979034$; then because of the parallels AP, ST , it is $EQ : ER :: AP : ST ::$ as the whole inscribed perimeter : to the circumscribed one, that is, as .9999979034 :



$1 : 6.2831788 : 6.2831920$ the perimeter of the circumscribed polygon. Now, the

PROBLEM VIII.

To find the Length of any Arc of a Circle.

MULTIPLY the decimal $\cdot 01745$ by the degrees in the given arc, and that product by the radius of the circle, for the length of the arc.*

Ex. 1. To find the length of an arc of 30 degrees, the radius being 9 feet. Ans. 4.7115.

Ex. 2. To find the length of an arc of $12^\circ 10'$, or $12\frac{1}{6}^\circ$, the radius being 10 feet. Ans. 2.1231.

PROBLEM IX.

To find the Area of a Circle.†

RULE I. **MULTIPLY** half the circumference by half the diameter. Or multiply the whole circumference by the whole diameter, and take $\frac{1}{4}$ of the product.

RULE

the circumference of the circle being greater than the perimeter of the inner polygon, but less than that of the outer, it must consequently be greater than

6.2831783,

but less than 6.2831920,

and must therefore be nearly equal $\frac{1}{2}$ their sum, or

6.2831854,*

which in fact is true to the last figure, which should be a 3 instead of the 4.

Hence, the circumference being 6.2831854 when the diameter is 2, it will be the half of that, or 3.1415927, when the diameter is 1, to which the ratio in the rule, viz. 1 to 3.1416 is very near. Also the other ratio in the rule, 7 to 22 or 1 to $3\frac{1}{7} = 3.1428$ &c. is another near approximation.

* It having been found, in the demonstration of the foregoing problem, that when the radius of a circle is 1, the length of the whole circumference is 6.2831854, which consists of 360 degrees; therefore as $360^\circ : 6.2831854 :: 1^\circ : \cdot 01745$ &c. the length of the arc of 1 degree. Hence the decimal $\cdot 01745$ multiplied by any number of degrees, will give the length of the arc of those degrees. And because the circumferences and arcs are in proportion as the diameters, or as the radii of the circles, therefore as the radius 1 is to any other radius r , so is the length of the arc above mentioned, to $\cdot 01745 \times$ degrees in the arc $\times r$, which is the length of that arc, as in the rule.

† The first rule is proved in the Geom. theor. 94.

And the 2d and 3d rules are deduced from the first rule, in this manner.—By that rule, $dc, \div 4$ is the area, when d denotes the diameter, and c the circumference.

RULE II. Square the diameter, and multiply that square by the decimal $\cdot 7854$, for the area.

RULE III. Square the circumference, and multiply that square by the decimal $\cdot 07958$.

Ex. 1. To find the area of a circle whose diameter is 10, and its circumference 31.416.

By Rule 1.	By Rule 2.	By Rule 3.
31.416	$\cdot 7854$	31.416
10	$10^2 = 100$	31.416
4)314.16	78.54	986.965
78.54	78.54	$\cdot 07958$
78.54	78.54	78.54

So that the area is 78.54 by all the three rules.

Ex. 2. To find the area of a circle, whose diameter is 7, and circumference 22. Ans. $38\frac{1}{2}$.

Ex. 3. How many square yards are in a circle whose diameter is $3\frac{1}{2}$ feet: Ans. 1.069.

Ex. 4. To find the area of a circle whose circumference is 12 feet. Ans. 11.4595.

PROBLEM X.

To find the Area of a Circular Ring, or of the Space included between the Circumferences of two Circles; the one being contained within the other.

TAKE the difference between the areas of the two circles, as found by the last problem, for the area of the ring.—Or,

ference. But, by prob. 7, c is $= 3.1416d$; therefore the said area $dc \div 4$, becomes $d \times 3.1416d \div 4 = .7854d^2$, which gives the second rule.—Also, by the same prob. 7, d is $= c \div 3.1416$; therefore again the same first area $dc \div 4$, becomes $c \div 3.1416 \times c \div 4 = c^2 \div 12.5664$, which is $= c^2 \times .07958$, by taking the reciprocal of 12.5664, or changing that divisor into the multiplier $\cdot 07958$; which gives the 3d rule.

Corol. Hence, the areas of different circles are in proportion to one another, as the square of their diameters, or as the square of their circumferences; as before proved in the Geom. theor. 93.

which

which is the same thing, subtract the square of the less diameter from the square of the greater, and multiply their difference by $\cdot 7054$.—Or lastly, multiply the sum of the diameters by the difference of the same, and that product by $\cdot 7854$; which is still the same thing, because the product of the sum and difference of any two quantities, is equal to the difference of their squares.

Ex. 1. The diameters of two concentric circles being 10 and 6, required the area of the ring contained between their circumferences.

Here $10 + 6 = 16$ the sum, and $10 - 6 = 4$ the diff.
Therefore $\cdot 7854 \times 16 \times 4 = \cdot 7854 \times 64 = 50\cdot 2656$.
the area.

Ex. 2. What is the area of the ring, the diameters of whose bounding circles are 10 and 20? **Ans.** $235\cdot 62$.

PROBLEM XI.

To find the Area of the Sector of a Circle.

RULE I. MULTIPLY the radius, or half the diameter, by half the arc of the sector, for the area. Or, multiply the whole diameter by the whole arc of the sector, and take $\frac{1}{4}$ of the product. The reason of which is the same as for the first rule to problem 9, for the whole circle.

RULE II. Compute the area of the whole circle: then say, as 360 is to the degrees in the arc of the sector, so is the area of the whole circle to the area of the sector.

This is evident, because the sector is proportional to the length of the arc, or to the degrees contained in it.

Ex. 1. To find the area of a circular sector, whose arc contains 18 degrees; the diameter being 3 feet?

1. By the 1st Rule.

First, $3\cdot 1416 \times 3 = 9\cdot 4248$, the circumference.

And $360 : 18 :: 9\cdot 4248 : \cdot 47124$, the length of the arc.

Then $\cdot 47124 \times 3 \div 4 = 1\cdot 41372 \div 4 = \cdot 35343$, the area.

2. By the 2d Rule.

First, $\cdot 7854 \times 3^2 = 7\cdot 0686$, the area of the whole circle.

Then, as $360 : 18 :: 7\cdot 0686 : \cdot 35343$, the area of the sector.

Ex. 2.

Ex. 2. To find the area of a sector, whose radius is 10, and arc 20. Ans. 100.

Ex. 3. Required the area of a sector, whose radius is 25, and its arc containing $147^{\circ} 29'$. Ans. 804.3986.

PROBLEM XII.

To find the Area of a Segment of a Circle.


RULE I. FIND the area of the sector having the same arc with the segment, by the last problem.

Find also the area of the triangle, formed by the chord of the segment and the two radii of the sector.

Then add these two together for the answer, when the segment is greater than a semicircle; or subtract them when it is less than a semicircle.—As is evident by inspection.

Ex. 1. To find the area of the segment ACBDA, its chord AB being 12, and the radius AE or CE 10.

First, As $AE : \sin. \angle D 90^{\circ} :: AD : \sin. 36^{\circ} 52' \frac{1}{2} = 36.87$ degrees, the degrees in the $\angle AEC$ or arc AC. Their double, 73.74 , are the degrees in the whole arc ACB.



Now $.7854 \times 400 = 314.16$, the area of the whole circle.

Therefore $360^{\circ} : 73.74 :: 314.16 : 64.3504$, area of the sector ACBE.

Again, $\sqrt{AE^2 - AD^2} = \sqrt{100 - 36} = \sqrt{64} = 8 = DE$.

Theref. $AD \times DE = 6 \times 8 = 48$, the area of the triangle AEB.

Hence sector ACBE — triangle AEB = 16.3504 , area of seg. ACBDA.

RULE II. Divide the height of the segment by the diameter, and find the quotient in the column of heights in the following tablet: Take out the corresponding area in the next column on the right hand; and multiply it by the square of the circle's diameter, for the area of the segment.*

Note.

* The truth of this rule depends on the principle of similar plane figures, which are to one another as the square of their like linear dimensions. The segments in the table are those of a circle whose diameter is 1; and the first column contains

Note. When the quotient is not found exactly in the table, proportion may be made between the next less and greater area, in the same manner as is done for logarithms, or any other table.

Table of the Areas of Circular Segments.

Height.	Area of the Segm.	Height.	Area of the Segm.	Height.	Area of the Segm.	Height.	Area of the Segm.	Height.	Area of the Segm.
·01	·00133	·11	·04701	·21	·11990	·31	·20738	·41	·30319
·02	·00375	·12	·05339	·22	·12811	·32	·21667	·42	·31304
·03	·00687	·13	·06000	·23	·13646	·33	·22603	·43	·32293
·04	·01054	·14	·06683	·24	·14494	·34	·23547	·44	·33284
·05	·01468	·15	·07387	·25	·15354	·35	·24498	·45	·34278
·06	·01924	·16	·08111	·26	·16226	·36	·25455	·46	·35274
·07	·02417	·17	·08853	·27	·17109	·37	·26418	·47	·36272
·08	·02941	·18	·09613	·28	·18002	·38	·27386	·48	·37270
·09	·03502	·19	·10390	·29	·18905	·39	·28359	·49	·38270
·10	·04088	·20	·11182	·30	·19817	·40	·29337	·50	·39270

Ex. 2. Taking the same example as before, in which are given the chord AB 12, and the radius 10, or diameter 20.

And having found, as above, $DE = 8$; then $CE - DE = CD = 10 - 8 = 2$. Hence, by the rule, $CD \div CE = 2 \div 20 = \cdot 1$ the tabular height. This being found in the first column of the table, the corresponding tabular area is $\cdot 04088$. Then $\cdot 04088 \times 20^2 = \cdot 04088 \times 400 = 16\cdot 352$, the area, nearly the same as before.

Ex. 3. What is the area of the segment, whose height is 18, and diameter of the circle 50? Ans. 636·375.

Ex. 4. Required the area of the segment whose chord is 16, the diameter being 20? Ans. 44·728.

contains the corresponding heights or versed sines divided by the diameter. Thus then, the area of the similar segment, taken from the table, and multiplied by the square of the diameter, gives the area of the segment to this diameter.

PROBLEM XIII.

To measure long Irregular Figures.

TAKE or measure the breadth at both ends, and at several places at equal distances. Then add together all these intermediate breadths and half the two extremes, which sum multiply by the length, and divide by the number of parts for the area.*

Note. If the perpendiculars or breadths be not at equal distances, compute all the parts separately, as so many trapezoids, and add them all together, for the whole area.


Or else, add all the perpendicular breadths together, and divide their sum by the number of them for the mean breadth, to multiply by the length; which will give the whole area, not far from the truth.

Ex. 1. The breadths of an irregular figure, at five equidistant places, being 8.2, 7.4, 9.2, 10.2, 8.6; and the whole length 39; required the area?

<table style="width: 100%; border-collapse: collapse;"> <tr><td style="text-align: right;">8.2</td><td style="width: 10px;"></td></tr> <tr><td style="text-align: right;">8.6</td><td style="width: 10px;"></td></tr> <tr><td colspan="2" style="border-top: 1px solid black;"></td></tr> <tr><td style="text-align: right;">2) 16.8 sum of the extremes.</td><td style="width: 10px;"></td></tr> <tr><td colspan="2" style="border-top: 1px solid black;"></td></tr> <tr><td style="text-align: right;">8.4 mean of the extremes.</td><td style="width: 10px;"></td></tr> <tr><td style="text-align: right;">7.4</td><td style="width: 10px;"></td></tr> <tr><td style="text-align: right;">9.2</td><td style="width: 10px;"></td></tr> <tr><td style="text-align: right;">10.2</td><td style="width: 10px;"></td></tr> <tr><td colspan="2" style="border-top: 1px solid black;"></td></tr> <tr><td style="text-align: right;">35.2 sum.</td><td style="width: 10px;"></td></tr> </table>	8.2		8.6				2) 16.8 sum of the extremes.				8.4 mean of the extremes.		7.4		9.2		10.2				35.2 sum.		<table style="width: 100%; border-collapse: collapse;"> <tr><td style="text-align: right;">35.2 sum.</td><td style="width: 10px;"></td></tr> <tr><td style="text-align: right;">39</td><td style="width: 10px;"></td></tr> <tr><td colspan="2" style="border-top: 1px solid black;"></td></tr> <tr><td style="text-align: right;">3168</td><td style="width: 10px;"></td></tr> <tr><td style="text-align: right;">1056</td><td style="width: 10px;"></td></tr> <tr><td colspan="2" style="border-top: 1px solid black;"></td></tr> <tr><td style="text-align: right;">4) 1372.8</td><td style="width: 10px;"></td></tr> <tr><td colspan="2" style="border-top: 1px solid black;"></td></tr> <tr><td style="text-align: right;">343.2 the area.</td><td style="width: 10px;"></td></tr> <tr><td colspan="2" style="border-top: 1px solid black;"></td></tr> </table>	35.2 sum.		39				3168		1056				4) 1372.8				343.2 the area.			
8.2																																											
8.6																																											
2) 16.8 sum of the extremes.																																											
8.4 mean of the extremes.																																											
7.4																																											
9.2																																											
10.2																																											
35.2 sum.																																											
35.2 sum.																																											
39																																											
3168																																											
1056																																											
4) 1372.8																																											
343.2 the area.																																											

Ex.

* This rule is made out as follows:—Let ABCD be the irregular piece; having the several breadths AD, EF, GH, IK, BC, at the equal distances AE, EG, GI, IB. Let the several breadths in order be denoted by the corresponding letters a, b, c, d, e, and the whole length AB by l; then compute the areas of



the parts into which the figure is divided by the perpendiculars, as so many trapezoids, by prob. 3, and add them all together. Thus, the sum of the parts is,

$$\frac{a+b}{2} \times AE + \frac{b+c}{2} \times EG + \frac{c+d}{2} \times GI + \frac{d+e}{2} \times IB = \frac{a+b}{2} \times \frac{1}{4}l + \frac{b+c}{2} \times \frac{1}{4}l + \frac{c+d}{2} \times \frac{1}{4}l + \frac{d+e}{2} \times \frac{1}{4}l = (\frac{1}{2}a + b + c + d + \frac{1}{2}e) \times \frac{1}{4}l = (m + b + c + d) \frac{1}{4}l$$

which is the whole area, agreeing with the rule: m being

Ex. 2. The length of an irregular figure being 34, and the breadths at six equidistant places 17·4, 20·6, 14 2, 16·5, 20·1, 24·4; what is the area? Ans. 1550·64.

PROBLEM XIV.

To find the Area of an Ellipsis or Oval.

MULTIPLY the longest diameter, or axis by the shortest; then multiply the product by the decimal ·7854, for the area. As appears from cor. 2, theor. 3, of the Ellipse, in the Conic Sections.

Ex. 1. Required the area of an ellipse whose two axes are 70 and 50. Ans. 2748·9.

Ex. 2. To find the area of the oval whose two axes are 24 and 18. Ans. 339·2928.

PROBLEM XV.

To find the Area of any Elliptic Segment.

FIND the area of a corresponding circular segment, having the same height and the same vertical axis or diameter. Then say, as the said vertical axis is to the other axis, parallel to the segment's base, so is the area of the circular segment before found, to the area of the elliptic segment sought. This rule also comes from cor. 2, theor 3 of the Ellipse.

Otherwise thus. Divide the height of the segment by the vertical axis of the ellipse; and find, in the table of circular segments to prob. 12, the circular segment having the above quotient for its versed sine: then multiply all together, this segment and the two axes of the ellipse, for the area.

Ex. 1. To find the area of the elliptic segment, whose height is 20, the vertical axis being 70, and the parallel axis 50.

being the arithmetical mean between the extremes, or half the sum of them both, and 4 the number of the parts. And the same for any other number of parts whatever.

Here

Here $20 \div 70$ gives $28\frac{4}{7}$ the quotient or versed sine ; to which in the table answers the seg. $\cdot 18518$

then 70

12·96260

50

648·13000 the area.

Ex. 2. Required the area of an elliptic segment, cut off parallel to the shorter axis ; the height being 10, and the two axes 25 and 35. Ans. 162·03.

Ex. 3. To find the area of the elliptic segment, cut off parallel to the longer axis ; the height being 5, and the axis 25 and 35. Ans. 97·8425.

PROBLEM XVI.

To find the Area of a Parabola, or its Segment.

MULTIPLY the base by the perpendicular height ; then take two-thirds of the product for the area. As is proved in theorem 17 of the Parabola, in the Conic Sections.

Ex. 1. To find the area of a parabola ; the height being 2, and the base 12.

Here $2 \times 12 = 24$. Then $\frac{2}{3}$ of $24 = 16$, is the area.

Ex. 2. Required the area of the parabola, whose height is 10, and its base 16. Ans. $106\frac{2}{3}$.



MENSURATION OF SOLIDS.

BY the Mensuration of Solids are determined the spaces included by contiguous surfaces ; and the sum of the measures of these including surfaces, is the whole surface or superficies of the body.

The measure of a solid, is called its solidity, capacity, or content.

Solids are measured by cubes, whose sides are inches, or feet, or yards, &c. And hence the solidity of a body is said to be so many cubic inches, feet, yards, &c. as will fill its capacity or space, or another of an equal magnitude.

The

The least solid measure is the cubic inch, other cubes being taken from it according to the proportion in the following table, which is formed by cubing the linear proportions.

Table of Cubic or Solid Measures.

1728	cubic inches make	1 cubic foot
27	cubic feet -	1 cubic yard
$166\frac{2}{3}$	cubic yards -	1 cubic pole
64000	cubic poles -	1 cubic furlong
512	cubic furlongs -	1 cubic mile

PROBLEM I.

To find the Superficies of a Prism or Cylinder.

MULTIPLY the perimeter of one end of the prism by the length or height of the solid, and the product will be the surface of all its sides. To which add also the area of the two ends of the prism, when required *

Or, compute the areas of all the sides and ends separately, and add them all together.

Ex. 1. To find the surface of a cube, the length of each side being 20 feet. Ans. 2400 feet.

Ex. 2. To find the whole surface of a triangular prism, whose length is 20 feet, and each side of its end or base 13 inches. Ans. 91.948 feet.

Ex. 3. To find the convex surface of a round prism, or cylinder, whose length is 20 feet, and the diameter of its base is 2 feet. Ans. 125.664.

Ex. 4. What must be paid for lining a rectangular cistern with lead, at 2*d.* a pound weight, the thickness of the lead being such as to weigh 7*lb.* for each square foot of surface; the inside dimensions of the cistern being as follow, viz. the length 3 feet 2 inches, the breadth 2 feet 8 inches, and depth 2 feet 6 inches? Ans. 2*l.* 3*s.* 10 $\frac{1}{2}$ *d.*

* The truth of this will easily appear, by considering that the sides of any prism are parallelograms, whose common length is the same as the length of the solid, and their breadths taken all together make up the perimeter of the ends of the same.

And the rule is evidently the same for the surface of a cylinder.

PROBLEM

Note. For a cube, take the cube of its side by multiplying this twice by itself; and for a parallelepipedon, multiply the length, breadth, and depth all together, for the content.

Ex. 1. To find the solid content of a cube, whose side is 24 inches. Ans. 13824.

Ex. 2: How many cubic feet are in a block of marble, its length being 3 feet 2 inches, breadth 2 feet 8 inches, and thickness 2 feet 6 inches? Ans. $21\frac{1}{8}$.


Ex. 3. How many gallons of water will the cistern contain, whose dimensions are the same as in the last example, when 282 cubic inches are contained in one gallon? Ans. $129\frac{1}{4}$.

Ex. 4. Required the solidity of a triangular prism, whose length is 10 feet, and the three sides of its triangular end or base are 3. 4. 5 feet. Ans. 60.

Ex. 5. Required the content of a round pillar, or cylinder, whose length is 20 feet, and circumference 5 feet 6 inches. Ans. $48\cdot1459$ feet.

the solid to be measured, and the cube p the solid measuring unit, its side being 1 inch, or 1 foot, &c. : also, let the length and breadth of the base, $ABCD$ and also the height AH , be each divided into spaces equal to the length of the base of the cube p , namely, here 3 in the length and 2 in the breadth, making 3 times 3 or 6 squares in the base AC , each equal to the base of the cube p . Hence it is manifest that the parallelepipedon will contain the cube p , as many times as the base AC contains the base of the cube, repeated as often as the height AH contains the height of the cube. That is, the content of any parallelepipedon is found, by multiplying the area of the base by the altitude of that solid.

And, because all prisms and cylinders are equal to parallelepipedons of equal bases and altitudes, by Geom. theor. 108, it follows that the rule is general for all such solids, whatever the figure of the base may be.



PROBLEM V.

To find the Content of any Pyramid or Cone.

FIND the area of the base, and multiply that area by the perpendicular height; then take $\frac{1}{3}$ of the product for the content.*

Ex. 1. Required the solidity of the square pyramid, each side of its base being 30, and its perpendicular height 25.

Ans. 7500.

Ex. 2. To find the content of a triangular pyramid whose perpendicular height is 30, and each side of the base 3.

Ans. 38 97117.

Ex. 3. To find the content of a triangular pyramid, its height being 14 feet 6 inches, and the three sides of its base 5, 6, 7 feet.

Ans. 71.0352.

Ex. 4. What is the content of a pentagonal pyramid its height being 12 feet, and each side of its base 2 feet?

Ans. 27.5276.

Ex. 5. What is the content of the hexagonal pyramid, whose height is 6.4 feet, and each side of its base 6 inches?

Ans. 1.38564 feet.

Ex. 6. Required the content of a cone, its height being $10\frac{1}{2}$ feet and the circumference of its base 9 feet.

Ans. 22.56093.

PROBLEM VI.

To find the Solidity of the Frustrum of a Cone or Pyramid.

ADD into one sum, the areas of the two ends, and the mean proportional between them: and take $\frac{1}{3}$ of that sum for a mean area; which being multiplied by the perpendicular height or length of the frustrum will give its content †

Note.

* This rule follows from that of the prism, because any pyramid is $\frac{1}{3}$ of a prism of equal base and altitude; by Geom. theor. 115, cor. 1 and 2.

† Let ABCD be any pyramid, of which BCDGFE is a frustrum. And put a^2 for the area of the base BCD, b^2 the area of the top EFG, h the height HH of the frustrum.


Note. This general rule may be otherwise expressed, as follows, when the ends of the frustum are circles or regular polygons. In this latter case, square one side of each polygon, and also multiply the one side by the other; add all these three products together; then multiply their sum by the tabular area proper to the polygon, and take one-third of the product for the mean area, to be multiplied by the length, to give the solid content. And in the case of the frustum of a cone, the ends being circles, square the diameter or the circumference of each end, and also multiply the same two dimensions together; then take the sum of the three products and multiply it by the proper tabular number, viz. by $\cdot 7854$ when the diameters are used, or by $\cdot 07958$ in using the circumferences; then taking one-third of the product to multiply by the length, for the content.

Ex. 1. To find the number of solid feet in a piece of timber, whose bases are squares, each side of the greater end being 15 inches, and each side of the less end 6 inches; also, the length or perpendicular altitude 24 feet. Ans. $19\frac{1}{2}$.

Ex. 2 Required the content of a pentagonal frustum, whose height is 5 feet, each side of the base 18 inches, and each side of the top or less end 6 inches. Ans. $9\cdot 31925$ feet.

tum, and c the height AI of the top part above it. Then $c + h = AH$ is the height of the whole pyramid.

Hence, by the last prob. $\frac{1}{3} a^2 (c + h)$ is the content of the whole pyramid $ABCD$, and $\frac{1}{3} b^2 c$ the content of the top part $AIEFG$; therefore the difference $\frac{1}{3} a^2 (c + h) - \frac{1}{3} b^2 c$ is the content of the frustum $BCDGFIE$. But the quantity c being no dimension of the frustum, it must be expelled from this formula, by substituting its value, found in the following manner. By Geom. theor. 112, $a^2 : b^2 :: (c + h)^2 : c^2$, or $a : b :: c + h : c$, hence (Geom. th. 69) $a - b : b :: h : c$, and $a - b : a :: b : c + h$; hence therefore $c = \frac{bh}{a - b}$ and $c + h = \frac{ah}{a - b}$; then these values of c and $c + h$ being substituted for them in the expression for the content of the frustum, gives that content $= \frac{1}{3} a^2 \times \frac{ah}{a - b} - \frac{1}{3} b^2 \times \frac{bh}{a - b} = \frac{1}{3} h \times \frac{a^3 - b^3}{a - b} = \frac{1}{3} h \times (a^2 + ab + b^2)$; which is the rule above given; ab being the mean between a^2 and b^2 .



Ex. 3.

Ex. 3. To find the content of a conic frustum, the altitude being 18, the greatest diameter 8, and the least diameter 4.
 Ans. 527·7888.

Ex. 4. What is the solidity of the frustum of a cone, the altitude being 25, also the circumference at the greater end being 20, and at the less end 10?
 Ans. 464·216.

Ex. 5. If a cask, which is two equal conic frustums joined together at the bases, have its bung diameter 28 inches, the head diameter 20 inches, and length 40 inches; how many gallons of wine will it hold.
 Ans. 79·0613.

PROBLEM VII.


To find the Surface of a Sphere, or any Segment.

RULE 1. MULTIPLY the circumference of the sphere by its diameter, and the product will be the whole surface of it.*

RULE II.

* These rules come from the following theorems, for the surface of a sphere, viz. That the said surface is equal to the curve surface of its circumscribing cylinder; or that it is equal to 4 great circles of the same sphere, or of the same diameter; which are thus proved.

Let ABCD be a cylinder, circumscribing the sphere EFGH; the former generated by the rotation of the rectangle FBCH about the axis or diameter FH; and the latter by the rotation of the semicircle FGH about the same diameter FH. Draw two lines KL, MN, perpendicular to the axis intercepting the parts LN, OP, of the cylinder and sphere; then will the ring or cylindric surface generated by the rotation of LN, be equal to the ring or spherical surface generated by the arc OP. For first, suppose the parallels KL and MN to be indefinitely near together; drawing IO, and also OQ parallel to LN. Then, the two triangles IKO, OQP, being equiangular, it is, as OP : OQ or LN :: IO or KL : KO :: circumference described by KL : circumf. described by KO; therefore the rectangle OP x circumf. of KO is equal to the rectangle LN x circumf. of KL; that is, the ring described by OP on the sphere, is equal to the ring described by LN on the cylinder.



And

RULE II. Square the diameter and multiply that square by 3·1416, for the surface.

Rule III. Square the circumference ; then either multiply that square by the decimal ·3183, or divide it by 3·1416, for the surface.

Note. For the surface of a segment or frustum, multiply the whole circumference of the sphere by the height of the part required.

Ex. 1. Required the convex superficies of a sphere, whose diameter is 7, and circumference 22 Ans. 154.

Ex. 2. Required the superficies of a globe, whose diameter is 24 inches. Ans. 1809·5616.

Ex. 3. Required the area of the whole surface of the earth, its diameter being 7918·7 miles. Ans. 196994111 sq. miles.

Ex. 4. The axis of a sphere being 42 inches, what is the convex superficies of the segment whose height is 9 inches ? Ans. 1187·5248 inches.

Ex. 5. Required the convex surface of a spherical zone, whose breadth or height is 2 feet, and cut from a sphere of $12\frac{1}{2}$ feet diameter. Ans. 78·54 feet.

And as this is every where the case, therefore the sums of any corresponding number of these are also equal ; that is, the whole surface of the sphere, described by the whole semicircle FGH , is equal to the whole curve surface of the cylinder, described by the height bc ; as well as the surface of any segment described by fo , equal to the surface of the corresponding segment described by bl .

Corol. 1. Hence the surface of the sphere is equal to 4 of its great circles, or equal to the circumference $efgh$, or of dc , multiplied by the height bc , or by the diameter fh .

Corol. 2. Hence also, the surface of any such part as a segment or frustum, or zone, is equal to the same circumference of the sphere, multiplied by the height of the said part. And consequently such spherical curve surfaces are to one another in the same proportion as their altitudes.

PROBLEM VIII.

To find the Solidity of a Sphere or Globe.

RULE I. Multiply the surface by the diameter, and take $\frac{1}{6}$ of the product for the content.* Or, which is the same thing, multiply the square of the diameter, by the circumference, and take $\frac{1}{6}$ of the product.

RULE II. Take the cube of the diameter, and multiply it by the decimal .5236, for the content.

RULE III. Cube the circumference, and multiply by .01688.

Ex. 1. To find the content of a sphere whose axis is 12.
 Ans. 904.7808.

Ex. 2. To find the solid content of the globe of the earth supposing its diameter to be 7918.7, and consequently its circumference 24877.4 miles.
 Ans. 260002677535 miles.

PROBLEM IX.

To find the Solid Content of a Spherical Segment.

† **RULE I.** From 3 times the diameter of the sphere take double


* For, put $d =$ the diameter, $c =$ the circumference, and $s =$ the surface of the sphere, or of its circumscribing cylinder: also, $a =$ the number 3.1416.

Then, $\frac{1}{4}s$ is = the base of the cylinder, or one great circle of the sphere; and d is the height of the cylinder; therefore $\frac{1}{4}ds$ is the content of the cylinder. But $\frac{2}{3}$ of the cylinder is the sphere, by th. 117, Geom. that is, $\frac{2}{3}$ of $\frac{1}{4}ds$, or $\frac{1}{6}ds$ is the sphere; which is the first rule.

Again, because the surface s is = ad^2 ; therefore $\frac{1}{6}ds = \frac{1}{6}ad^3 = .5236d^3$, is the content, as in the 2d rule. Also d being = $c \div a$ therefore $\frac{1}{6}ad^3 = \frac{1}{6}c^3 \div a^2 = .01688$, the 3d rule for the content.

† By corol. 3, of theor. 117, Geom. it appears that the spheric segment PFN, is equal to the difference between the cylinder ABLO, and the conic frustum ABMQ.

But, putting $d = AB$ or FH the diameter of the sphere or cylinder, $h = FK$ the height of the segment, $r = PK$ the radius of its base, and $a = 3.1416$; then the content of the cone ABI is = $\frac{1}{4}ad^2 \times \frac{1}{3}FI = \frac{1}{12}ad^2$; and by the similar cones ABI, QMI, as $FI^3 : KI^3 :: \frac{1}{12}ad^3 : \frac{1}{12}ad^3 \times$



double the height of the segment : then multiply the remainder by the square of the height, and the product by the decimal .5236, for the content.

RULE II. To 3 times the square of the radius of the segment's base, add the square of its height ; then multiply the sum by the height, and the product by .5236, for the content.

Ex. 1. To find the content of a spherical segment, of 2 feet in height, cut from a sphere of 4 feet diameter.

Ans. 41.888.

Ex. 2. What is the solidity of the segment of a sphere, its height being 9, and the diameter of its base 20 ?

Ans. 1795.4244.

Note. The general rules for measuring all sorts of figures having been now delivered, we may next proceed to apply them to the several practical uses in life, as follows.

$(\frac{\frac{1}{2}d-h}{\frac{1}{2}d})^3$ = the cone QMI ; therefore the cone ABI — the cone QMI = $\frac{1}{24}ad^3$
 — $\frac{1}{24}ad^3 \times (\frac{\frac{1}{2}d-h}{\frac{1}{2}d})^3$ = $\frac{1}{4}ad^2 h - \frac{1}{2}adh^2 + \frac{1}{3}ah^3$ is = the conic frustum
 ABMQ.

And $\frac{1}{4}ad^2 h$ is = the cylinder ABLO.

Then the difference of these two is $\frac{1}{2}adh^2 - \frac{1}{3}ah^3 = \frac{1}{6}ah^2 \times (3d - 2h)$, for the spheric segment PFN ; which is the first rule.

Again, because $PK^2 = FK \times KH$ (COR. to theor. 87, Geom.) or $r^2 = h(d-h)$, therefore $d = \frac{r^2}{h} + h$, and $3d - 2h = \frac{3r^2}{h} + h = \frac{3r^2 + h^2}{h}$; which being substituted in the former rule, it becomes $\frac{1}{6}ah^2 \times \frac{3r^2 + h^2}{h} = \frac{1}{6}ah^2 \times (3r^2 + h^2)$, which is the 2d rule.

Note. By subtracting a segment from half a sphere, or from another segment, the content of any frustum or zone may be found.

LAND SURVEYING.

SECTION I.

DESCRIPTION AND USE OF THE INSTRUMENTS.

1. OF THE CHAIN.

LAND is measured with a chain, called Gunter's Chain, from its inventor, the length of which is 4 poles, or 22 yards, or 66 feet. It consists of 100 equal links; and the length of each link is therefore $\frac{22}{100}$ of a yard, or $\frac{66}{100}$ of a foot, or 7.92 inches.

Land is estimated in acres, roods, and perches. An acre is equal to 10 square chains, or as much as 10 chains in length and 1 chain in breadth. Or, in yards, it is $22 \times 22 = 4840$ square yards. Or, in poles, it is $40 \times 4 = 160$ square poles. Or, in links, it is $1000 \times 100 = 100000$ square links: these being all the same quantity.

Also, an acre is divided into 4 parts called roods, and a rood into 40 parts called perches, which are square poles, or the square of a pole of $5\frac{1}{2}$ yards long, or the square of $\frac{1}{4}$ of a chain, or of 25 links, which is 625 square links. So that the divisions of land measure, will be thus:

$$\begin{aligned} 625 \text{ sq. links} &= 1 \text{ pole or perch} \\ 40 \text{ perches} &= 1 \text{ rood} \\ 4 \text{ roods} &= 1 \text{ acre.} \end{aligned}$$

The length of lines, measured with a chain, are best set down in links as integers, every chain in length being 100 links; and not in chains and decimals. Therefore, after the content is found, it will be in square links; then cut off five of the figures on the right hand for decimals, and the rest will be acres. These decimals are then multiplied by 4 for roods, and the decimals of these again by 40 for perches.

EXAM. Suppose the length of a rectangular piece of ground be 792 links, and its breadth 385; to find the area in acres, roods, and perches.

792	3.04920
385	4
3960	.19680
6336	40
2376	7.87200
3.04920	3.04920

Ans. 3 acres, 0 roods, 7 perches.

2. OF THE PLAIN TABLE.

THIS instrument consists of a plane rectangular board, of any convenient size : the centre of which, when used, is fixed by means of screws to a three-legged stand, having a ball and socket, or other joint, at the top, by means of which, when the legs are fixed on the ground, the table is inclined in any direction.

To the table belong various parts, as follow :

1. A frame of wood, made to fit round its edges, and to be taken off, for the convenience of putting a sheet of paper on the table. One side of this frame is usually divided into equal parts, for drawing lines across the table, parallel or perpendicular to the sides ; and the other side of the frame is divided into 360 degrees, to a centre in the middle of the table ; by means of which the table may be used as a theodolite, &c.

2. A magnetic needle and compass, either screwed into the side of the table, or fixed beneath its centre, to point out the directions, and to be a check on the sights.

3. An index, which is a brass two-foot scale, with either a small telescope, or open sights set perpendicularly on the ends. These sights and one edge of the index are in the same plane, and that is called the fiducial edge of the index.

To use this instrument, take a sheet of paper which will cover it, and wet it to make it expand ; then spread it flat on the table, pressing down the frame on the edges, to stretch it and keep it fixed there ; and when the paper is become dry, it will, by contracting again, stretch itself smooth and flat from any cramps and unevenness. On this paper is to be drawn the plan or form of the thing measured.

Thus, begin at any proper part of the ground, and make a point on a convenient part of the paper or table, to represent that place on the ground ; then fix in that point one leg of the compasses, or a fine steel pin, and apply to it the fiducial edge of the index, moving it round till through the sights you perceive some remarkable object, as the corner of a field, &c. ; and from the station-point draw a line with the point of the compasses along the fiducial edge of the index, which is called setting or taking the object : then set another object or corner, and draw its line ; do the same by another ; and so on, till as many objects are taken as may be thought fit. Then measure from the station towards as many of the objects as may be necessary, but not more, taking the requisite offsets to corners or crooks in the hedges, laying the measures down on their respective lines on the table.

Then

Then at any convenient place measured to, fix the table in the same position, and set the objects which appear from that place : and so on, as before. And thus continue till the work is finished, measuring such lines only as are necessary, and determining as many as may be by intersecting lines of direction drawn from different stations.

Of shifting the Paper on the Plain Table.

When one paper is full, and there is occasion for more ; draw a line in any manner through the farthest point of the last station line, to which the work can be conveniently laid down ; then take the sheet off the table, and fix another on, drawing a line over it, in a part the most convenient for the rest of the work ; then fold or cut the old sheet by the line drawn on it, applying the edge to the line on the new sheet, and, as they lie in that position, continue the last station line on the new paper, placing on it the rest of the measure, beginning at where the old sheet left off. And so on from sheet to sheet.

When the work is done, and you would fasten all the sheets together into one piece, or rough plan, the aforesaid lines are to be accurately joined together, in the same manner as when the lines were transferred from the old sheets to the new ones. But it is to be noted, that if the said joining lines, on the old and new sheets, have not the same inclination to the side of the table, the needle will not point to the original degree when the table is rectified ; and if the needle be required to respect still the same degree of the compass, the easiest way of drawing the lines in the same position, is to draw them both parallel to the same sides of the table, by means of the equal divisions marked on the other two sides.

3. OF THE THEODOLITE.

THE theodolite is a brazen circular ring, divided into 360 degrees, &c. and having an index with sights, or a telescope, placed on the centre, about which the index is moveable ; also a compass fixed to the centre, to point out courses and check the sights ; the whole being fixed by the centre on a stand of a convenient height for use.

In using this instrument, an exact account, or field-book, of all measures and things necessary to be remarked in the plan, must be kept, from which to make out the plan on returning home from the ground.

Begin at such part of the ground, and measure in such directions as are judged most convenient ; taking angles or directions to objects, and measuring such distances as appear necessary,

necessary, under the same restrictions as in the use of the plain table. And it is safest to fix the theodolite in the original position at every station, by means of fore and back objects, and the compass, exactly as in using the plain table; registering the number of degrees cut off by the index when directed to each object; and, at any station placing the index at the same degree as when the direction towards that station was taken from the last preceding one, to fix the theodolite there in the original position.

The best method of laying down the aforesaid lines of direction, is to describe a pretty large circle; then quarter it, and lay on it the several numbers of degrees cut off by the index in each direction, and drawing lines from the centre to all these marked points in the circle. Then, by means of a parallel ruler draw from station to station, lines parallel to the aforesaid lines drawn from the centre to the respective points in the circumference.

4. OF THE CROSS.

The cross consists of two pair of sights set at right angles to each other, on a staff having a sharp point at the bottom, to fix in the ground.

The cross is very useful to measure small and crooked pieces of ground. The method is, to measure a base or chief line, usually in the longest direction of the piece, from corner to corner; and while measuring it, finding the places where perpendiculars would fall on this line, from the several corners and bends in the boundary of the piece, with the cross, by fixing it, by trials, on such parts of the line, as that through one pair of the sight both ends of the line may appear, and through the other pair the corresponding bends or corners; and then measuring the lengths of the said perpendiculars.

REMARKS.

Besides the fore-mentioned instruments, which are most commonly used, there are some others; as,

The perambulator, used for measuring roads, and other great distances, level ground and by the sides of rivers. It has a wheel of $8\frac{1}{4}$ feet, or half a pole, in circumference, by the turning of which the machine goes forward: and the distance measured is pointed out by an index, which is moved round by clock work.

Levels, with telescopic or other sights, are used to find the level between place and place, or how much one place is higher or lower than another. And in measuring any sloping or oblique line, either ascending or descending, a small pocket

pocket level is useful for showing how many links for each chain are to be deducted, to reduce the line to the horizontal length.

An offset-staff is a very useful instrument, for measuring the offsets and other short distances. It is 10 links in length, being divided and marked at each of the 10 links.

Ten small arrows, or rods of iron, or wood, are used to mark the end of every chain length, in measuring lines. And sometimes pickets, or staves with flags, are set up as marks or objects of direction.

Various scales are also used in protracting and measuring on the plan or paper; such as plane scales, line or chords, protractor, compasses, reducing scale, parallel and perpendicular rules, &c. Of plane scales there should be several sizes, as a chain in 1 inch, a chain in $\frac{3}{4}$ of an inch, a chain in $\frac{1}{2}$ an inch, &c. And of these, the best for use are those that are laid on the very edges of the ivory scale, to mark off distances, without compasses.

SECTION II.

THE PRACTICE OF SURVEYING.

THIS part contains the several works proper to be done in the field, or the ways of measuring by all the instruments, and in all situations.

PROBLEM I.

To Measure a Line or Distance.

To measure a line on the ground with the chain: Having provided a chain, with 10 small arrows, or rods, to fix one into the ground, as a mark, at the end of every chain; two persons take hold of the chain, one at each end of it; and all the 10 arrows are taken by one of them, who goes foremost, and is called the leader; the other being called the follower, for distinction's sake.

A picket, or station-staff being set up in the direction of the line to be measured, if there do not appear some marks naturally in that direction, they measure straight towards it, the leader fixing down an arrow at the end of every chain, which the follower always takes up, as he comes at it, till all the ten arrows are used. They are then all returned to the leader, to use over again. And thus the arrows are changed from the one to the other at every 10 chains' length, till the whole line is finished; then the number of changes


of the arrows shows the number of tens, to which the follower adds the arrows he holds in his hand, and the number of links of another chain over to the mark or end of the line. So, if there have been 3 changes of the arrows, and the follower hold 6 arrows and the end of the line cut off 45 links more, the whole length of the line is set down in links thus, 3645.

When the ground is not level, but either ascending or descending; at every chain length, lay the offset-staff, or link-staff, down in the slope of the chain, on which lay the small pocket level, to show how many links or parts the slope line is longer than the true level one; then draw the chain forward so many links or parts, which reduces the line to the horizontal direction.

PROBLEM II.

To take Angles and Bearings.

LET B and c be two objects, or two pickets set up perpendicular; and let it be required to take their bearings, or the angles formed between them at any station A.



1. *With the Plain Table.*

The table being covered with a paper, and fixed on its stand; plant it at the station A, and fix a fine pin, or a foot of the compasses, in a proper point of the paper, to represent the place A: Close by the side of this pin lay the fiducial edge of the index, and turn it about, still touching the pin, till one object B can be seen through the sights: then by the fiducial edge of the index draw a line. In the same manner draw another line in the direction of the other object c. And it is done.

2. *With the Theodolite, &c.*

Direct the fixed sights along one of the lines, as AB, by turning the instrument about till the mark B is seen through these sights; and there screw the instrument fast. Then turn the moveable index round, till through its sights the other mark c is seen. Then the degrees cut by the index, on the graduated limb or ring of the instrument, show the quantity of the angle.

3. *With*

3. *With the Magnetic Needle and Compass.*

Turn the instrument or compass so, that the north end of the needle point to the flower-de-luce. Then direct the sights to one mark as B, and note the degrees cut by the needle. Next direct the sights to the other mark c, and note again the degrees cut by the needle. Then their sum or difference, as the case may be, will give the quantity of the angle BAC.

4. *By Measurement with the Chain, &c.*


Measure one chain length, or any other length, along both directions, as to b and c. Then measure the distance bc, and it is done.—This is easily transferred to paper, by making a triangle abc with these three lengths, and then measuring the angle a.

PROBLEM III.

To Survey a Triangular Field ABC.

1. *By the Chain.*

AP 794
 AB 1321
 PC 826



Having set up marks at the corners, which is to be done in all cases where there are not marks naturally; measure with the chain from A to P, where a perpendicular would fall from the angle c, and set up a mark at P, noting down the distance AP. Then complete the distance AB, by measuring from P to B. Having set down this measure, return to P, and measure the perpendicular PC. And thus, having the base and perpendicular, the area from them is easily found. Or having the place P of the perpendicular, the triangle is easily constructed.

Or, measure all the three sides with the chain, and note them down. From which the content is easily found, or the figure is constructed.

2. *By taking some of the Angles.*

Measure two sides AB, AC, and the angle A between them. Or measure one side AB, and the two adjacent angles A and B. From either of these ways the figure is easily planned; then by measuring the perpendicular CP on the plan, and multiplying it by half AB, the content is found.


PROBLEM

PROBLEM IV.

To Measure a Four-sided Field.

1. By the Chain.


AE	214		210	DE
AF	362		306	BF
AC	592			



Measure along one of the diagonals, as AC; and either the two perpendiculars DE, BF, as in the last problem; or else the sides AB, BC, CD, DA. From either of which the figure may be planned and computed as before directed.

Otherwise by the Chain.

AP	110		352	PC
AQ	745		595	QD
AB	1110			



Measure on the longest side, the distances AP, AQ, AB; and the perpendiculars PC, QD.

2. By taking some of the Angles.

Measure the diagonal AC (see the last fig. but one), and the angles CAB, CAD, ACB, ACD.—Or measure the four sides, and any one of the angles, as BAD.

Thus.			Or thus.	
AC	591		AB	486
CAB	37° 20'		BC	394
CAD	41 15		CD	410
ACB	72 25		DA	462
ACD	54 40		BAD	78° 35'


PROBLEM V.

To Survey any Field by the Chain only.

HAVING set up marks at the corners, where necessary, of the proposed field ABCDEFG, walk over the ground, and consider how it can best be divided in triangles and trapeziums; and measure them separately, as in the last two problems. Thus, the following figure is divided into the two trapeziums ABCG, GDEF, and the triangle GCD. Then, in the first trapezium, beginning at A, measure the diagonal AC, and the two

two perpendiculars gm, bn. Then the base gc, and the perpendicular bq. Lastly, the diagonal df, and the two perpendiculars pe, og. All which measures write against the corresponding parts of a rough figure drawn to resemble the figure surveyed, or set them down in any other form you choose.

				Thus;			
am	135	150	mg				
an	410	180	nb				
ac	550						
<hr/>							
cq	152	230	qd				
cg	440						
<hr/>							
fo	237	120	og				
fp	288	80	pe				
fd	520						



Or thus

Measure all the sides AB, BC, CD, DE, EF, FG, GA ; and the diagonals AC, CG, GD, DF.


Otherwise.

Many pieces of land may be very well surveyed, by measuring any base line, either within or without them, with the perpendiculars let fall on it from every corner. For they are by those means divided into several triangles and trapezoids, all whose parallel sides are perpendicular to the base line ; and the sum of these triangles and trapeziums will be equal to the figure proposed if the base line fall within it ; if not, the sum of the parts which are without being taken from the sum of the whole which are both within and without, will leave the area of the figure proposed.

In pieces that are not very large, it will be sufficiently exact to find the points, in the base line, where the several perpendiculars will fall, by means of the cross, or even by judging by the eye only, and from thence measuring to the corners for the lengths of the perpendiculars.—And it will be most convenient to draw the line so as that all the perpendiculars may fall within the figure.

Thus. in the following figure, beginning at A, and measuring along the line AG, the distances and perpendiculars on the right and left are as below.


Ab	315	350	bB
Ac	440	70	cc
Ad	585	320	dd
Ae	610	50	eE
Af	990	470	fF
AG	1020	0	



PROBLEM VI.
To Measure the Offsets.


Ahikman being a crooked hedge, or brook, &c. From A measure in a straight direction along the side of it to B. And in measuring along this line AB, observe when you are directly opposite any bends or corners of the boundary, as at c, d, e, &c. ; and from these measure the perpendicular offsets ch, di, &c. with the offset-staff, if they are not very large, otherwise with the chain itself ; and the work is done. The register, or field-book, may be as follows :

Offs. left.	Base line AB	
	0	⊙ A
ch	62	45 AC
di	84	220 Ad
ek	70	340 Ae
fl	98	510 Af
gm	57	634 Ag
Bn	91	785 AB



PROBLEM VII.
To survey any Field with the Plain Table.
1. *From one Station.*


PLANT the table at any angle as c, from which all the other angles, or marks set up, can be seen ; turn the table about till the needle point to the flower-de-luce ; and there screw it fast. Make a point for c on the paper on the table, and lay the edge of the index to c, turning it about c till through the sights you see the mark D : and by the edge of the index draw a dry or obscure line : then measure the distance cD, and lay that distance down on the line cD. Then turn the index about the point d, till the mark E be seen through the sights,



sights, by which draw a line, and measure the distance to E, laying it on the line from c to E. In like manner determine the positions of CA and CB, by turning the sights successively to A and B; and lay the lengths of those lines down. Then connect the points, by drawing the black lines CD, DE, EA, AB, BC, for the boundaries of the field.

2. *From a Station Within the Field.*

When all the other parts cannot be seen from one angle, choose some place O within, or even without, if more convenient, from which the other parts can be seen. Plant the table at O, then fix it with the needle north, and mark the point O on it. Apply the index successively to O, turning it round with the sights to each angle, A, B, C, D, E, drawing dry lines to them by the edge of the index; then measuring the distance OA, OB, &c. and laying them down on those lines. Lastly, draw the boundaries AB, BC, CD, DE, EA.



3. *By going Round the Figure.*

When the figure is a wood, or water, or when from some other obstruction you cannot measure lines across it; begin at any point A, and measure around it either within or without the figure, and draw the directions of all the sides, thus: Plant the table at A; turn it with the needle to the north or flower-de-luce; fix it, and mark the point A. Apply the index to A, turning it till you can see the point E, and there draw a line: then the point B, and there draw a line: then measure these lines, and lay them down from A to E and B. Next move the table to B, lay the index along the line AB, and turn the table about till you can see the mark A, and screw fast the table; in which position also the needle will again point to the flower-de-luce, as it will do indeed at every station when the table is in the right position. Here turn the index about B till through the sights you see the mark c; there draw a line, measure BC, and lay the distance on that line after you have set down the table at c. Turn it then again into its proper position, and in like manner find the next line CD. And so on quite around by E, to A again. Then the proof of the work will be the joining at A: for if the work be all right, the last direction EA on the ground, will pass exactly through the point A on the paper; and the measured distance will also reach exactly to A. If these do not coincide, or nearly so, some error has been committed, and the work must be examined over again.

PROBLEM VIII.

To Survey a Field with the Theodolite, &c.

1. *From One Point or Station.*


WHEN all the angles can be seen from one point, as the angle c (first fig. to last prob.) place the instrument at c , and turn it about, till through the fixed sights you see the mark B , and there fix it. Then turn the moveable index about till the mark A be seen through the sights, and note the degrees cut on the instrument. Next turn the index successively to E and D , noting the degrees cut off at each; which gives all the angles BCA , BCE , BCD . Lastly measure the lines CB , CA , CE , CD ; and enter the measures in a field-book, or rather against the corresponding parts of a rough figure drawn by guess to resemble the field.

1. *From a point Within or Without.*

Plant the instrument at O (last fig.) and turn it about till the fixed sights point to any object, as A ; and there screw it fast. Then turn the moveable index round till the sights point successively to the other points E , D , C , B , noting the degrees cut off at each of them; which gives all the angles round the point O . Lastly measure the distances OA , OB , OC , OD , OE , noting them down as before, and the work is done.

3. *By going Round the Field.*

By measuring round, either within or without the field, proceed thus. Having set up marks at B , c , &c. near the corners as usual, plant the instrument at any point A , and turn it till the fixed index be in the direction AB , and there screw it fast: then turn the moveable index to the direction AF ; and the degrees



cut off will be the angle A . Measure the line AB , and plant the instrument at B , and there in the same manner observe the angle c . Then measure BC , and observe the angle C . Then measure the distance CD , and take the angle D . Then measure DE , and take the angle E . Then measure EF , and take the angle F . And lastly measure the distance FA .

To prove the work; add all the inward angles A , B , C , &c. together; for when the work is right, their sum will be equal to twice as many right angles as the figure has sides, wanting 4 right angles. But when there is an angle, as F , that bends inwards, and you measure the external angle, which

which is less than two right angles, subtract it from 4 right angles, or 360 degrees, to give the internal angle greater than a semicircle or 180 degrees.


Otherwise.

Instead of observing the internal angles, we may take the external angles, formed without the figure by producing the sides farther out. And in this case, when the work is right, their sum altogether will be equal to 360 degrees. But when one of them, as F, runs inwards, subtract it from the sum of the rest, to leave 360 degrees.

PROBLEM IX.


To Survey a Field with Crooked Hedges, &c.

WITH any of the instruments, measure the lengths and positions of imaginary lines running as near the sides of the field as you can; and, in going along them, measure the off-sets in the manner before taught; then you will have the plan on the paper in using the plain table, drawing the crooked hedges through the ends of the off-sets; but in surveying with the theodolite, or other instrument, set down the measures properly in a field-book, or memorandum-book, and plan them after returning from the field, by laying down all the lines and angles.



So, in surveying the piece ABCDE, set up marks a, b, c, d, dividing it so as to have as few sides as may be. Then begin at any station, a, and measure the lines ab, bc, cd, da, taking their positions, or the angles a, b, c, d; and, in going along the lines, measure all the off-sets, as at m, n, o, p, &c. along every station-line.

And this is done either within the field, or without, as may be most convenient. When there are obstructions within, as wood, water, hills, &c. then measure without, as in the next following figure.




PROBLEM X.

To Survey a Field, or any other Thing, by Two Stations.

This is performed by choosing two stations from which all the marks and objects can be seen; then measuring the distance between the stations. and at each station taking the angles formed by every object from the station line or distance.

The two stations may be taken either within the bounds, or in one of the sides, or in the direction of two of the objects, or quite at a distance and without the bounds of the objects or part to be surveyed.

In this manner, not only grounds may be surveyed, without even entering them, but a map may be taken of the principal parts of a county, or the chief places of a town. or any part of a river or coast surveyed, or any other inaccessible objects; by taking two stations, on two towers, or two hills, or such-like.



PROBLEM XI.

To Survey a Large Estate.

If the estate be very large, and contain a great number of fields, it cannot well be done by surveying all the fields singly

singly, and then putting them together ; nor can it be done by taking all the angles and boundaries that enclose it. For in these cases, any small errors will be so much increased, as to render it very much distorted. But proceed as below.

1. Walk over the estate two or three times, in order to get a perfect idea of it, or till you can keep the figure of it pretty well in mind. And to help your memory, draw an eye-draught of it on paper. or at least of the principal parts of it, to guide you ; setting the names within the fields in that draught.

2. Choose two or more eminent places in the estate, for stations, from which all the principal parts of it can be seen : selecting these stations as far distant from one another as convenient.

3. Take such angles, between the stations, as you think necessary, and measure the distances from station to station, always in a right line : these things must be done, till you get as many angles and lines as are sufficient for determining all the points of station. And in measuring any of these station-distances mark accurately where these lines meet with any hedges, ditches, roads, lanes, paths, rivulets, &c. ; and where any remarkable object is placed, by measuring its distance from the station-line ; and where a perpendicular from it cuts that line. And thus as you go along any main station-line, take offsets to the ends of all hedges, and to any pond, house, mill, bridge, &c. noting every thing down that is remarkable.

4. As to the inner parts of the estate, they must be determined, in like manner, by new station-lines : for after the main stations are determined, and every thing adjoining to them, then the estate must be subdivided into two or three parts by new station-lines ; taking inner stations at proper places, where you can have the best view. Measure these station-lines as you did the first, and all their intersections with hedges, and offsets to such objects as appear. Then proceed to survey the adjoining fields, by taking the angles that the sides make with the station-line, at the intersections, and measuring the distances to each corner, from the intersections. For the station-lines will be the bases to all the future operations ; the situation of all parts being entirely dependent on them ; and therefore they should be taken of as great length as possible ; and it is best for them to run along some of the hedges or boundaries of one or more fields, or to pass through some of their angles. All things being determined for these stations, you must take more inner stations, and continue to divide and subdivide till at last you come to single fields ; repeating the same work for the inner
stations

stations as for the outer ones, till all is done; and close the work as often as you can, and in as few lines as possible.

5. An estate may be so situated that the whole cannot be surveyed together; because one part of the estate cannot be seen from another. In this case, you may divide it into three or four parts, and survey the parts separately, as if they were lands belonging to different persons; and at last join them together.

6. As it is necessary to protract or lay down the work as you proceed in it, you must have a scale of a due length to do it by. To get such a scale, measure the whole length of the estate in chains; then consider how many inches long the map is to be; and from these will be known how many chains you must have in an inch; then make the scale accordingly, or choose one already made.

PROBLEM XII.

To survey a County, or large Tract of Land.

1. CHOOSE two, three, or four eminent places, for stations; such as the tops of high hills or mountains, towers, or church steeples which may be seen from one another; from which most of the towns and other places of note may also be seen; and so as to be as far distant from one another as possible. On these places raise beacons, or long poles, with flags of different colours flying at them, so as to be visible from all the other stations.

2. At all the places which you would set down in the map, plant long poles, with flags at them of several colours to distinguish the places from one another; fixing them on the tops of church steeples, or the tops of houses; or in the centres of smaller towns and villages.

These marks then being set up at a convenient number of places, and such as may be seen from both stations; go to one of these stations, and, with an instrument to take angles, standing at that station, take all the angles between the other station and each of these marks. Then go to the other station, and take all the angles between the first station and each of the former marks, setting them down with the others, each against its fellow with the same colour. You may, if convenient, also take the angles at some third station, which may serve to prove the work, if the three lines intersect in that point where any mark stands. The marks must stand till the observations are finished at both stations; and then they may be taken down, and set up at new places. The same operations must be performed, at both stations, for these new places; and the like for others. The instrument for taking

taking angles must be an exceeding good one, made on purpose with telescopic sights, and of a good length of radius.

3. And though it be not absolutely necessary to measure any distance, because, a stationary line being laid down from any scale, all the other lines will be proportional to it; yet it is better to measure some of the lines, to ascertain the distances of places in miles, and to know how many geometrical miles there are in any length; as also from thence to make a scale to measure any distance in miles. In measuring any distance, it will not be exact enough to go along the high roads; which by reason of their turnings and windings, hardly ever lie in a right line between the stations; which must cause endless reductions, and require great trouble to make it a right line; for which reason it can never be exact. But a better way is to measure in a straight line with a chain, between station and station, over hills and dales, or level fields, and all obstacles. Only in case of water, woods, towns, rocks, banks, &c. where we cannot pass, such parts of the line must be measured by the methods of inaccessible distances; and besides allowing for ascents and descents, when they are met with. A good compass, that shows the bearing of the two stations, will always direct us to go straight, when the two stations cannot be seen; and in the progress, if we can go straight, offsets may be taken to any remarkable places, likewise noting the intersection of the station-line with all roads, rivers, &c.

4. From all the stations, and in the whole progress, we must be very particular in observing sea-coast, river-mouths, towns, castles, houses, churches, mills, trees, rocks, sands, roads, bridges, fords, ferries, woods, hills, mountains, rills, brooks, parks, beacons, sluices, floodgates, locks, &c. and in general every thing that is remarkable.


5. After we have done with the first and main station-lines, which command the whole county; we must then take inner stations at some places already determined; which will divide the whole into several partitions: and from these stations we must determine the places of as many of the remaining towns as we can. And if any remain in that part, we must take more stations, at some places already determined, from which we may determine the rest. And thus go through all the parts of the county, taking station after station, till we have determined the whole. And in general the station-distances must always pass through such remarkable points as have been determined before, by the former stations.

PROBLEM XIII.

To Survey a Town or City.

THIS may be done with any of the instruments for taking angles, but best of all with the plain table, where every minute part is drawn while in sight. Instead of the common surveying or Gunter's chain, it will be best, for this purpose, to have a chain 50 feet long, divided into 50 links of one foot each, and an offset-staff of 10 feet long.

Begin at the meeting of two or more of the principal streets, through which we can have the longest prospects, to get the longest station-lines: there having fixed the instrument, draw lines of direction along those streets, using two men as marks, or poles set in wooden pedestals, or perhaps some remarkable places in the houses at the farther ends, as windows, doors, corners, &c. Measure these lines with the chain, taking offsets with the staff, at all corners of streets, bendings, or windings, and to all remarkable things, as churches, markets, halls, colleges, eminent houses, &c. Then remove the instrument to another station, along one of these lines; and there repeat the same process as before. And so on till the whole is finished.



Thus, fix the instrument at A, and draw lines in the direction of all the streets meeting there; then measure AB, noting the street on the left at m. At the second station B, draw the directions of the streets meeting there; and measure from B to c, noting the places of the streets at n and o as you pass by them. At the third station c, take the direction of all the streets meeting there, and measure cd. At D do the same, and measure DE noting the place of the cross streets at p. And in this manner go through all the principal streets. This done, proceed to the smaller and intermediate streets; and lastly to the lanes, alleys, courts, yards, and every part that it may be thought proper to represent in the plan.

PROBLEM

PROBLEM XIV.

To lay down the Plan of any Survey.

If the survey was taken with the plain table, we have a rough plan of it already on the paper which covered the table. But if the survey was with any other instrument, a plan of it is to be drawn from the measures that were taken in the survey; and first of all a rough plan on paper.

To do this, you must have a set of proper instruments, for laying down both lines and angles, &c.; as scales of various sizes (the more of them, and the more accurate the better), scales of chords, protractors, perpendicular and parallel rulers, &c. Diagonal scales are best for the lines, because they extend to three figures, or chains, and links, which are 100 parts of chains. But in using the diagonal scale, a pair of compasses must be employed, to take off the lengths of the principal lines very accurately. But a scale with a thin edge divided, is much readier for laying down the perpendicular offsets to crooked hedges, and for marking the places of those offsets on the station-line; which is done at only one application of the edge of the scale to that line, and then pricking off all at once the distances along it. Angles are to be laid down either with a good scale of chords, which is perhaps the most accurate way, or with a large protractor, which is much readier when many angles are to be laid down at one point, as they are pricked off all at once round the edge of the protractor.

In general, all lines and angles must be laid down on the plan in the same order in which they were measured in the field, and in which they are written in the field-book; laying down first the angles for the position of lines, next the lengths of the lines, with the places of the offsets, and then the lengths of the offsets themselves, all with dry or obscure lines; then a black line drawn through the extremities of all the offsets, will be the hedge or bounding line of the field, &c. After the principal bounds and lines are laid down, and made to fit or close properly, proceed next to the smaller objects, till you have entered every thing that ought to appear in the plan, as houses, brooks, trees, hills, gates, stiles, roads, lanes, mills, bridges, woodlands, &c. &c.

The north side of a map or plan is commonly placed uppermost, and a meridian is some where drawn, with the compass or flower-de-luce pointing north. Also, in a vacant part, a scale of equal parts or chains is drawn, with the title of the map in conspicuous characters, and embellished with a compartment. Hills are shadowed to distinguish them in the map. Colour the hedges with different colours; represent hilly grounds

grounds by broken hills and valleys ; draw single dotted lines for foot-paths, and double ones for horse or carriage roads. Write the name of each field and remarkable place within it, and, if you choose, its content in acres, roods, and perches.

In a very large estate, or a county, draw vertical and horizontal lines through the map, denoting the spaces between them by letters placed at the top, and bottom, and sides, for readily finding any field or other object mentioned in a table.

In mapping counties, and estates that have uneven grounds of hills and valleys, reduce all oblique lines, measured up-hill and down-hill, to horizontal straight lines, if that was not done during the survey, before they were entered in the field-book, by making a proper allowance to shorten them. For which purpose there is commonly a small table engraven on some of the instruments for surveying.

THE NEW METHOD OF SURVEYING,

PROBLEM XV.

To Survey and Plan by the New Method.

IN the former method of measuring a large estate, the accuracy of it depends both on the correctness of the instruments, and on the care in taking the angles. To avoid the errors incident to such a multitude of angles, other methods have of late years been used by some few skilful surveyors : the most practical, expeditious, and correct, seems to be the following, which is performed, without taking angles, by measuring with the chain only.

Choose two or more eminences, as grand stations, and measure a principal base line from one station to another ; noting every hedge, brook, or other remarkable object, as you pass by it ; measuring also such short perpendicular lines to the bends of hedges as may be near at hand. From the extremities of this base line, or from any convenient parts of the same, go off with other lines to some remarkable object situated towards the sides of the estate, without regarding the angles they make with the base line or with one another ; still remembering to note every hedge, brook, or other object, that you pass by. These lines, when laid down by intersections, will, with the base line, form a grand triangle on the estate ; several of which, if need be, being thus measured and laid down, you may proceed to form other smaller triangles and trapezoids on the sides of the former : and so on till you finish with the enclosures individually. By which means a kind of skeleton of the estate may first be obtained,
and

and the chief lines serve as the bases of such triangles and trapezoids as are necessary to fill up all the interior parts.

The field-book is ruled into three columns, as usual. In the middle one are set down the distances on the chain-line, at which any mark, offset, or other observation, is made; and in the right and left hand columns are entered the offsets and observations made on the right and left hand respectively of the chain-line; sketching on the sides the shape or resemblance of the fences or boundaries.

It is of great advantage, both for brevity and perspicuity, to begin at the bottom of the leaf, and write upwards; denoting the crossing of fences, by lines drawn across the middle column, or only a part of such a line on the right and left opposite the figures, to avoid confusion; and the corners of fields, and other remarkable turns in the fences where offsets are taken to, by lines joining in the manner the fences do; as will be best seen by comparing the book with the plan annexed to the field-book following, p. 450.

The letter in the left-hand corner at the beginning of every line, is the mark or place measured *from*; and that at the right hand corner at the end, is the mark measured *to*: But when it is not convenient to go exactly from a mark, the place measured from is described *such a distance from one mark towards another*; and where a former mark is not measured to, the exact place is ascertained by saying, turn to the right or left hand, *such a distance to such a mark*, it being always understood that those distances are taken in the chain-line.

The characters used are, $\{$ for *turn to the right hand*, $\}$ for *turn to the left hand*, and \sim placed over an offset, to show that it is not taken at right angles with the chain-line, but in the direction of some straight fence; being chiefly used when crossing their directions; which is a better way of obtaining their true places than by offsets at right angles.

When a line is measured whose position is determined, either by former work (as in the case of producing a given line, or measuring from one known place or mark to another) or by itself (as in the third side of the triangle), it is called a *fast line*, and a double line across the book is drawn at the conclusion of it; but if its position is not determined (as in the second side of the triangle), it is called a *loose line*, and a single line is drawn across the book. When a line becomes determined in position, and is afterwards continued farther, a double line half through the book is drawn.

When a loose line is measured, it becomes absolutely necessary to measure some other line that will determine its position. Thus, the first line *ah*, or *bh*, being the base of a

triangle is always determined ; but the position of the second side hj , does not become determined, till the third side jb is measured ; then the position of both is determined, and the triangle may be constructed.

At the beginning of a line, to fix a loose line to the mark or place measured from, the sign of turning to the right or left hand must be added, as at h in the second, and j in the third line ; otherwise a stranger, when laying down the work, may as easily construct the triangle hjb on the wrong side of the line ah , as on the right one ; but this error cannot be fallen into, if the sign above named be carefully observed.

In choosing a line to fix a loose one, care must be taken that it does not make a very acute or obtuse angle ; as in the triangle pbr , by the angle at b being very obtuse, a small deviation from truth, even the breadth of a point at p or r would make the error at b when constructed, very considerable ; but by constructing the triangle pbq , such a deviation is of no consequence.

Where the words *leave off* are written in the field-book, it signifies that the taking of offsets is from thence discontinued ; and of course something is wanting between that and the next offset, to be afterwards determined by measuring some other line.


The field-book for this method, and the plan drawn from it, are contained in the four following pages, engraven on copper-plates ; answerable to which, the pupil is to draw a plan from the measures in the field-book, of a larger size, viz. to a scale of a double size will be convenient, such a scale being also found on most instruments. In doing this, begin at the commencement of the field-book, or bottom of the first page and draw the first line ah in any direction at pleasure, and then the next two sides of the first triangle bhj by sweeping intersecting arcs ; and so all the triangles in the same manner, after each other in their order ; and afterwards setting the perpendicular and other offsets at their proper places, and through the ends of them drawing the bounding fences.

Note. That the field-book begins at the bottom of the first page, and reads up to the top ; hence it goes to the bottom of the next page, and to the top ; and thence it passes from the bottom of the third page to the top which is the end of the field-book. The several marks measured to or from, are here denoted by the letters of the alphabet, first the small ones a , b , c , d , &c. and after them the capitals A , B , C , D , &c. But, instead of these letters, some surveyors use the numbers in order, 1, 2, 3, 4, &c.

<p><i>n</i></p>	<p>1310 836 684</p>	<p>756 to <i>e</i> 56 50</p>
<p><i>m</i></p>	<p>1480 960 930 700 400</p>	<p>90 to <i>f</i> 24 <i>n</i> 48 30</p>
<p><i>k</i></p>	<p>1430 1290 1004 980 610 280</p>	<p>to <i>i</i> 40 36 <i>m</i> 24 32</p>
<p><i>l</i></p>	<p>1820 1464 1050 920 650 350 0</p>	<p>to <i>l</i> 22 32 60 48 14</p>
<p><i>j</i></p>	<p>3074 2494 2100 0 2072 54 80 1730 1530 1420 56 + 30 1170 52 620 32 280</p>	<p>to <i>b</i> <i>l</i> <i>k</i> 40</p>
<p><i>h</i></p>	<p>2574 2494 2000 1860 1840 50 1794 34 + 50 1464 70 1328 96 1240 52 + 34 1130 34 860 66 190</p>	<p>to <i>j</i> 44 50 <i>i</i></p>
<p><i>a</i></p>	<p>4450 3570 2620 2610 2210 2080 1640 1550 1510 990 806</p>	<p><i>h</i> <i>g</i> <i>f</i> <i>e</i> <i>d</i> <i>c</i> <i>b</i></p>

		768	to A
		526	70
		496	
	70	460	
	40	124	
D	┌	100	
		455	D
		400	76
		48	10
C			
		600	to r
		432	C
	56	160	
B		36	
	44		
B	┌	152	to q
		480	B
	24	160	
p			
	0	1700	
		1560	┌ 44 to x
		980	
		885	A
	44	666	
	79	310	x
d	60	236	
		2148	┌ 480 to b
		1950	y
		1836	
	126	1724	
	60	1600	
	30	1480	x
	0	1320	
	50	1110	
		1080	
		840	w
u	┌	750	50
		4440	36
		4420	v
		3884	u
		3380	60
		2992	90
		2692	t
	120	2624	
		2592	
		2500	s
		2070	56
		1900	leave off
		1840	r
	60	1770	
		1320	q
		808	p
	leave off	650	
	40	360	
	80	170	
	20		
v	┌	220	0
		190	46
h produced from i			

		580	to v
		40 500	
		70 300	
F		76 100	
		420	to F
J		20 150	
		954	J
		15 850	
		740	to E
		30 490	
		340	60
		0 280	
I		20 170	50
		725	to H
		672	0
		70 150	0
a		50 13	
		1160	to v
		32 1000	
		890	
		780	32
		590	40
		570	I
		530	40
		376	H
		256	150
		190	64
G		144	130
		1676	G
		1676	30
		396	24
		632	
		620	50
180 from u towards v		588	F
		620	to f
D		488	32
		2260	
		2250	E
		2210	
		2050	
		2030	
		1990	130 to w
		1552	180
		1380	96
		950	110
		860	8



Whole Content A. R. P.
103, 2, 10.

OF THE OLD KIND OF FIELD-BOOK.

In surveying with the plain table, a field-book is not used, as every thing is drawn on the table immediately when it is measured. But in surveying with the theodolite, or any other instrument, some kind of a field-book must be used, to write down in it a register or account of all that is done and occurs relative to the survey in hand

This book every one contrives and rules as he thinks fittest for himself. The following is a specimen of a form which has been formerly used. It is ruled into three columns, as below.


Here $\odot 1$ is the first station, where the angle or bearing is $105^\circ 25'$. On the left, at 73 links in the distance or principal line, is an offset of 92; and at 610 an offset of 24 to a cross hedge. On the right at 0. or the beginning, an offset 25 to the corner of the field; at 248 Brown's boundary hedge commences; at 610 an offset 35; and at 954, the end of the first line, the 0 denotes its terminating in the hedge. And so on for the other stations.

A line is drawn under the work, at the end of every station line, to prevent confusion.

Form of this Field-Book.

Offsets and remarks on the left.	Stations, Bearings, and Distances.	Offsets and Remarks on the right.
	$\odot 1$ $105^\circ 25'$	
80	00	25 corner
92	73	
a cross hedge 24	248	Brown's hedge
	610	35
	954	00
	$\odot 2$ $53^\circ 10'$	
house corner 51	25	21
	120	29 a tree
34	734	40 a stile
	$\odot 3$ $67^\circ 20'$	
a brook 30	61	35
	248	
foot path 16	639	16 a spring
cross hedge 18	810	
	973	20 a pond
		Then

Then the plan, on a small scale drawn from the above field-book, will be as in the following figure. But the pupil may draw a plan of 3 or 4 times the size on his paper book. The dotted lines denote the 3 chain or measured lines, and the black lines the boundaries on the right and left.




But some skilful surveyors now make use of a different method for the field-book, namely, beginning at the bottom of the page and writing upwards; sketching also a neat boundary on either hand, resembling the parts near the measured lines as they pass along; an example of which will be given further on, in the method of surveying a large estate.

In smaller surveys and measurements, a good way of setting down the work, is, to draw by the eye on a piece of paper, a figure resembling that which is to be measured; and so writing the dimensions, as they are found, against the corresponding parts of the figure. And this method may be practised to a considerable extent, even in the larger surveys.

Another specimen of a field-book, with its plan, is as follows; being a single field, surveyed with the chain, and the theodolite for taking angles; which the pupil will likewise draw of a larger size.

	⊙ A			⊙ C
	82° 55'			57° 10'
0	0		35	268
40	230		50	470
48	572		0	846
30	860		30	1140
	⊙ B			⊙ D
	130° 35'			
0	238		40	0
20	520		25	117
			45	312
			0	554



SECTION III.

OF COMPUTING AND DIVIDING.

PROBLEM XVI.

To Compute the Contents of Fields.

1. COMPUTE the contents of the figures as divided into triangles, or trapeziums, by the proper rules for these figures laid down in measuring; multiplying the perpendiculars by the diagonals or bases, both in links, and divide by 2; the quotient is acres, after having cut off five figures on the right for decimals. Then bring these decimals to roods and perches, by multiplying first by 4, and then by 40. An example of which is given in the description of the chain, pag. 429.

2. In small and separate pieces, it is usual to compute their contents from the measures of the lines taken in surveying them, without making a correct plan of them.

3. In pieces bounded by very crooked and winding hedges, measured by offsets, all the parts between the offsets are most accurately measured separately as small trapezoids.

4. Sometimes such pieces as that last mentioned, are computed by finding a mean breadth, by adding all the offsets together, and dividing the sum by the number of them, accounting that for one of them where the boundary meets the station-line, (which increases the number of them by 1, for the divisor, though it does not increase the sum or quantity to be divided); then multiply the length by that mean breadth.

5. But in larger pieces and whole estates, consisting of many fields, it is the common practice to make a rough plan of the whole, and from it compute the contents, quite independent of the measures of the lines and angles that were taken in surveying

surveying. For then new lines are drawn in the fields on the plan, so as to divide them into trapeziums and triangles, the bases and perpendiculars of which are measured on the plan by means of the scale from which it was drawn, and so multiplied together for the contents. In this way, the work is very expeditiously done, and sufficiently correct; for such dimensions are taken as afford the most easy method of calculation; and among a number of parts, thus taken and applied to a scale, though it be likely that some of the parts will be taken a small matter too little, and others too great, yet they will, on the whole, in all probability, very nearly balance one another, and give a sufficiently accurate result. After all the fields and particular parts are thus computed separately, and added all together into one sum: calculate the whole estate independent of the fields by dividing it into large and arbitrary triangles and trapeziums, and add these all together. Then if this sum be equal to the former, or nearly so, the work is right; but if the sums have any considerable difference, it is wrong, and they must be examined, and re-computed, till they nearly agree.


6. But the chief art in computing, consists in finding the contents of pieces bounded by curved or very irregular lines, or in reducing such crooked sides of fields or boundaries to straight lines, that shall inclose the same or equal area with those crooked sides, and so obtain the area of the curved figure by means of the right-lined one, which will commonly be a trapezium. Now this reducing the crooked sides to straight ones, is very easily and accurately performed in this manner:—Apply the straight edge of a thin, clear piece of lanthorn-horn to the crooked line, which is to be reduced, in such a manner, that the small parts cut off from the crooked figure by it, may be equal to those which are taken in: which equality of the parts included and excluded you will presently be able to judge of very nicely by a little practice; then with a pencil, or point of a tracer, draw a line by the straight edge of the horn. Do the same by the other sides of the field or figure. So shall you have a straight-sided figure equal to the curved one; the content of which, being computed as before directed, will be the content of the crooked figure proposed.

Or, instead of the straight edge of the horn, a horse hair, or fine thread, may be applied across the crooked sides in the same manner: and the easiest way of using the thread, is to string a small slender bow with it, either of wire or cane, or whalebone, or such like slender elastic matter; for the bow keeping it always stretched, it can be easily and neatly applied with one hand, while the other is at liberty to make two marks by the side of it, to draw the straight line by.

EXAMPLE

EXAMPLE.

Thus, let it be required to find the contents of the same figure as in Prob. ix, page 441, to a scale of 4 chains to an inch.



Draw the 4 dotted straight lines AB, BC, CD, DA, cutting off equal quantities on both sides of them: which they do as near as the eye can judge: so is the crooked figure reduced to an equivalent right-lined one of 4 sides. ABCD. Then draw the diagonal BD, which, by applying a proper scale to it, measures suppose 1256. Also the perpendicular or nearest distance from A to this diagonal, measures 456; and the distance of C from it, is 428.

Then, half the sum of 456 and 428, multiplied by the diagonal 1256, gives 55515² square links, or 5 acres, 2 roods, 8 perches, the content of the trapezium, or of the irregular crooked piece.

As a general example of this practice. let the contents be computed of all the fields separately in the foregoing plan in page 452, and by adding the contents altogether, the whole sum or content of the estate will be found nearly equal to 103 $\frac{1}{2}$ acres. Then, to prove the work, divide the whole plan into two parts, by a pencil line drawn across it any way near the middle, as from the corner *l* on the right, to the corner near *s* on the left; then by computing these two large parts separately, their sum must be nearly equal to the former sum, when the work is all right.

PROBLEM XVII.

To Transfer a Plan to Another Paper, &c.

AFTER the rough plan is completed, and a fair one is wanted; this may be done by any of the following methods.

First

First Method.—Lay the rough plan on the clean paper, keeping them always pressed flat and close together, by weights laid on them. Then, with the point of a fine pin or pricker, prick through all the corners of the plan to be copied. Take them asunder, and connect the pricked points, on the clean paper, with lines; and it is done. This method is only to be practised in plans of such figures as are small and tolerably regular, or bounded by right lines.

Second Method.—Rub the back of the rough plan over with black-lead powder; and lay this blacked part on the clean paper on which the plan is to be copied, and in the proper position. Then, with the blunt point of some hard substance, as brass or such-like, trace over the lines of the whole plan; pressing the tracer so much, as that the black lead under the lines may be transferred to the clean paper: after which, take off the rough plan, and trace over the leaden marks with common ink, or with Indian ink—Or, instead of blacking the rough plan, we may keep constantly a blacked paper to lay between the plans.

Third Method—Another method of copying plans, is by means of squares. This is performed by dividing both ends and sides of the plan which is to be copied into any convenient number of equal parts, and connecting the corresponding points of division with lines: which will divide the plan into a number of small squares. Then divide the paper, on which the plan is to be copied, into the same number of squares, each equal to the former when the plan is to be copied of the same size, but greater or less than the others, in the proportion in which the plan is to be increased or diminished, when of a different size. Lastly, copy into the clean squares the parts contained in the corresponding squares of the old plan; and you will have the copy, either of the same size, or greater or less in any proportion.

Fourth Method.—A fourth method is by the instrument called a pentagraph, which also copies the plan in any size required.

Fifth Method.—But the neatest method of any, at least in copying from a fair plan, is this. Procure a copying frame or glass, made in this manner; namely, a large square of the best window glass, set in a broad frame of wood, which can be raised up to any angle, when the lower side of it rests on a table. Set this frame up to any angle before you, facing a strong light; fix the old plan and clean paper together, with several pins quite around, to keep them together, the clean
paper

paper being laid uppermost, and over the face of the plan to be copied. Lay them, with the back of the old plan, on the glass; namely, that part which you intend to begin at to copy first, and by means of the light shining through the papers you will very distinctly perceive every line of the plan through the clean paper. In this state then trace all the lines on the paper with a pencil. Having drawn that part which covers the glass, slide another part over the glass, and copy it in the same manner. Then another part. And so on, till the whole is copied. Then take them asunder, and trace all the pencil lines over with a fine pen and Indian ink, or with common ink. And thus you may copy the finest plan without injuring it in the least.

OF ARTIFICERS' WORKS,

AND

TIMBER MEASURING.

1. OF THE CARPENTER'S OR SLIDING RULE.

THE Carpenter's or Sliding Rule, is an instrument much used in measuring of timber and artificers' works, both for taking the dimensions, and computing the contents.

The instrument consists of two equal pieces, each a foot in length, which are connected together by a folding joint.

One side or face of the rule, is divided into inches, and eighths, or half-quarters. On the same face also are several plane scales, divided into twelfth parts by diagonal lines; which are used in planning dimensions that are taken in feet and inches. The edge of the rule is commonly divided decimally, or into tenths; namely, each foot into ten equal parts, and each of these into ten parts again: so that by means of this last scale, dimensions are taken in feet, tenths, and hundredths, and multiplied as common decimal numbers, which is the best way.

On the one part of the other face are four lines, marked A, B, C, D; the two middle ones B and C being on a slider, which runs in a groove made in the stock. The same numbers serve for both these two middle lines, the one being above the numbers, and the other below.

These four lines are logarithmic ones, and the three A, B, C, which are all equal to one another, are double lines, as they proceed twice over from 1 to 10. The other or lowest line, D, is a single one, proceeding from 4 to 40. It is also called the girt line, from its use in computing the contents of trees and timber; and on it are marked wg at 17.15, and AG at 18.95, the wine and ale gage points, to make this instrument serve the purpose of a gaging rule.

On the other part of this face, there is a table of the value of a load, or 50 cubic feet, of timber, at all prices, from 6 pence to 2 shillings a foot.

When 1 at the beginning of any line is accounted 1, then the 1 in the middle will be 10, and the 10 at the end 100; but when one at the beginning is counted 10, then the one in the middle is 100, and the 10 at the end 1000; and so on. And all the smaller divisions are altered proportionally.

II. ARTIFICERS' WORK.

ARTIFICERS compute the contents of their works by several different measures. As,

Glazing and masonry, by the foot; Painting, plastering, paving, &c. by the yard, of 9 square feet; Flooring, partitioning, roofing, tiling, &c. by the square of 100 square feet:

And brickwork, either by the yard of 9 square feet, or by the perch, or square rod or pole, containing $272\frac{1}{4}$ square feet, or $30\frac{1}{4}$ square yards, being the square of the rod or pole of $16\frac{1}{2}$ feet or $5\frac{1}{2}$ yards long.

As this number $272\frac{1}{4}$ is troublesome to divide by, the $\frac{1}{4}$ is often omitted in practice, and the content in feet divided only by the 272.

All works, whether superficial or solid, are computed by the rules proper to the figure of them, whether it be a triangle, or rectangle, a parallelopiped, or any other figure.

III. BRICKLAYERS' WORK.

BRICKWORK is estimated at the rate of a brick and a half thick. So that if a wall be more or less than this standard thickness, it must be reduced to it, as follows:

Multiply the superficial content of the wall by the number of half bricks in the thickness, and divide the product by 3.

The

The dimensions of a building may be taken by measuring half round on the outside and half round it on the inside ; the sum of these two gives the compass of the wall, to be multiplied by the height, for the content of the materials.

Chimneys are commonly measured as if they were solid, deducting only the vacuity from the hearth to the mantle, on account of the trouble of them. All windows, doors, &c. are to be deducted out of the contents of the walls in which they are placed.

EXAMPLES.

EXAM. 1. How many yards and rods of standard brick-work are in a wall whose length or compass is 57 feet 3 inches, and height 24 feet 6 inches ; the wall being $2\frac{1}{2}$ bricks or 5 half-bricks thick ? Ans. 8 rods, $17\frac{2}{3}$ yards.

EXAM. 2. Required the content of a wall 62 feet 6 inches long, and 14 feet 8 inches high, and $2\frac{1}{2}$ bricks thick ? Ans. 169·753 yards.

EXAM. 3. A triangular gable is raised $17\frac{1}{2}$ feet high, on an end wall whose length is 24 feet 9 inches, the thickness being 2 bricks ; required the reduced content ? Ans. 32·08 $\frac{1}{3}$ yards.

EXAM. 4. The end wall of a house is 28 feet 10 inches long, and 55 feet 8 inches high, to the eaves ; 20 feet high is $2\frac{1}{2}$ bricks thick, other 20 feet high is 2 bricks thick, and the remaining 15 feet 8 inches is $1\frac{1}{2}$ brick thick ; above which is a triangular gable, of 1 brick thick : which rises 42 courses of bricks, of which every 4 courses make a foot. What is the whole content in standard measure ? Ans. 253·626 yards.

IV. MASONS' WORK.

To masonry belong all sorts of stone-work ; and the measure made use of is a foot, either superficial or solid.

Walls, columns, blocks of stone or marble, &c. are measured by the cubic foot ; and pavements, slabs, chimney-pieces, &c. by the superficial or square foot.

Cubic or solid measure is used for the materials, and square measure for the workmanship.

In the solid measure, the true length, breadth, and thickness are taken, and multiplied continually together. In the superficial, there must be taken the length and breadth of every part of the projection which is seen without the general upright face of the building.

EXAMPLES.

EXAMPLES.

EXAM. 1. Required the solid content of a wall, 53 feet 6 inches long, 12 feet 3 inches high, and 2 feet thick ?

Ans. 1310 $\frac{3}{4}$ feet,

EXAM. 2. What is the solid content of a wall, the length being 24 feet 3 inches, height 10 feet 9 inches, and 2 feet thick ?

Ans. 521.375 feet.

EXAM. 3. Required the value of a marble slab, at 8s. per foot ; the length being 5 feet 7 inches, and breadth 1 foot 10 inches ?

Ans. 4l. 1s. 10 $\frac{1}{2}$ d.

EXAM. 4. In a chimney-piece, suppose the length of the mantle and slab, each 4 feet 6 inches

breadth of both together - 3 2

length of each jamb - - 4 4

breadth of both together - 1 9

Required the superficial content ? Ans. 21 feet 10 inches.

V. CARPENTERS' AND JOINERS' WORK.

To this branch belongs all the wood-work of a house, such as flooring, partitioning, roofing, &c.

Large and plain articles are usually measured by the square foot or yard, &c. ; but enriched mouldings, and some other articles, are often estimated by running or lineal measure ; and some things are rated by the piece.

In measuring of Joists, take the dimensions of one joist, and multiply its content by the number of them ; considering that each end is let into the wall about $\frac{2}{3}$ of the thickness, as it ought to be.

Partitions are measured from wall to wall for one dimension, and from floor to floor, as far as they extend, for the other.

The measure of Centering for Cellars is found by making a string pass over the surface of the arch for the breadth, and taking the length of the cellar for the length : but in groin centering, it is usual to allow double measure, on account of their extraordinary trouble.

In Roofing, the dimensions as to length, breadth, and depth, are taken as in flooring joists, and the contents computed the same way.

In Floor-boarding, take the length of the room for one dimension, and the breadth for the other, to multiply together for the content.

For Stair-cases, take the breadth of all the steps, by making a line

a line ply close over them, from the top to the bottom, and multiply the length of this line by the length of a step, for the whole area.—By the length of a step is meant the length of the front and the returns at the two ends; and by the breadth is to be understood the girts of its two outer surfaces, or the tread and riser.

For the Balustrade, take the whole length of the upper part of the hand-rail, and girt over its end till it meet the top of the newel post, for the one dimension; and twice the length of the baluster on the landing, with the girt of the hand-rail, for the other dimension.

For Wainscoting, take the compass of the room for the one dimension; and the height from the floor to the ceiling, making the string ply close into all the mouldings, for the other.

For Doors, take the height and the breadth, to multiply them together for the area.—If the door be panned on both sides, take double its measure for the workmanship; but if one side only be panned, take the area and its half for the workmanship.—*For the Surrounding Architrave*, girt it about the uppermost part, for its length; and measure over it, as far as it can be seen when the door is open, for the breadth.

Window-shutters, Bases, &c. are measured in like manner.

In measuring of Joiners' work, the string is made to ply close into all the mouldings, and to every part of the work over which it passes.

EXAMPLES.

EXAM. 1. Required the content of a floor, 48 feet 6 inches long, and 24 feet 3 inches broad? Ans. 11 sq. 76 $\frac{1}{2}$ feet.

EXAM. 2. A floor being 36 feet 3 inches long, and 16 feet 6 inches broad, how many squares are in it? Ans. 5 sq. 98 $\frac{1}{2}$ feet.

EXAM. 3. How many squares are there in 173 feet 10 inches in length, and 10 feet 7 inches height, of partitioning? Ans. 18.3973 squares.

EXAM. 4. What cost the roofing of a house at 10s. 6d. a square; the length within the walls being 52 feet 8 inches, and the breadth 30 feet 6 inches; reckoning the roof $\frac{2}{3}$ of the flat? Ans. 12l. 12s. 11 $\frac{3}{4}$ d.

EXAM.

EXAM. 5. To how much, at 6s. per square yard, amounts the wainscoting of a room; the height, taking in the cornice and mouldings, being 12 feet 6 inches, and the whole compass 83 feet 8 inches; also the three window-shutters are each 7 feet 8 inches by 3 feet 6 inches, and the door 7 feet by 3 feet 6 inches; the doors and shutters, being worked on both sides, are reckoned work and half work?

Ans. 36l. 12s. 2½d.

VI. SLATERS' AND TILERS' WORK.

In these articles, the content of a roof is found by multiplying the length of the ridge by the girt over from eaves to eaves; making allowance in this girt for the double row of slates at the bottom, or for how much one row of slates or tiles is laid over another.

When the roof is of a true pitch, that is, forming a right angle at top; then the breadth of the building, with its half added, is the girt over both sides nearly.

In angles formed in a roof, running from the ridge to the eaves, when the angle bends inwards, it is called a valley; but when outwards, it is called a hip.

Deductions are made for chimney shafts or window holes.

EXAMPLES.

EXAM. 1. Required the content of a slated roof, the length being 45 feet 9 inches, and the whole girt 34 feet 3 inches?

Ans. $174\frac{5}{8}$ yards.

EXAM. 2. To how much amounts the tiling of a house, at 25s. 6d. per square; the length being 43 feet 10 inches, and the breadth on the flat 27 feet 5 inches; also the eaves projecting 16 inches on each side, and the roof of a true pitch?

Ans. 24l. 9s. 5¼d.

VII. PLASTERERS' WORK.

PLASTERERS' work is of two kinds; namely, ceiling, which is plastering on laths; and rendering, which is plastering on walls: which are measured separately.

The contents are estimated either by the foot or the yard, or the square, of 100 feet. Inriched mouldings, &c. are rated by running or lineal measure.

Deductions are made for chimneys, doors, windows, &c.

EXAMPLES.

EXAM. 1. How many yards contains the ceiling which is 43 feet 3 inches long, and 25 feet 6 inches broad ?

Ans. 122½.

EXAM. 2. To how much amounts the ceiling of a room, at 10*d.* per yard; the length being 21 feet 8 inches, and the breadth 14 feet 10 inches ?

Ans. 1*l.* 9*s.* 8¾*d.*

EXAM. 3. The length of a room is 18 feet 6 inches, the breadth 12 feet 3 inches, and height 10 feet 6 inches; to how much amounts the ceiling and rendering, the former at 8*d.* and the latter at 3*d.* per yard; allowing for the door of 7 feet by 3 feet 8, and a fire-place of 5 feet square ?

Ans. 1*l.* 13*s.* 3¼*d.*

EXAM. 4. Required the quantity of plastering in a room, the length being 14 feet 5 inches, breadth 13 feet 2 inches, and height 9 feet 3 inches to the under side of the cornice, which girts 8½ inches, and projects 5 inches from the wall on the upper part next the ceiling; deducting only for a door 7 feet by 4 ?

Ans. 53 yards 5 feet 3½ inches of rendering

18	5	6	of ceiling
	39	0½	of cornice.

VIII. PAINTERS' WORK.

PAINTERS' work is computed in square yards. Every part is measured where the colour lies; and the measuring line is forced into all the mouldings and corners.

Windows are done at so much a piece. And it is usual to allow double measure for carved mouldings, &c.

EXAMPLES.

EXAM. 1. How many yards of painting contains the room which is 65 feet 6 inches in compass, and 12 feet 4 inches high ?

Ans. 89½ yards.

EXAM. 2. The length of a room being 20 feet, its breadth 14 feet

14 feet 6 inches, and height 10 feet 4 inches; how many yards of painting are in it, deducting a fire-place of 4 feet by 4 feet 4 inches, and two windows each 6 feet by 3 feet 2 inches?

Ans. $73\frac{2}{7}$ yards.

EXAM. 3. What cost the painting of a room, at 6*d.* per yard; its length being 24 feet 6 inches, its breadth 16 feet 3 inches, and height 12 feet 9 inches; also the door is 7 feet by 3 feet 6, and the window-shutters to two windows each 7 feet 9 by 3 feet 6; but the breaks of the windows themselves are 8 feet 6 inches high, and 1 foot 3 inches deep; including also the window sills or seats, and the soffits above, the dimensions of which are known from the other dimensions: but deducting the fire-place of 5 feet by 5 feet 6?

Ans. 3*l.* 3*s.* $10\frac{3}{4}$ *d.*

IX. GLAZIERS' WORK.

GLAZIERS take their dimensions, either in feet, inches, and parts, or feet, tenths, and hundredths. And they compute their work in square feet.

In taking the length and breadth of a window, the cross bars between the squares are included. Also windows of round or oval forms are measured as square, measuring them to their greatest length and breadth, on account of the waste in cutting the glass.

EXAMPLES.

EXAM. 1. How many square feet contains the window which is 4.25 feet long, and 2.75 feet broad?

Ans. $11\frac{2}{3}$.

EXAM. 2. What will the glazing of a triangle sky-light come to, at 10*d.* per foot; the base being 12 feet 6 inches, and the perpendicular height 6 feet 9 inches?

Ans. 1*l.* 15*s.* $1\frac{3}{4}$ *d.*

EXAM. 3. There is a house with three tiers of windows, three windows in each tier, their common breadth 3 feet 11 inches:

now the height of the first tier is 7 feet 10 inches

of the second 6 8

of the third 5 4

Required the expense of glazing at 14*d.* per foot?

Ans. 13*l.* 11*s.* $10\frac{1}{2}$ *d.*

EXAM.

EXAM. 4. Required the expense of glazing the windows of a house at 13*d.* a foot; there being three stories, and three windows in each story:

the height of the lower tier is 7 feet 9 inches

of the middle 6 6

of the upper 5 3 $\frac{1}{4}$

and of an oval window over the door 1 10 $\frac{1}{2}$

the common breadth of all the windows being 3 feet 9 inches?

Ans. 12*l.* 5*s.* 6*d.*

X. PAVERS' WORK.

PAVERS' work is done by the square yard. And the content is found by multiplying the length by the breadth.

EXAMPLES.

EXAM. 1. What cost the paving a foot-path, at 3*s.* 4*d.* a yard; the length being 35 feet 4 inches, and breadth 8 feet 3 inches? Ans. 5*l.* 7*s.* 11 $\frac{1}{2}$ *d.*

EXAM. 2. What cost the paving a court, at 3*s.* 2*d.* per yard; the length being 27 feet 10 inches, and the breadth 14 feet 9 inches? Ans. 7*l.* 4*s.* 5 $\frac{1}{2}$ *d.*

EXAM. 3. What will be the expense of paving a rectangular court-yard, whose length is 63 feet, and breadth 45 feet; in which there is laid a foot-path of 5 feet 3 inches broad, running the whole length, with broad stones, at 3*s.* a yard; the rest being paved with pebbles at 2*s.* 6*d.* a yard?

Ans. 40*l.* 5*s.* 10 $\frac{1}{2}$ *d.*

XI. PLUMBERS' WORK.

PLUMBERS' work is rated at so much a pound, or else by the hundred weight of 112 pounds.

Sheet lead, used in roofing, guttering, &c. is from 6 to 10 lb. to the square foot. And a pipe of an inch bore is commonly 13 or 14 lb. to the yard in length.

EXAMPLES.

EXAM. 1. How much weighs the lead which is 39 feet 6

inches long, and 3 feet 3 inches broad, at $8\frac{1}{2}$ lb. to the square foot ?

Ans. $1091\frac{3}{16}$ lb.

EXAM. 2. What cost the covering and guttering a roof with lead, at 18s. the cwt; the length of the roof being 43 feet, and breadth or girt over it 32 feet; the guttering 57 feet long, and 2 feet wide; the former 9.331 lb. and the latter 7.373 lb. to the square foot ?

Ans. 115l. 9s. $1\frac{1}{2}$ d.

XII. TIMBER MEASURING.

PROBLEM I.

To find the Area, or Superficial Content, of a Board or Plank.

MULTIPLY the length by the mean breadth.

Note. When the board is tapering, add the breadths at the two ends together, and take half the sum for the mean breadth. Or else take the mean breadth in the middle.

By the Sliding Rule.

Set 12 on B to the breadth in inches on A: then against the length in feet on B, is the content on A, in feet and fractional parts.

EXAMPLES.

EXAM. 1. What is the value of a plank, at $1\frac{1}{2}$ d. per foot, whose length is 12 feet 6 inches, and mean breadth 11 inches ?

Ans. 1s. 5d.

EXAM. 2. Required the content of a board, whose length is 11 feet 2 inches, and breadth 1 foot 10 inches ?

Ans. 20 feet 5 inches 8".

EXAM. 3. What is the value of a plank, which is 12 feet 9 inches long, and 1 foot 3 inches broad, at $2\frac{1}{2}$ d. a foot ?

Ans. 3s. $3\frac{3}{4}$ d.

EXAM. 4. Required the value of 5 oaken planks at 3d. per foot, each of them being $17\frac{1}{2}$ feet long; and their several breadths as follows, namely, two of $13\frac{1}{2}$ inches in the middle, one of $14\frac{1}{2}$ inches in the middle, and the two remaining ones, each 18 inches at the broader end, and $11\frac{1}{4}$ at the narrower ?

Ans. 1l. 5s. $9\frac{1}{2}$ d.

PROBLEM

PROBLEM II.

To find the Solid Content of Squared or Four-sided Timber.

MULTIPLY the mean breadth by the mean thickness, and the product again by the length, for the content nearly.

By the Sliding Rule.

$\begin{array}{cccc} & C & D & D & C \\ \text{As length} & : & 12 \text{ or } 10 & : & \text{quarter girt} & : & \text{solidity.} \end{array}$

That is, as the length in feet on *c*, is to 12 on *D*, when the quarter girt is in inches, or to 10 on *D*, when it is in tenths of feet; so is the quarter girt on *D*, to the content on *c*.

Note 1. If the tree taper regularly from the one end to the other; either take the mean breadth and thickness in the middle, or take the dimensions at the two ends, and half their sum will be the mean dimensions; which multiplied as above, will give the content nearly.

2. If the piece do not taper regularly, but be unequally thick in some parts and small in others; take several different dimensions, add them all together, and divide their sum by the number of them, for the mean dimensions.

EXAMPLES.

EXAM. 1. The length of a piece of timber is 18 feet 6 inches, the breadths at the greater and less end 1 foot 6 inches and 1 foot 3 inches, and the thickness at the greater and less end 1 foot 3 inches and 1 foot; required the solid content?

Ans. 28 feet 7 inches.

EXAM. 2. What is the content of the piece of timber, whose length is $24\frac{1}{2}$ feet, and the mean breadth and thickness each 1.04 feet?

Ans. $26\frac{1}{2}$ feet.

EXAM. 3. Required the content of a piece of timber, whose length is 20.38 feet, and its ends unequal squares, the sides of the greater being $19\frac{1}{8}$ inches, and the side of the less $9\frac{7}{8}$ inches?

Ans. 29.7562 feet.

EXAM. 4. Required the content of the piece of timber, whose length is 27.36 feet; at the greater end the breadth is 1.78, and thickness 1.23; and at the less end the breadth is 1.04, and thickness 0.91 feet?

Ans. 41.278 feet.

PROBLEM III.

To find the Solidity of Round or Unsquared Timber.

MULTIPLY the square of the quarter girt, or of $\frac{1}{4}$ of the mean circumference, by the length, for the content.

By the Sliding Rule.

As the length upon *c* : 12 or 10 upon *D* : :
quarter girt, in 12ths or 10ths, on *D* : content on *c*.

Note 1. When the tree is tapering, take the mean dimensions as in the former problems, either by girting it in the middle, for the mean girt, or at the two ends, and take half the sum of the two ; or by girting it in several places, then adding all the girts together, and dividing the sum by the number of them, for the mean girt. But when the tree is very irregular, divide it into several lengths, and find the content of each part separately.

2. This rule, which is commonly used, gives the answer about $\frac{1}{4}$ less than the true quantity in the tree, or nearly what the quantity would be, after the tree is hewed square in the usual way : so that it seems intended to make an allowance for the squaring of the tree.

EXAMPLES.

EXAM. 1. A piece of round timber being 9 feet 6 inches long, and its mean quarter girt 42 inches ; what is the content ?
Ans. 116 $\frac{1}{3}$ feet.

EXAM. 2. The length of a tree is 24 feet, its girt at the thicker end 14 feet, and at the smaller end 2 feet ; required the content ?
Ans. 96 feet.

EXAM. 3. What is the content of a tree, whose mean girt is 3.15 feet, and length 14 feet 6 inches ?

Ans. 8.9922 feet.

EXAM. 4. Required the content of a tree, whose length is 17 $\frac{1}{4}$ feet, which girts in five different places as follows, namely, in the first place 9.43 feet, in the second 7.92, in the third 6.15, in the fourth 4.74, and in the fifth 3.16 ?

Ans. 42.519525.

CONIC SECTIONS.

DEFINITIONS.


1. **CONIC SECTIONS** are the figures made by a plane cutting a cone.

2. According to the different positions of the cutting plane, there arise five different figures or sections, namely, a triangle, a circle, an ellipsis, an hyperbola, and a parabola: the three last of which only are peculiarly called Conic Sections.


3. If the cutting plane pass through the vertex of the cone, and any part of the base, the section will evidently be a triangle; as VAB .



4. If the plane cut the cone parallel to the base, or make no angle with it, the section will be a circle; as ABD .




5. The section DAB is an ellipse when the cone is cut obliquely through both sides, or when the plane is inclined to the base in a less angle than the side of the cone is.




6. The

6. The section is a parabola, when the cone is cut by a plane parallel to the side, or when the cutting plane and the side of the cone make equal angles with the base.



7. The section is an hyperbola, when the cutting plane makes a greater angle with the base than the side of the cone makes.

8. And if all the sides of the cone be continued through the vertex, forming an opposite equal cone, and the plane be also continued to cut the opposite cone, this latter section will be the opposite hyperbola to the former; as dbe .




9. The Vertices of any section, are the points where the cutting plane meets the opposite sides of the cone, or the sides of the vertex triangular section; as A and B.

Hence the ellipse and the opposite hyperbolas, have each two vertices; but the parabola only one; unless we consider the other as at an infinite distance.


10. The Axis, or Transverse Diameter, of a conic section, is the line or distance AB between the vertices.

Hence the axis of a parabola is infinite in length, ab being only a part of it.


Ellipse.



Hyperbolas.



Parabola.



11. The Centre c is the middle of the axis.

Hence the centre of a parabola is infinitely distant from the vertex. And of an ellipse, the axis and centre lie within the curve; but of an hyperbola, without.

12. A Diameter is any right line, as AB or DE , drawn through the centre, and terminated on each side by the curve; and the extremities of the diameter, or its intersections with the curve, are its vertices.

Hence all the diameters of a parabola are parallel to the axis, and infinite in length. And hence also every diameter of the ellipse and hyperbola have two vertices; but of the parabola, only one; unless we consider the other as at an infinite distance.

13. The Conjugate to any diameter, is the line drawn through the centre, and parallel to the tangent of the curve at the vertex of the diameter. So, FG , parallel to the tangent at D , is the conjugate to DE : and HI , parallel to the tangent at A , is the conjugate to AB .

Hence the conjugate HI , of the axis AB , is perpendicular to it.

14. An Ordinate to any diameter, is a line parallel to its conjugate, or to the tangent at its vertex, and terminated by the diameter and curve. So DK , EL , are ordinates to the axis AB ; and MN , NO , ordinates to the diameter DE .

Hence the ordinates of the axis are perpendicular to it.

15. An Absciss is a part of any diameter contained between its vertex and an ordinate to it; as AK or BK , or DN or EN .


Hence, in the ellipse and hyperbola, every ordinate has two determinate abscisses; but in the parabola, only one; the other vertex of the diameter being infinitely distant.

16. The Parameter of any diameter, is a third proportional to that diameter and its conjugate.

17. The

17. The Focus is the point in the axis where the ordinate is equal to half the parameter. As k and l , where dk or el is equal to the semi-parameter. The name focus being given to this point from the peculiar property of it mentioned in the corol. to theor. 9 in the Ellipse and Hyperbola following, and to theor. 6 in the Parabola.

Hence, the ellipse and hyperbola have each two foci ; but the parabola only one.



18. If DAE , FBG , be two opposite hyperbolas, having AB for their first or transverse axis, and ab for their second or conjugate axis. And if dae , fbg , be two other opposite hyperbolas having the same axes, but in the contrary order, namely, ab their first axis, and AB their second ; then these two latter curves dae , fbg , are called the conjugate hyperbolas to the two former DAE , FBG , and each pair of opposite curves mutually conjugate to the other.

19. And if tangents be drawn to the four vertices of the curves, or extremities of the axes, forming the inscribed rectangle $HIKL$; the diagonals HCK , IL , of this rectangle, are called the asymptotes of the curves. And if these asymptotes intersect at right angles, or the inscribed rectangle be a square, or the two axes AB and ab be equal, then the hyperbolas are said to be right-angled, or equilateral.

SCHOLIUM.


The rectangle inscribed between the four conjugate hyperbolas, is similar to a rectangle circumscribed about an ellipse, by drawing tangents, in like manner, to the four extremities of the two axes ; and the asymptotes or diagonals in the hyperbola, are analogous to those in the ellipse, cutting this curve in similar points, and making that pair of conjugate diameters which are equal to each other. Also, the whole figure formed by the four hyperbolas, is, as it were, an ellipse turned inside out, cut open at the extremities D , E , F , G , of the said equal conjugate diameters, and those four points drawn out to an infinite distance ; the curvature being turned the contrary way, but the axes, and the rectangle passing through their extremities, continuing fixed.

OF THE ELLIPSE.

THEOREM I.

The squares of the Ordinates of the Axis are to each other as the Rectangles of their Abscisses.

LET AVB be a plane passing through the axis of the cone; AGIH another section of the cone perpendicular to the plane of the former; AB the axis of this elliptic section; and FG, HI, ordinates perpendicular to it. Then it will be, as $FG^2 : HI^2 :: AF \cdot FB : AH \cdot HB$.



For, through the ordinates FG, HI, draw the circular sections KGL, MIN, parallel to the base of the cone, having KL, MN, for their diameters, to which FG, HI, are ordinates, as well as to the axis of the ellipse.

Now, by the similar triangles AFL, AHN, and BFK, BHM,

$$\begin{aligned} \text{it is } AF : AH &:: FL : HN, \\ \text{and } FB : HB &:: KF : MH; \end{aligned}$$


hence, taking the rectangles of the corresponding terms, it is, the rect. $AF \cdot FB : AH \cdot HB :: KF \cdot FL : MH \cdot HN$.

But, by the circle, $KF \cdot FL = FG^2$, and $MH \cdot HN = HI^2$; Therefore the rect. $AF \cdot FB : AH \cdot HB :: FG^2 : HI^2$. Q. E. D.

THEOREM II.

As the Square of the Transverse Axis :
Is to the square of the Conjugate ::
So is the Rectangle of the Abscisses :
To the Square of their Ordinate.

That is, $AB^2 : ab^2$ or
 $AC^2 : ac^2 :: AD \cdot DB : DE^2$.



For, by theor. 1, $AC \cdot CB : AD \cdot DB :: ca^2 : DE^2$;
 But, if c be the centre, then $AC \cdot CB = AC^2$, and ca is the
 semi-conjugate.

Therefore $AC^2 : AD \cdot DB :: ac^2 : DE^2$;
 or, by permutation, $AC^2 : ac^2 :: AD \cdot DB : DE^2$;
 or, by quadrupling, $AB^2 : ab^2 :: AD \cdot DB : DE^2$. Q. E. D.

Corol. Or by div. $AB : \frac{ab^2}{AB} :: AD \cdot DB$ or $CA^2 - CD^2 : DE^2$, Q. E. D.


that is, $AB : p :: AD \cdot DB$ or $CA^2 - CD^2 : DE^2$;

where p is the parameter $\frac{ab^2}{AB}$, by the definition of it.

That is, As the transverse,
 Is to its parameter,
 So is the rectangle of the abscisses,
 To the square of their ordinate.

THEOREM III.

As the Square of the Conjugate Axis :
 Is to the Square of the Transverse Axis ::
 So is the Rectangle of the Abscisses of the Conjugate, or
 the Difference of the Squares of the Semi-conjugate and
 Distance of the Centre from any Ordinate of that Axis :
 To the Square of their Ordinate.



That is,
 $ca^2 : cb^2 :: ad \cdot db$ or $ca^2 - cd^2 : de^2$.

For, draw the ordinate ED to the transverse AB.
 Then, by theor. 2, $ca^2 : CA^2 :: DE^2 : AD \cdot DB$ or $CA^2 - CD^2$;
 or - - - - - $ca^2 : CA^2 :: cd^2 : CA^2 - de^2$.
 But - - - - - $ca^2 : CA^2 :: ca^2 : CA^2$;
 theref. by subtr. - $ca^2 : CA^2 :: ca^2 - cd^2$ or $ad \cdot db : de^2$.

Q. E. D.

Corol.

Corol. 1. If two circles be described on the two axes as diameters, the one inscribed within the ellipse, and the other circumscribed about it; then an ordinate in the circle will be to the corresponding ordinate in the ellipse, as the axis of this ordinate, is to the other axis.

That is, $CA : ca :: DG : DE,$
and $ca : CA :: dg : de.$

For, by the nature of the circle, $AD \cdot DB = DG^2$; theref. by the nature of the ellipse, $CA^2 : ca^2 :: AD \cdot DB$ or $DG^2 : DE^2,$
or $CA : CA :: DG : DE.$

In like manner - - $ca : CA :: dg : de.$

Also, by equality, - - $DG : DE$ or $cd :: de$ or $DC : dg.$

Therefore cgg is a continued straight line.


Corol. 2. Hence also, as the ellipse and circle are made up of the same number of corresponding ordinates, which are all in the same proportion of the two axes, it follows that the areas of the whole circle and ellipse, as also of any like parts of them, are in the same proportion of the two axes, or as the square of the diameter to the rectangle of the two axes; that is, the areas of the two circles, and of the ellipse, are as the square of each axis and the rectangle of the two; and therefore the ellipse is a mean proportional between the two circles.

THEOREM IV.

The Square of the Distance of the Focus from the Centre, is equal to the Difference of the Squares of the Semi-axes;

Or, the Square of the Distance between the Foci, is equal to the Difference of the Squares of the two Axes.

That is, $CF^2 = CA^2 - ca^2,$
or $Ff^2 = AB^2 - ab^2.$



For, to the focus F draw the ordinate FE ; which, by the definition, will be the semi-parameter. Then, by the nature of the curve - - $CA^2 : ca^2 :: CA^2 - CF^2 : FE^2$; and by the def. of the para. $CA^2 : ca^2 :: ca^2 : FE^2$; therefore - - $ca^2 = CA^2 - CF^2$; and by addit. and subtr. $CF^2 = CA^2 - ca^2$; or, by doubling, - - $Ff^2 = AB^2 - ab^2$;


Q. E. D.
Corol.

Corol. 1. Hence CI or $CA - FE$ is a 4th proportional to CA , CF , CD .

Corol. 2. And $fe - FE = 2CI$; that is, the difference between two lines drawn from the foci, to any point in the curve, is double the 4th proportional to CA , CF , CD .

Corol. 3. Hence is derived the common method of describing this curve mechanically by points, or with a thread thus:


In the transverse take the foci F, f , and any point I . Then with the radii AI, BI , and centres F, f , describe arcs intersecting in E , which will be a point in the curve. In like manner, assuming other points I , as many other points will be found in the curve. Then with a steady hand the curve line may be drawn through all the points of intersection E .



Or, take a thread of the length AB of the transverse axis, and fix its two ends in the foci F, f , by two pins. Then carry a pen or pencil round by the thread, keeping it always stretched, and its point will trace out the curve line.

THEOREM VI.

If from any Point I in the Axis produced, a Line IL be drawn touching the curve in one point L ; and the Ordinate LM be drawn; and if c be the Centre or Middle of AB : Then shall CM be to CI as the Square of AM to the Square of AI .



That is,
 $CM : CI :: AM^2 : AI^2$.

For, from the point I draw any other line IEH to cut the curve in two points E and H ; from which let fall the perpendiculars ED and HG ; and bisect DG in K .

Then, by theo. 1, $AD \cdot DB : AG \cdot GB :: DE^2 : GH^2$,
 and by sim. triangles, $ID^2 : IG^2 :: DE^2 : GH^2$;
 theref. by equality, $AD \cdot DB : AG \cdot GB :: ID^2 : IG^2$.

But $DB = CB + CD = AC + CD = AG + DC - CG = 2CK + AG$,
 and $GB = CB - CG = AC - CG = AD + DC - CG = 2CK + AD$;
 theref. $AD \cdot 2CK + AD \cdot AG : AG \cdot 2CK + AD \cdot AG :: ID^2 : IG^2$,
 and, by div. $DG \cdot 2CK : IG^2 - ID^2$ or $DG \cdot 2IK :: AD \cdot 2CK + AD \cdot AG : ID^2$,
or


or $2CK : 2IK :: AD \cdot 2CK + AD \cdot AG : ID^2$,
 or $AD \cdot 2CK : AD \cdot 2IK :: AD : 2CK + AD \cdot AG : ID^2$;
 theref. by div. $CK : IK :: AD \cdot AG : ID^2 - AD \cdot 2IK$,
 and, by comp. $CK : IC :: AD \cdot AG : ID^2 - AD \cdot ID + IA$,
 or $CK : CI :: AD \cdot AG : AI^2$.

But, when the line IH , by revolving about the point I , comes into the position of the tangent IL , then the points E and H meet in the point L , and the points D, K, G , coincide with the point M ; and then the last proportion becomes $CM : CI :: AM^2 : AI^2$. Q. E. D.

THEOREM VII. *17 2/11/7*

If a Tangent and Ordinate be drawn from any Point in the Curve, meeting the Transverse Axis; the Semi-transverse will be a Mean Proportional between the Distances of the said Two Intersections from the Centre.

That is,
 CA is a mean proportional between CD , and CT ;
 or CD, CA, CT , are continued proportionals.



For, by theor. 6, $CD : CT :: AD^2 : AT^2$.
 that is, $CD : CT :: (CA - CD)^2 : (CT - CA)^2$,
 or $CD : CT :: CD^2 + CA^2 : CA^2 + CT^2$,
 and $CD : DT :: CD^2 + CA^2 : CT^2 - CD^2$,
 or $CD : DT :: CD^2 + CA^2 : (CT + CD) \cdot DT$,
 or $CD^2 : CD \cdot DT :: CD^2 + CA^2 : CD \cdot DT + CT \cdot DT$,
 hence $CD^2 : CA^2 :: CD \cdot DT : CT \cdot DT$,
 and $CD^2 : CA^2 :: CD : CT$.
 therefore (th. 78, Geom.) $CD : CA :: CA : CT$. Q. E. D.


Corol. Since CT is always a third proportional to CD, CA ; if the points D, A , remain constant, then will the point T be constant also; and therefore all the tangents will meet in this point T , which are drawn from the point E , of every ellipse described on the same axis AB , where they are cut by the common ordinate DEE drawn from the point D .

THEOREM

THEOREM VIII.

If there be any Tangent meeting Four Perpendiculars to the Axis drawn from these four Points, namely, the Centre, the two Extremities of the Axis, and the Point of Contact; those Four Perpendiculars will be Proportionals.

That is,
 $AG : DE :: CH : BI.$



For, by theor. 7, $TC : AC :: AC : DC$,
 theref. by div. $TA : AD :: TC : AC$ or CB ,
 and by comp. $TA : TD :: TC : TB$,
 and by sim. tri. $AG : DE :: CH : BI.$ Q. E. D.


Corol. Hence TA, TD, TC, TB } are also proportionals.
 and TG, TE, TH, TI }

For these are as AG, DE, CH, BI , by similar triangles.

THEOREM IX.

If there be any Tangent, and two Lines drawn from the Foci to the Point of Contact; these two Lines will make equal Angles with the Tangent.

That is,
 the $\angle FET = \angle fee.$



For, draw the ordinate DE , and fe parallel to FE .

By cor. 1, theor. 5, $CA : CD :: CF : CA - FE$,
 and by theor. 7, $CA : CD :: CT : CA$;
 therefore $CT : CF :: CA : CA - FE$;
 and by add. and sub. $TF : Tf :: FE : 2CA - FE$ or fe by th. 5.
 But by simp. tri. $TF : Tf :: FE : fe$;
 therefore $fe = fe$, and conseq. $\angle e = \angle fee.$
 But, because FE is parallel to fe , the $\angle e = \angle FET$;
 therefore the $\angle FET = \angle fee.$ Q. E. D.


Corol.

Corol. As opticians find that the angle of incidence is equal to the angle of reflection, it appears from this theorem, that rays of light issuing from the one focus, and meeting the curve in every point, will be reflected into lines drawn from those points to the other focus. So the ray FE is reflected into FE . And this is the reason why the points F, f , are called the *foci*, or burning points.

THEOREM X.

All the Parallelograms circumscribed about an Ellipse are equal to one another, and each equal to the Rectangle of the two Axes.

That is,
the parallelogram $PQRS$ =
the rectangle $AB \cdot ab$.



Let EG, eg , be two conjugate diameters parallel to the sides of the parallelogram, and dividing it into four less and equal parallelograms. Also, draw the ordinates DE, de , and CK perpendicular to PQ ; and let the axis CA produced meet the sides of the parallelogram, produced if necessary, in T and t .

Then, by theor. 7,

$$CT : CA :: CA : CD,$$

and

$$ct : CA :: CA : cd;$$

theref. by equality,

$$CT : ct :: cd : CD;$$

but, by sim. triangles,

$$CT : ct :: TD : CD,$$

theref. by equality, †

$$TD : cd :: CD : CD,$$

and the rectangle,

$$TD \cdot DC \text{ is } = \text{the square } cd^2.$$

Again, by theor. 7,

$$CD : CA :: CA : CT,$$

or, by division,

$$CD : CA :: DA : AT,$$

and by composition,

$$CD : DB :: AD : DT;$$

conseq. the rectangle

$$CD \cdot DT = cd^2 = AD \cdot DB^*.$$

But, by theor. 2,

$$CA^2 : ca^2 :: (AD \cdot DB \text{ or }) cd^2 : DE^2,$$

therefore

$$CA : ca :: cd : DE;$$

* *Corol.* Because $cd^2 = AD \cdot DB = CA^2 - CD^2$,

therefore $CA^2 = CD^2 + cd^2$,


In like manner, $ca^2 = DE^2 + de^2$.

In like manner, $CA : ca :: CD : de,$
 or $ca : de :: CA : CD.$
 But, by theor. 7, $CT : CA :: CA : CD ;$
 theref. by equality, $CT : CA :: ca : de.$
 But, by sim. tri. $CT : CK :: ce : de ;$
 theref. by equality, $CK : CA :: ca : ce,$
 and the rectangle $CK . ce = CA . ca.$
 But the rect. $CK . ce =$ the parallelogram $CEPE,$
 theref. the rect. $CA . ca =$ the parallelogram $CEPE,$
 conseq. the rect. $AB . ab =$ the parallelogram $PQRS. \quad Q. E. D.$

THEOREM XI.

The Sum of the Squares of every Pair of Conjugate Diameters, is equal to the same constant Quantity, namely, the Sum of the Squares of the two Axes.

That is,
 $AB^2 + ab^2 = EG^2 + eg^2 ;$
 where $EG, eg,$ are any pair of conjugate diameters.




For, draw the ordinates $ED, eb.$
 Then, by cor. to theor. 10, $CA^2 = CD^2 + cd^2,$
 and $ca^2 = DE^2 + de^2 ;$
 therefore the sum $CA^2 + ca^2 = CD^2 + DE^2 + cd^2 + de^2.$
 But, by right-angled $\Delta s,$ $CE^2 = CD^2 + DE^2,$
 and $ce^2 = cd^2 + de^2 ;$
 therefore the sum $CE^2 + ce^2 = CD^2 + DE^2 + cd^2 + de^2.$
 consequently $CA^2 + ca^2 = CE^2 + ce^2 ;$
 or, by doubling, $AB^2 + ab^2 = EG^2 + eg^2. \quad Q. E. D.$

THEOREM XII.

The difference between the Semi-transverse and a Line drawn from the Focus to any point in the Curve, is equal to a Fourth Proportional to the Semi-transverse, the distance from the Centre to the Focus, and Distance from the Centre to the Ordinate belonging to that Point of the Curve.

That is,
 $AC - FE = CI$, or $FE = AI$;
 and $fE - AC = CI$, or $fE = BI$.
 Where $CA : CF :: CD : CI$ the 4th
 proportional to CA, CF, CD .




For, draw AG parallel and equal to ca the semi-conjugate ;
 and join CG meeting the ordinate DE in H .
 Then, by theor. 2 $CA^2 : AG^2 :: CA^2 - CD^2 : DE^2$;
 and, by sim. tri. $CA^2 : AG^2 :: CA^2 - CD^2 : AG^2 - DH$;
 consequently $DE^2 = AG^2 - DH^2 = ca^2 - DH^2$.
 Also $FD = CF \oslash CD$, and $FD^2 = CF^2 - 2CF \cdot CD + CD^2$;
 but by right-angled triangles, $FD^2 + DE^2 = FE^2$;
 therefore $FE^2 = CF^2 + ca^2 - 2CF \cdot CD + CD^2 - DH^2$.
 But by theor. 4, $ca^2 + CF = CA^2$;
 and, by supposition, $2CF \cdot CD = 2CA \cdot CI$;
 theref. $FE^2 = CA^2 - 2CA \cdot CI + CD^2 - DH^2$;
 But by supposition, $CA^2 : CD^2 :: CF^2$ or $CA^2 - AG^2 : CI^2$;
 and, by sim. tri. $CA^2 : CD^2 :: CA^2 - AG^2 : CD^2 - DH^2$;
 therefore - - - $CI^2 = CD^2 - DH^2$;
 consequently - - - $FE^2 = CA^2 - 2CA \cdot CI + CI^2$.
 And the root or side of this square is $FE = CA - CI = AI$.
 In the same manner is found $fE = CA + CI = BI$. Q. E. D.

THEOREM XIII.

If a Line be drawn from either Focus, Perpendicular to a
 Tangent to any Point of the curve ; the Distance of their
 Intersection from the Centre will be equal to the Semi-
 transverse Axis.

That is, if FP, fp
 be perpendicular to
 the tangent TPP ,
 then shall CP and cp
 be each equal to CA
 or CB .



For,

For, through the point of contact E draw FE , and fE meeting FP produced in G . Then, the $\angle GEP = \angle FEP$, being each equal to the $\angle fep$, and the angles at P being right, and the side PE being common, the two triangles GEP , FEP are equal in all respects, and so $GE = FE$, and $GP = FP$. Therefore, since $FP = \frac{1}{2}FG$, and $FC = \frac{1}{2}fG$, and the angle at F common, the side CP will be $= \frac{1}{2}fG$ or $\frac{1}{2}AB$, that is $CP = CA$ or CB . And in the same manner $cp = CA$ or CB . Q. E. D.

Corol. 1. A circle described on the transverse axis, as a diameter, will pass through the points P, p ; because all the lines CA, CP, cp, CB , being equal, will be radii of the circle.


Corol. 2. CP is parallel to fE , and cp parallel to FE .

Corol. 3. If at the intersections of any tangent, with the circumscribed circle, perpendiculars to the tangent be drawn, they will meet the transverse axis in the two foci. That is, the perpendiculars PF, pf give the foci F, f .

THEOREM XIV.

The equal Ordinates, or the Ordinates at equal Distances from the Centre, on the opposite Sides and Ends of an Ellipse, have their Extremities connected by one Right Line passing through the Centre, and that Line is bisected by the Centre.

That is, if $CD = CG$, or the ordinate $DE = GH$; then shall $CE = CH$, and ECH will be a right line.



For, when $CD = CG$, then also is $DE = GH$ by cor. 2, th. 1. But the $\angle D = \angle G$, being both right angles; therefore the third side $CE = CH$, and the $\angle DCE = \angle GCH$, and consequently ECH is a right line.

Corol.

Corol. 1. And, conversely. if ech be a right line passing through the centre; then shall it be bisected by the centre, or have $ce=ch$; also de will be $=gh$, and $cd=cg$.


Corol. 2. Hence also. if two tangents be drawn to the two ends, e, h of any diameter eh ; they will be parallel to each other, and will cut the axis at equal angles, and at equal distances from the centre. For, the two cd, ca being equal to the two cg, cb , the third proportionals ct, cs will be equal also; then the two sides ce, ct being equal to the two ch, cs , and the included angle ect equal to the included angle hcs , all the other corresponding parts are equal: and so the $\angle T = \angle s$, and te parallel to hs .

Corol. 3. And hence the four tangents, at the four extremities of any two conjugate diameters form a parallelogram circumscribing the ellipse, and the pairs of opposite sides are each equal to the corresponding parallel conjugate diameters. For, if the diameter eh be drawn parallel to the tangent te or hs , it will be the conjugate to eh by the definition; and the tangents to e, h will be parallel to each other, and to the diameter eh for the same reason.

THEOREM XV.

If two Ordinates ED, ed be drawn from the Extremities E, e of two Conjugate Diameters, and Tangents be drawn to the same Extremities, and meeting the Axis produced in T and R ;

Then shall cd be a mean proportional between cd, dr ,
and cd a mean proportional between cd, dt .



For, by theor. 7,
and by the same,
theref. by equality,
But by sim. tri.
theref. by equality,
In like manner,

$$\begin{aligned} CD : CA &:: CA : CT, \\ cd : CA &:: CA : CR ; \\ CD : cd &:: CR : CT, \\ DT : cd &:: CT : CR ; \\ CD : cd &:: cd : DT. \\ cd : CD &:: CD : dr. \end{aligned}$$

Q. E. D.
Corol.

Corol. 1. Hence $CD : cd :: CR : CT$.

Corol. 2. Hence also $CD : cd :: de : DE$.

And the rectangle $CD \cdot DE = cd \cdot de$, or $\triangle CDE = \triangle cde$.

Corol. 3. Also $cd^2 = CD \cdot DT$,

and $CD^2 = cd \cdot dr$.

Or cd a mean proportional between CD, DT ;
and CD a mean proportional between cd, dr .

THEOREM XVI.

The same Figure being constructed as in the last Theorem, each Ordinate will divide the Axis, and the semi-axis added to the external Part, in the same Ratio.

[See the last fig.]

That is, $DA : DT :: DC : DB$,

and $dA : dr :: dc : dB$.

For, by theor. 7, $CD : CA :: CA : CT$,

and by div. $CD : CA :: AD : AT$,

and by comp. $CD : DB :: AD : DT$,

or, - - - - - $DA : DT :: DC : DB$.

In like manner, $dA : dr :: dc : dB$.

Q. E. D.

Corol. 1. Hence, and from cor. 3 to the last, it is,

$cd^2 = CD \cdot DT = AD \cdot DB = CA^2 - CD^2$, *3^d Prop.*

$CD^2 = CD \cdot dr = Ad \cdot dB = CA^2 - cd^2$.

Corol. 2. Hence also, $CA^2 = CD^2 + cd^2$,

and $ca^2 = DE^2 + de^2$.

Corol. 3. Further, because $CA^2 : ca^2 :: AD \cdot DB$ or $cd^2 : DE^2$,

therefore $CA : ca :: cd : DE$.

likewise $CA : CA :: CD : de$.

THEOREM XVII.


If from any Point in the Curve there be drawn an Ordinate, and a Perpendicular to the Curve, or to the Tangent at that point : Then, the

Dist. on the Trans. between the Centre and Ordinate, CD :

Will be to the Dist. PD ::

As sq. of the Trans. Axis :

To sq. of the Conjugate.



That is,

$CA^2 : ca^2 :: DC : DP$.


For,

For, by theor. 2, $CA^2 : ca^2 :: AD \cdot DB : DE^2$;
 But, by rt. angled Δ s, the rect. $TD \cdot DP = DE^2$;
 and, by cor. 1, theor. 16. $CD \cdot DT = AD \cdot DB$;
 therefore - - $CA^2 : ca^2 :: TD \cdot DC : TD \cdot DP$,
 or - - - - $AC^2 : ca^2 :: DC : DP$. Q. E. D.

THEOREM XVIII.

If there be Two Tangents drawn, the One to the Extremity of the Transverse, and the other to the Extremity of any other diameter, each meeting the other's diameter produced; the two Tangential Triangles so formed will be equal.

That is,
 the triangle $CET =$ the
 triangle CAN .



For, draw the ordinate DE . Then
 By sim. triangles, $CD : CA :: CE : CN$?
 but, by theor. 7, $CD : CA :: CA : CT$;
 theref. by equal. $CA : CT :: CE : CN$.

The two triangles CET, CAN have then the angle c common, and the sides about that angle reciprocally proportional; those triangles are therefore equal, namely, the $\Delta CET = \Delta CAN$.


Corol. 1. From each of the equal tri. CET, CAN , take the common space $CAPE$, and there remains the external $\Delta PAT = \Delta PNE$.

Corol. 2. Also from the equal triangles CET, CAN , take the common triangle CED , and there remains the $\Delta TED =$ trapez. $ANED$.

THEOREM XIX.

The same being supposed as in the last Proposition; then any Lines kq, qg , drawn parallel to the two Tangents, shall also cut off equal Spaces. That is,

ΔKQG



$\Delta KQG = \text{trapez. ANHG}$,
and $\Delta Kqg = \text{trapez. ANhg}$.

For draw the ordinate DE. Then

The three sim. triangles CAN, CDE, CGH,
are to each other as CA^2, CD^2, CG^2 ;

th. by div. the trap. ANED : trap. ANHG :: $CA^2 - CD^2 : CA^2 - CG^2$.

But, by theor. 1, $DE^2 : GQ^2 :: CA^2 - CD^2 : CA^2 - CG^2$,

theref. by equ. trap. ANED : trap. ANHG :: $DE^2 : GQ^2$.

but, by sim. Δ s, tri. TED : tri. KQG :: $DE^2 : GQ^2$;

theref. by equality, ANED : TED :: ANHG : KQG.

But, by cor. 2, theor. 18, the trap. ANED = Δ TED ;

and therefore the trap. ANHG = Δ KQG.

In like manner the trap. ANhg = Δ Kqg. Q. E. D.

Corol. 1. The three spaces ANHG, TEHG, KQG are all equal.

Corol. 2. From the equals ANHG, KQG,

take the equals ANhg, Kqg.

and there remains ghHG = gqQG.

Corol. 3. And from the equals ghHG, gqQG,

take the common space gqLHG,


and there remains the Δ LQH = Δ Lqh.

Corol. 4. Again from the equals KQG, TEHG,

take the common space KLHG,

and there remains TELK = Δ LQH.

Corol. 5. And when,
by the lines KQ, GH,
moving with a parallel
motion, KQ comes into
the position IR, where
CR is the conjugate to
CA ; then



the triangle KQG becomes the triangle IRC,
and the space ANHG becomes the triangle ANC ;
and therefore the Δ IRC = Δ ANC = Δ TEC.

Corol. 6. Also when the lines KQ and HQ, by moving with a
parallel motion, come into the position ce, me,


the

the triangle LQH becomes the triangle cem ,
and the space $TELK$ becomes the triangle TEC ;
and theref. the $\triangle cem = \triangle TEC = \triangle ANC = \triangle IRC$.

THEOREM XX.

Any Diameter bisects all its Double Ordinates, or the Lines drawn Parallel to the Tangent at its Vertex, or to its Conjugate Diameter.

That is, if qg be parallel to the tangent TE , or to ce , then shall $LQ = Lq$.



For, draw qg perpendicular to the transverse. Then by cor. 3, theor. 19, the $\triangle LQH = \triangle Lqh$; but these triangles are also equiangular; consequently their like sides are equal, or $LQ = Lq$.

Corol. Any diameter divides the ellipse into two equal parts.

For, the ordinates on each side being equal to each other, and equal in number; all the ordinates, or the area, on one side of the diameter, is equal to all the ordinates, or the area, on the other side of it.


THEOREM XXI.

As the Square of any Diameter :
Is to the Square of its Conjugate : :
So is the Rectangle of any two Abscisses :
To the Square of their Ordinate.

That is, $CE^2 : ce^2 :: EL \cdot LG$ or $CE^2 - CL^2 : LQ^2$.

For, draw the tangent TE , and produce the ordinate QL to the transverse at K . Also draw QH , em perpendicular to the transverse, and meeting EG in H and m .

Then similar triangles being as the squares of their like sides, it is,



by

by sim. triangles, $\triangle CET : \triangle CLK :: CE^2 : CL^2$;
 or, by division, $\triangle CET : \text{trap. TELK} :: CE^2 : CE^2 - CL^2$.
 Again, by sim. tri. $\triangle CEM : \triangle LQH :: ce^2 : LQ^2$.

But, by cor. 5, theor. 19, the $\triangle CEM = \triangle CET$,
 and, by cor. 4, theor. 19, the $\triangle LQH = \text{trap. TELK}$;
 theref. by equality, $CE^2 : ce^2 :: CE^2 - CL^2 : LQ^2$,
 OR - - - $CE^2 : ce^2 :: EL . LG : LQ^2$. Q. E. D.

Corol. 1. The squares of the ordinates to any diameter, are to one another as the rectangles of their respective abscisses, or as the difference of the squares of the semidiameter and of the distance between the ordinate and centre. For they are all in the same ratio of CE^2 to ce^2 .


Corol. 2. The above being the same property as that belonging to the two axes, all the other properties before laid down for the axes, may be understood of any two conjugate diameters whatever using only the oblique ordinates of these diameters, instead of the perpendicular ordinates of the axes ; namely, all the properties in theorems 6, 7, 8, 14, 15, 16, 18 and 19.

THEOREM XXII.

If any Two lines, that any where intersect each other, meet the Curve each in Two Points ; then

The Rectangle of the Segments of the one :
 Is to the Rectangle of the Segments of the other : :
 As the Square of the Diam. Parallel to the former :
 To the Square of the Diam. Parallel to the latter.

That is, if CR and cr, be Parallel to any two Lines PHQ, pHq ; then shall
 $CR^2 : cr^2 :: PH . HQ : PH . Hq$.



For, draw the diameter CHE, and the tangent TE, and its parallels PK, RI, MH, meeting the conjugate of the diameter CR in the points T, K, I, M. Then, because similar triangles are as the squares of their like sides, it is,

by sim. triangles, $CR^2 : GP^2 :: \triangle CRI : \triangle GPK$,
 and - - - $CR^2 : GH^2 :: \triangle CRI : \triangle GHM$;
 theref by division, $CR^2 : GP^2 - GH^2 :: CRI : KPHM$.
 Again, by sim. tri. $CE^2 : CH^2 :: \triangle CTE : \triangle CMH$;
 and by division, $CE^2 : CE^2 - CH^2 :: \triangle CTE : TEHM$.

But, by cor. 5, theor. 19, the $\triangle CTE = \triangle CIR$,
 and by cor. 1, theor. 19, $TEHC = KPHG$, or $TEHM = KPHM$;
 theref. by equ. $CE^2 : CE^2 - CH^2 :: CR^2 : GP^2 - GH^2$ or $PH : HQ$.
 In like manner $CE^2 : CE^2 - CH^2 :: cr^2 : ph \cdot hq$.
 Theref. by equ. $CR^2 : cr^2 :: PH \cdot HQ : ph \cdot hq$. Q. E. D.


Corol. 1. In like manner, if any other lines $p'h'q'$, parallel to cr or to pq , meet PHQ ; since the rectangles PHQ , $p'h'q'$ are also in the same ratio of CR^2 to cr^2 ; therefore rect. $PHQ : phq :: PH'Q : p'h'q'$.

Also, if another line $f'hq'$ be drawn parallel to pq or CR ; because the rectangles $f'hq'$ $p'hq'$ are still in the same ratio, therefore, in general, the rect. $PHQ : phq :: f'hq' : p'hq'$.

That is, the rectangles of the parts of two parallel lines, are to one another, as the rectangles of the parts of two other parallel lines, any where intersecting the former.

Corol. 2. And when any of the lines only touch the curve, instead of cutting it, the rectangles of such become squares, and the general property still attends them.

That is,
 $CR^2 : cr^2 :: TE^2 : te^2$,
 or $CR : cr :: TE : te$.
 and $CR : cr :: TE : te$.




Corol. 3. And hence $TE : te :: TE : te$.

OF THE HYPERBOLA.

THEOREM I.

The Squares of the Ordinates of the Axis are to each other as the Rectangles of their Abscisses.

LET AVB be a plane passing through the vertex and axis of the opposite cones; $AGIH$ another section of them perpendicular to the plane of the former; AB the axis of the hyperbolic sections; and FG, HI , ordinates perpendicular to it. Then it will be, as $FG^2 : HI^2 :: AF \cdot FB : AH \cdot HB$.



For, through the ordinates FG, HI , draw the circular sections KGL, MIN , parallel to the base of the cone, having KL, MN , for their diameters, to which FG, HI , are ordinates, as well to the axis of the hyperbola.

Now, by the similar triangles AFL, AHN , and BFK, BHM ,

$$\text{it is } AF : AH :: FL : HN,$$

$$\text{and } EB : HB :: KF : MH;$$

hence, taking the rectangles of the corresponding terms, it is, the rect. $AF \cdot FB : AH \cdot HB :: KF \cdot FL : MH \cdot HN$.


But, by the circle, $KF \cdot FL = FG^2$, and $MH \cdot HN = HI^2$;

Therefore the rect. $AF \cdot FB : AH \cdot HB :: FG^2 : HI^2$. Q. E. D.

THEOREM II.

As the Square of the Transverse Axis :
Is to the square of the Conjugate ::
So is the Rectangle of the Abscisses :
To the Square of their Ordinate.

That is, $AB^2 : ab^2$ or
 $AC^2 : ac^2 :: AD \cdot DB : DE^2$.



For

For, by theor. 1, $AC \cdot CB : AD \cdot DB :: ca^2 : DE^2$;
 But, if c be the centre, then $AC \cdot CB = AC^2$, and ca is the semi-conj.

Therefore $AC^2 : AD \cdot DB :: ac^2 : DE^2$;
 or, by permutation, $AC^2 : ac^2 :: AD \cdot DB : DE^2$;
 or, by doubling, $AB^2 : ab^2 :: AD \cdot DB : DE^2$. Q. E. D.

Corol. Or by div. $AB : \frac{ab^2}{AB} :: AD \cdot DB \text{ OR } CD^2 - CA^2 : DE^2$,

that is, $AB : p :: AD \cdot DB \text{ OR } CD^2 - CA^2 : DE^2$;


where p is the parameter $\frac{ab^2}{AB}$, by the definition of it.

That is, As the transverse,
 Is to its parameter,
 So is the rectangle of the abscisses,
 To the square of their ordinate.

THEOREM III.

As the Square of the Conjugate Axis :
 To the Square of the Transverse Axis : :
 The Sum of the Squares of the Semi-conjugate, and
 Distance of the Centre from any Ordinate of the Axis :
 The Square of their Ordinate.

That is,
 $ca^2 : CA^2 :: ca^2 + cd^2 : de^2$.



For, draw the ordinate ED to the transverse AB .

Then, by theor. 1, $ca^2 : CA^2 :: DE^2 : AD \cdot DB \text{ OR } CD^2 - CA^2$,
 or - - - - - $ca^2 : CA^2 :: cd^2 : de^2 - CA^2$.
 But - - - - - $ca^2 : CA^2 :: ca^2 : CA^2$,
 theref. by compos. $ca^2 : CA^2 :: ca^2 + cd^2 : de^2$.
 In like manner, $CA^2 : CA^2 :: CA^2 + CD^2 : de^2$. Q. E. D.


Corol. By the last theor. $ca^2 : CA^2 :: CD^2 - CA^2 : DE^2$,
 and by this theor. $CA^2 : CA^2 :: CD^2 + CA^2 : de^2$,
 therefore - - $DE^2 : de^2 :: CD^2 - CA^2 : CD^2 + CA^2$.
 In like manner, $de^2 : de^2 :: cd^2 - ca^2 : cd^2 + ca^2$.

THEOREM

THEOREM IV.

The Square of the Distance of the Focus from the Centre, is equal to the Sum of the Squares of the Semi-axes.
 Or, the Square of the Distance between the Foci, is equal to the Sum of the Squares of the two Axes.

That is,
 $CF^2 = CA^2 + ca^2$, OR
 $Ff^2 = AB^2 + ab^2$.



For, to the focus F draw the ordinate FE ; which, by the definition, will be the semi-parameter. Then, by the nature of the curve $CA^2 : ca^2 :: CF^2 - CA^2 : FE^2$, and by the def. of the para $CA^2 : ca^2 :: ca^2 : FE^2$; therefore $ca^2 = CF^2 - CA^2$; and by addition, $CF^2 = CA^2 + ca^2$; or, by doubling, $Ff^2 = AB^2 + ab^2$; Q. E. D.

Corol. 1 The two semi-axes, and the focal distance from the centre, are the sides of a right-angled triangle *CAA*; and the distance *AA* is = *CF* the focal distance.


Corol. 2 The conjugate semi-axes, *ca*, is a mean proportional between *AF*, *FB*. or between *af*, *fB*, the distances of either focus from the two vertices.

For $ca^2 = CF^2 - CA^2 = CF + CA \cdot CF - CA = AF \cdot FB$.

THEOREM V.

The Difference of two Lines drawn from the two Foci, to meet at any Point in the Curve, is equal to the Transverse Axis.

That is,
 $fE - FE = AB$.



For, draw *AG* parallel and equal to *ca* the semi-conjugate ; and join *CG* meeting the ordinate *DE* produced in *H* ; also take *CI* a 4th proportional to *CA*, *CF*, *CD*.

Then,

Then, by th. 2, $CA^2 : AG^2 :: CD^2 - CA^2 : DE^2$;
 and by sim. Δ s, $CA^2 : AG^2 :: CD^2 - CA^2 : DH^2 - AG$;
 consequently, $DE^2 = DH^2 - AG^2 = DH^2 - CA^2$.

Also, $FD = CF \oslash CD$, and $FD^2 = CF^2 - 2CF \cdot CD + CD^2$;
 and, by right-angled triangles, $FE^2 = FD^2 + DE^2$;
 therefore $FE^2 = CF^2 - CA^2 - 2CF \cdot CD + CD^2 + DH^2$.

But, by theor 4, $CF^2 - CA^2 = CA^2$,
 and by supposition, $2CF \cdot CD = 2CA \cdot CI$;
 therel. $FE^2 = CA^2 - 2CA \cdot CI + CD^2 + DH^2$.

Again, by suppos. $CA^2 : CD^2 :: CF^2$ or $CA^2 + AG^2 \cdot CI^2$;
 and, by sim. tri. $CA^2 : CD^2 :: CA^2 + AG^2 : CD^2 + DH^2$;
 therefore - - - $CI^2 = CD^2 + DH^2 = CH^2$;
 consequently $FE^2 = CA^2 - 2CA \cdot CI + CI^2$.

And the root or side of this square is $FE = CI - CA = AI$.

In the same manner it is found that $fe = CI + CA = BI$.

Conseq. by subtract. $fe - FE = BI - AI = AB$. Q. E. D.

Corol. 1. Hence $CH = CI$ is a 4th proportional to CA, CF, CD .

Corol. 2. And $fe + FE = 2CH$ or $2CI$; or FE, CH, fe . are in continued arithmetical progression, the common difference being CA the semi-transverse.

Corol. 3. Hence is derived the common method of describing this curve mechanically by points, thus :

In the transverse AB , produced, take the foci F, f , and any point I . Then with the radii AI, BI , and centres F, f , describe arcs intersecting in E , which will be a point in the curve. In like manner, assuming other points I , as many other points will be found in the curve


Then, with a steady hand, the curve line may be drawn through all the points of intersection E .

In the same manner are constructed the other two or conjugate hyperbolas, using the axis ab instead of AB .

THEOREM VI.

If from any Point I in the Axis, a Line IL be drawn touching the Curve in one point L ; and the Ordinate LM be drawn ; and if c be the Centre or the Middle of AB : Then shall CM be to CI as the Square of AM to the Square of AI .

That is,
 $CM : CI :: AM^2 : AI^2$.



For, from the point I draw any line IEH to cut the curve in two points E and H; from which let fall the perps. ED HG; and bisect DG in K.

Then, by theor. 1, $AD \cdot DB : AG \cdot GB :: DE^2 : GH^2$,

and by sim. triangles, $ID^2 : IG^2 :: DE^2 : GH^2$;

theref. by equality, $AD \cdot DB : AG \cdot GB :: ID^2 : IG^2$.

But $DB = CB + CD = CB + CD = CG + CD - AG = 2CK - AG$,

and $GB = ~~BC~~ + CG = CA + CG = CG + CD - AD = 2CK - AD$;

theref. $AD \cdot 2CK - AD \cdot AG : AG \cdot 2CK - AD \cdot AG :: ID^2 : IG^2$,

and, by div. $DG \cdot 2CK : IG^2 - ID^2$ or $DG \cdot 2IK :: AD \cdot 2CK - AD \cdot AG : ID^2$,

or $2CK : 2IK :: AD \cdot 2CK - AD \cdot AG : ID^2$,

or $AD \cdot 2CK : AD \cdot 2IK :: AD \cdot 2CK - AD \cdot AG : ID^2$;

theref. by div. $CK : IK :: AD \cdot AG : AD \cdot 2IK - ID^2$,

and, by div. $CK : CI :: AD \cdot AG : ID^2 - AD \cdot ID + IA$,

or $CK : CI :: AD \cdot AG : AI^2$.


But, when the line IH, by revolving about the point I, comes into the position of the tangent IL, then the points E and H meet in the point L, and the points D, K, G. coincide with the point M; and then the last proportion becomes $OM : CI :: AM^2 : AI^2$. Q. E. D.

THEOREM VII.

If a Tangent and Ordinate be drawn from any Point in the Curve, meeting the Transverse Axis; the Semi-transverse will be a Mean Proportional between the Distances of the said Two Intersections from the centre.

That is,

CA is a mean proportional between CD and CT; or CD, CA, CT, are continued proportionals.



For, by th. 6, $CD : CT :: AD^2 : AT^2$.

that is, $CD : CT :: (CD - CA)^2 : (CA - CT)^2$,

or $CD : CT :: CD^2 + CA^2 : CA^2 + CT^2$,

and $CD : DT :: CD^2 + CA^2 : CD^2 - CT^2$,

or $CD : DT :: CD^2 + CA^2 : (CD + CT) \cdot DT$,

or $CD^2 : CD \cdot DT :: CD^2 + CA^2 : CD \cdot DT + CT \cdot TD$;

hence $CD^2 : CA^2 :: CD \cdot DT : CT \cdot TD$,

and $CD^2 : CA^2 :: CD : CT$.

therefore (th. 78, Geom.) $CD^2 : CA^2 :: CA : CT$.


Q. E. D.
Corol.

Corol. Since ct is always a third proportional to cd, ca ; if the points d, a , remain constant, then will the point t be constant also; and therefore all the tangents will meet in this point t , which are drawn from the point e . of every hyperbola described on the same axis ab , where they are cut by the common ordinate dfe drawn from the point d .

THEOREM VIII.

If there be any Tangent meeting Four Perpendiculars to the Axis drawn from these four Points, namely, the Centre, the two Extremities of the Axis, and the Point of Contact; those Four Perpendiculars will be Proportionals.

That is,
 $AG : DE :: CH : BI.$



For, by theor. 7, $TC : AC :: AC : DC$,
 theref. by div. $TA : AD :: TC : AC$ or CB ,
 and by comp. $TA : TD :: TC : TB$,
 and by sim. tri. $AG : DE :: CH : BI.$ Q. E. D.


Corol. Hence TA, TD, TC, TB } are also proportionals.
 and TG, TE, TH, TI }

For these are as AG, DE, CH, BI , by similar triangles.

THEOREM IX.

If there be any Tangent, and two Lines drawn from the Foci to the Point of Contact; these two Lines will make equal Angles with the Tangent.

That is,
 the $\angle FET = \angle fEe.$



For, draw the ordinate DE , and fe parallel to FE .
 By cor. 1, theor. 5, $CA : CD :: CF : CA + FE$,
 and by th. 7, $CA : CD : CT : CA$;

therefore


therefore - $CT : CF :: CA : CA + FE$;
 and by add. and sub. $TF : Tf :: FE : 2CA + FE$ or fe by th. 5.
 But by sim. tri. $TF : Tf :: FE : fe$;
 therefore - $fe = fe$, and conseq. $\angle e = \angle fee$.
 But, because FE is parallel to fe , the $\angle e = \angle FET$;
 therefore the $\angle FET = \angle fee$. Q. E. D.

Corol. As opticians find that the angle of incidence is equal to the angle of reflection, it appears, from this proposition, that rays of light issuing from the one focus, and meeting the curve in every point, will be reflected into lines drawn from the other focus. So the ray FE is reflected into FE . And this is the reason why the points F, f , are called *foci*, or burning points.

THEOREM X.

All the Parallelograms inscribed between the four Conjugate Hyperbolas are equal to one another, and each equal to the Rectangle of the two Axes.

That is,
 the parallelogram $PQRS =$
 the rectangle $AB \cdot ab$.



Let EG, eg , be two conjugate diameters parallel to the sides of the parallelogram, and dividing it into four less and equal parallelograms. Also, draw the ordinates DE, de , and CK perpendicular to rq ; and let the axis produced meet the sides of the parallelograms, produced, if necessary, in t and t .

Then, by theor. 7, $CT : CA :: CA : CD$,
 and $ct : CA :: CA : cd$;
 theref. by equality, $CT : ct :: cd : CD$;
 but, by sim. triangles, $CT : ct :: TD : cd$,
 theref. by equality, $TD : cd :: cd : CD$,
 and the rectangle, $TD : DC$ is $=$ the square cd^2 .
 Again, by theor. 7, $CD : CA :: CA : CT$,
 or, by division, $CD : CA :: DA : AT$,
 and, by composition, $CD : DB :: DA : DT$;
 conseq. the rectangle $CD \cdot DT = cd^2 = AD \cdot DE^*$.

* *Corol.* Because $cd^2 = AD \cdot DB = CD^2 - CA^2$,

therefore $CA^2 = CD^2 - cd^2$,


In like manner, $ca^2 = de^2 - DE^2$.

But, by theor. 2,	$CA^2 : ca^2 :: (AD \cdot DB \text{ OR}) cd^2 : DE^2,$
therefore	$CA : ca :: cd : DE ;$
In like manner,	$CA : ca :: CD : de,$
or	$ca : de :: CA : CD.$
But, by theor. 7,	$CT : CA :: CA : CD ;$
theref. by equality,	$CT : CA :: ca : de.$
But, by sim. tri.	$CT : CK :: ce : de ;$
theref. by equality,	$CK : CA :: ca : ce,$
and the rectangle	$CK \cdot ce = CA \cdot ca.$
But the rect.	$CK \cdot ce = \text{the parallelogram CEPE},$
theref. the rect.	$CA \cdot ca = \text{the parallelogram CEPE},$
conseq. the rect.	$AB \cdot ab = \text{the paral. PQRS.} \quad \text{Q. E. D.}$

THEOREM XI.

The Difference of the Squares of every Pair of Conjugate Diameters, is equal to the same constant Quantity, namely, the Difference of the Squares of the two Axes.

That is,
 $AB^2 - ab^2 = EG^2 - eg^2 ;$
 where EG, eg, are any conjugate diameters.




For, draw the ordinates ED, ed.

Then, by cor. to theor. 10, $ea^2 = CD^2 - cd^2,$
 and $ca^2 = de^2 - DE^2 ;$
 theref. the difference $CA^2 - ca^2 = CD^2 + DE^2 - cd^2 - de^2.$
 But, by right-angled Δ s, $CE^2 = CD^2 + DE^2,$
 and $ce^2 = cd^2 + de^2 ;$
 theref. the difference $CE^2 - ce^2 = CD^2 + DE^2 - cd^2 - de^2.$
 consequently $CA^2 - ca^2 = CE^2 - ce^2 ;$
 or, by doubling, $AB^2 - ab^2 = EG^2 - eg^2. \quad \text{Q. E. D.}$

THEOREM XII.

All the Parallelograms are equal which are formed between the Asymptotes and Curve, by Lines drawn Parallel to the Asymptotes.

That is, the lines GE, EK, AP, AQ, being parallel to the asymptotes CH, cl ; then the paral. CGEK = paral. CPAQ.



For,

For, let *A* be the vertex of the curve, or extremity of the semi-transverse axis *AC*, perp. to which draw *AL* or *Al*, which will be equal to the semi-conjugate, by definition 19. Also, draw *HEDEh* parallel to *LI*.

Then, by theor. 2, $CA^2 : AL^2 :: CD^2 - CA^2 : DE^2$,
 and, by parallels, $CA^2 : AL^2 :: CD^2 : DH^2$;
 theref. by subtract. $CA^2 : AL^2 :: CA^2 : DH^2 - DE^2$ or
 rect. *HE . Eh* ;
 conseq. the square $AL^2 =$ the rect. *HE . Eh*.

But, by sim. tri. $PA : AL :: GE : EH$,
 and, by the same, $QA : Al :: EK : Eh$;
 theref. by comp. $PA . AQ : AL^2 :: GE . EK : HE . Eh$;
 and, because $AL^2 = HE . Eh$, theref. $PA . AQ = GE . EK$.


But the parallelograms *CGEK*, *CPAQ*, being equiangular, are as the rectangles *GE . EK* and *PA . AQ*.

Therefore the parallelogram *GK* = the paral. *PQ*.

That is, all the inscribed parallelograms are equal to one another. Q. E. D.

Corol. 1. Because the rectangle *GEK* or *CGE* is constant, therefore *GE* is reciprocally as *CG*, or $CG : CP :: PA : GE$. And hence the asymptote continually approaches towards the curve, but never meets it: for *GE* decreases continually as *CG* increases: and it is always of *some* magnitude, except when *CG* is supposed to be infinitely great, for then *GE* is infinitely small, or nothing. So that the asymptote *CG* may be considered as a tangent to the curve at a point infinitely distant from *c*.

Corol. 2. If the abscisses *CD*, *CE*, *CG*, &c. taken on the one asymptote, be in geometrical progression increasing; then shall the ordinates *DH*, *DI*, *GK*, &c. parallel to the other asymptote, be a decreasing geometrical progression, having the same




ratio. For, all the rectangles *CPH*, *CEI*, *CGK*, &c. being equal, the ordinates *DH*, *EI*, *GK*, &c. are reciprocally as the abscisses, *CD*, *CE*, *CG*, &c. which are geometricals. And the reciprocals of geometricals are also geometricals, and in the same ratio, but decreasing, or in converse order.

THEOREM XIII.

The three following Spaces, between the Asymptotes and the Curve, are equal; namely, the Sector or Trilinear Space contained by an Arc of the Curve and two Radii, or Lines drawn from its Extremities to the Centre; and each of the two Quadrilaterals, contained by the said Arc, and two Lines drawn from its Extremities parallel to one Asymptote, and the intercepted Part of the other Asymptote.

That is,

The sector $CAE = PAEG = QAEK$,
all standing on the same arc AE .



For, by theor. 12. $CPAQ = CGEK$;
subtract the common space $CGIQ$,
there remains the paral. $PI =$ the par. IK ;
to each add the trilinear IAE , then
the sum is the quadr. $PAEG = QAEK$.


Again, from the quadrilateral $CAEK$
take the equal triangles CAQ, CEK ,
and there remains the sector $CAE = QAEK$.
Therefore $CAE = QAEK = PAEG$. Q. E. D.

THEOREM XIV.

The Sum or Difference of the Semi-transverse and a Line drawn from the Focus to any point in the Curve, is equal to a Fourth Proportional to the Semi-transverse, the distance from the Centre to the Focus, and the Distance from the Centre to the Ordinate belonging to that Point of the Curve.

That is,

$FE + AC = CI$, or $FE = AI$;
and $FE - AC = CI$, or $FE = BI$.
Where $CA : CF :: CD : CI$ the 4th
propor. to CA, CF, CD .



For,

For, draw AG parallel and equal to ca the semi-conjugate ; and join CG meeting the ordinate DE produced in H .

Then, by theor. 2 $CA^2 : AG^2 :: CD^2 - CA^2 : DE^2$;
and, by sim. Δ s, $CA^2 : AG^2 :: CD^2 - CA^2 : DH^2 - AG^2$;
consequently $DE^2 = DH^2 - AG^2 = DH^2 - ca^2$.

Also $FD = CF \cup CD$, and $FD^2 = CF^2 - 2CF \cdot CD + CD^2$;
but, by right-angled triangles, $FD^2 + DE^2 = FE^2$;
therefore $FE^2 = CF^2 - ca^2 - 2CF \cdot CD + CD^2 + DH^2$.

But by theor. 4, $CF^2 - ca^2 = CA^2$;
and, by supposition, $2CF \cdot CD = 2CA \cdot CI$;
theref. $FE^2 = CA^2 - 2CA \cdot CI + CD^2 + DH^2$;

But, by supposition, $CA^2 : CD^2 :: CF^2$ or $CA^2 + AG^2 : CI^2$;
and, by sim. Δ s, $CA^2 : CD^2 :: CA^2 + AG^2 : CD^2 + DH^2$;
therefore $CI^2 = CD^2 + DH^2 = CH^2$;
consequently $FE^2 = CA^2 - 2CA \cdot CI + CI^2$.


And the root or side of this square is $FE = CI - CA = AI$.
In the same manner is found $fE = CI + CA = BI$. Q. E. D.

Corol. From the demonstration it appears, that $DE^2 = DH^2 - AG^2 = DH^2 - ca^2$. Consequently DH is every where greater than DE ; and so the asymptote CGH never meets the curve, though they be ever so far produced ; but DH and DE approach nearer and nearer to a ratio of equality as they recede farther from the vertex, till at an infinite distance they become equal, and the asymptote is a tangent to the curve at an infinite distance from the vertex.

THEOREM XV.

If a Line be drawn from either Focus, Perpendicular to a Tangent to any Point of the curve ; the Distance of their Intersection from the Centre will be equal to the Semi-transverse Axis.

That is, if FP, fp be perpendicular to the tangent TPP , then shall CP and cp be each equal to to CA or CB .



For,

For, through the point of contact E draw FE , and fe meeting FP produced in G . Then, the $\angle GEP = \angle FEP$, being each equal to the $\angle fep$, and the angles at P being right, and the side PE being common, the two triangles GEP , FEP are equal in all respects, and so $GE = FE$, and $GP = FP$. Therefore, since $FP = \frac{1}{2}FG$, and $FC = \frac{1}{2}fg$, and the angle at F common, the side CP will be $= \frac{1}{2}fg$ or $\frac{1}{2}AB$, that is $CP = CA$ or CB .

And in the same manner $CP = CA$ or CB . Q. E. D.

Corol. 1. A circle described on the transverse axis, as a diameter, will pass through the points P, p ; because all the lines CA, CP, CB , being equal, will be radii of the circle.


Corol. 2. CP is parallel to fe , and CP parallel to FE .

Corol. 3. If at the intersections of any tangent, with the circumscribed circle, perpendiculars to the tangent be drawn, they will meet the transverse axis in the two foci. That is, the perpendiculars PF, pf give the foci F, f .

THEOREM XVI.

The equal Ordinates, or the Ordinates at equal Distances from the Centre, on the opposite Sides and Ends of an Hyperbola, have their Extremities connected by one Right Line passing through the Centre, and that Line is bisected by the Centre.

That is, if $CD = CG$, or the ordinate $DE = GH$; then shall $CE = CH$, and ECH will be a right line.



For, when $CD = CG$, then also is $DE = GH$ by cor. 2, th. 1. But the $\angle D = \angle G$, being both right angles; therefore the third side $CE = CH$, and the $\angle DCE = \angle GCH$, and consequently ECH is a right line.

Corol. 1. And, conversely if ECH be a right line passing through the centre; then shall it be bisected by the centre, or have $CE = CH$; also DE will be $= GH$, and $CD = CG$.

Corol. 2. Hence also, if two tangents be drawn to the two ends E, H of any diameter EH ; they will be parallel to each other,


other, and will cut the axis at equal angles, and at equal distances from the centre. For, the two cd , ca being equal to the two cg , cb , the third proportionals ct , cs will be equal also; then the two sides ce , ct being equal to the two ch , cs , and the included angle ect equal to the included angle hcs , all the other corresponding parts are equal: and so the $\angle t = \angle s$, and te parallel to hs .

Corol. 3. And hence the four tangents, at the four extremities of any two conjugate diameters, form a parallelogram inscribed between the hyperbolas, and the pairs of opposite sides are each equal to the corresponding parallel conjugate diameters.—For, if the diameter eh be drawn parallel to the tangent te or hs , it will be the conjugate to eh by the definition; and the tangents to eh will be parallel to each other, and to the diameter eh for the same reason.

THEOREM XVII.

If two Ordinates ED , ed be drawn from the Extremities E , e , of two Conjugate Diameters, and Tangents be drawn to the same Extremities, and meeting the Axis produced in r and t ;

Then shall cd be a mean proportional between cd , dr , and cd a mean proportional between cd , dt .



For, by theor. 7, $CD : CA :: CA : CT$,
 and by the same, $cd : CA :: CA : CR$;
 theref. by equality, $CD : cd :: CR : CT$,
 But by sim. tri. $DT : cd :: CT : CR$;
 theref. by equality, $CD : cd :: cd : DT$.
 In like manner, $cd : CD :: CD : dr$.

Q. E. D.

Corol. 1. Hence $cd : cd :: cr : ct$.

Corol. 2. Hence also $cd : cd :: de : DE$.

And the rect. $cd \cdot DE = cd \cdot de$, or $\triangle CDE = \triangle cde$.

Corol. 3. Also $cd^2 = CD \cdot DT$, and $cd^2 = cd \cdot dr$.

Or cd a mean proportional between CD , DT ;
 and cd a mean proportional between cd , dr .

THEOREM

THEOREM XVIII.

The same Figure being constructed as in the last Proposition, each Ordinate will divide the Axis, and the semi-axis added to the external Part, in the same Ratio.

[See the last fig.]

That is, $DA : DT :: DC : DE$,
and $da : dr :: dc : db$.

For, by theor. 7, $CD : CA :: CA : CT$,
and by div. $CD : CA :: AD : AT$,
and by comp. $CD : DB :: AD : DT$,
or, - - - - - $DA : DT :: DC : DB$.
In like manner, $da : dr :: dc : db$.

Q. E. D.

Corol. 1. Hence, and from cor. 3 to the last prop. it is,
 $cd^2 = CD \cdot DT = AD \cdot DB = CD^2 - CA^2$,
and $cd \cdot dr = Ad \cdot db = CA^2 - cd^2$.

Corol. 2. Hence also, $CA^2 = CD^2 - cd^2$, and $ca^2 = de^2 - DE^2$.

Corol. 3. Farther, because $CA^2 : ca^2 :: AD \cdot DB : CD^2 : DE^2$,
therefore $CA : ca :: cd : DE$.
likewise $CA : ca :: ed : de$.

THEOREM XIX.


If from any Point in the Curve there be drawn an Ordinate, and a Perpendicular to the Curve, or to the Tangent at that point: Then, the

Dist. on the Trans. between the Centre and Ordinate, CD :

Will be to the Dist. PA ::

As Square of Trans. Axis :

To Square of the Conjugate.



That is,

$$CA^2 : ca^2 :: DC : DP.$$

For, by theor. 2, $CA^2 : ca^2 :: AD \cdot DB : DE^2$,

But, by rt. angled Δ s, the rect. $TD \cdot DP = DE^2$;

and, by cor. 1, theor. 18. $CD \cdot DT = AD \cdot DB$;

therefore - - $CA^2 : ca^2 :: TD \cdot DC : TD \cdot DP$,


or - - - - $CA^2 : ca^2 :: DC : DP$. Q. E. D.

THEOREM

THEOREM XX.

If there be Two Tangents drawn, the One to the Extremity of the Transverse, and the other to the Extremity of any other Diameter, each meeting the other's diameter produced; the two Tangential Triangles so formed, will be equal.

That is,
the triangle $CET =$
the triangle CAN .



For, draw the ordinate DE . Then
By sim. triangles, $CD : CA :: CE : CN$;
but, by theor. 7, $CD : CA :: CA : CT$;
theref. by equal. $CA : CT :: CE : CN$.

The two triangles CET, CAN have then the angle c common, and the sides about that angle reciprocally proportional; those triangles are therefore equal, viz. the $\triangle CET = \triangle CAN$. Q. E. D.


Corol. 1. Take each of the equal tri. CET, CAN , from the common space $CAPE$, and there remains the external $\triangle PAT = \triangle PNE$.

Corol. 2. Also take the equal triangles CET, CAN , from the common triangle CED , and there remains the $\triangle TED =$ trapez. $ANED$.

THEOREM XXI.

The same being supposed as in the last Proposition; then any Lines kq, gq , drawn parallel to the two Tangents, shall also cut off equal Spaces.

That is,
the $\triangle KQG =$ trapez. $ANHG$,
and $\triangle kqg =$ trapez. $anhg$.



For, draw the ordinate DE . Then
The three sim. triangles CAN, CDE, CGH ,

are to each other as CA^2, CD^2, CG^2 ;
 th. by div. the trap. $ANED$: trap. $ANHG$:: $CD^2 - CA^2$: $CG^2 - CA^2$.
 But, by theor. 1, DE^2 : GQ^2 :: $CD^2 - CA^2$: $CG^2 - CA^2$,
 theref. by equ. trap. $ANED$: trap. $ANHG$:: DE^2 : GQ^2 .
 but. by sim. Δ s, tri. TED : tri. KQG :: DE^2 : GQ^2 ;
 theref. by equal. $ANED$: TED :: $ANHG$: KQG .
 But, by cor. 2, theor. 20. the trap. $ANED = \Delta TED$;
 and therefore the trap. $ANHG = \Delta KQG$.

In like manner the trap. $ANhg = \Delta Kqg$. Q. E. D.


Corol. 1. The three spaces $ANHG, TEHG, KQG$ are all equal.

Corol. 2. From the equals $ANHG, KQG$,
 take the equals $ANhg, Kqg$.
 and there remains $ghHG = gqQG$.

Corol. 3. And from the equals $ghHG, gqQG$,
 take the common space $gqLHG$,
 and there remains the $\Delta LQH = \Delta Lqh$.

Corol. 4. Again, from the equals $KQG, TEHG$,
 take the common space $KLHG$,
 and there remains $TELK = \Delta LQH$.

Corol. 5. And when, by
 the lines kq, gh , moving
 with a parallel motion, kq
 comes into the position IR ,
 where eR is the conjugate to
 σA ; then



the triangle kqG becomes the triangle IRC ,
 and the space $ANHG$ becomes the triangle ANC ;
 and therefore the $\Delta IRC = \Delta ANC = \Delta TEC$.


Corol. 6. Also when the lines kq and hq , by moving with a
 parallel motion, come into the position ce, me ,
 the triangle LQH becomes the triangle cem ,
 and the space $TELK$ becomes the triangle TEC ;
 and theref. the $\Delta cem = \Delta TEC = \Delta ANC = \Delta IRC$.

THEOREM XXII.

Any Diameter bisects all its Double Ordinates, or the Lines
 drawn Parallel to the Tangent at its Vertex, or to its Con-
 jugate Diameter.

That

That is, if qg be parallel to the tangent TE , or to ce , then shall $LQ = Lq$.



For, draw QH, qh perpendicular to the transverse. Then by cor. 3, theor. 21, the $\triangle LQH = \triangle Lqh$; but these triangles are also equiangular; consequently their like sides are equal, or $LQ = Lq$.

Corol. 1. Any diameter divides the hyperbola into two equal parts.

For, the ordinates on each side being equal to each other, and equal in number; all the ordinates, or the area, on one side of the diameter, is equal to all the ordinates, or the area, on the other side of it.


Corol. 2. In like manner, if the ordinate be produced to the conjugate hyperbolas at q', q' , it may be proved that $LQ' = Lq'$. Or if the tangent TE be produced, then $EV = EW$. Also the diameter $GCEH$ bisects all lines drawn parallel to TE or qg , and limited either by one hyperbola, or by its two conjugate hyperbolas.

THEOREM XXIII.

As the Square of any Diameter :
Is to the Square of its Conjugate : :
So is the Rectangle of any two Abscisses :
To the Square of their Ordinate.

That is, $CE^2 : ce^2 :: EL \cdot LG \text{ or } CL^2 - CE^2 : LQ^2$.

For, draw the tangent TE , and produce the ordinate QL to the transverse at K . Also draw QH, EM perpendicular to the transverse, and meeting EG in H and M . Then, similar triangles being as the squares of their like sides, it is,



by

by sim. triangles, $\triangle CET : \triangle CLK :: CE^2 : CL^2$;
 or, by division, $\triangle CET : \text{trap. TELK} :: ce^2 : CL^2 - CE^2$.
 Again, by sim. tri. $\triangle cem : \triangle LQH :: ce^2 : LQ^2$.
 But, by cor. 5, theor. 21, the $\triangle cem = \triangle CET$,
 and, by cor. 4, theor. 21, the $\triangle LQH = \text{trap. TELK}$;
 theref. by equality, $CE^2 : ce^2 :: CL^2 - CE^2 : LQ^2$,
 or $CE^2 : ce^2 :: EL \cdot LG : LQ^2$. Q. E. D.

Corol. 1. The squares of the ordinates to any diameter, are to one another as the rectangles of their respective abscisses, or as the difference of the squares of the semi-diameter and of the distance between the ordinate and centre. For they are all in the same ratio of CE^2 to ce^2 .

Corol. 2. The above being the same property as that belonging to the two axes, all the other properties before laid down, for the axes, may be understood of any two conjugate diameters whatever, using only the oblique ordinates of these diameters instead of the perpendicular ordinates of the axes ; namely, all the properties in theorems 6, 7, 8, 16, 17, 20, 21,

Corol. 3. Likewise, when the ordinates are continued to the conjugate hyperbolas at q' , q' , the same properties still obtain, substituting only the sum for the difference of the squares of CE and CL ,

That is, $CE^2 : ce^2 :: CL^2 + CE^2 : LQ'^2$.

And so $LQ^2 : LQ'^2 :: CL^2 - CE^2 : CL^2 + CE^2$.

Corol. 4. When by the motion of Lq' parallel to itself, that line coincides with EV , the last corollary becomes

$$CE^2 : ce^2 :: 2CE^2 : EV^2,$$

$$\text{or } ce^2 : EV^2 :: 1 : 2,$$

$$\text{or } ce : EV :: 1 : \sqrt{2},$$

or as the side of a square to its diagonal.

That is, in all conjugate hyperbolas, and all their diameters, any diameter is to its parallel tangent, in the constant ratio of the side of a square to its diagonal.

THEOREM XXIV.

If any Two Lines, that any where intersect each other, meet the Curve each in Two Points ; then

The Rectangle of the Segments of the one :


Is to the Rectangle of the Segments of the other : :

As the Square of the Diam. Parallel to the former :

To the Square of the Diam. Parallel to the latter.

That

That is, if CR and cr be Parallel to any two Lines PHQ , pHq ; then shall $CR^2 : cr^2 :: PH \cdot HQ : pH \cdot Hq$.



For, draw the diameter $CH E$, and the tangent TE , and its parallels PK , RI , MH , meeting the conjugate of the diameter CR in the points T , K , I , M . Then, because similar triangles are as the squares of their like sides, it is,

by sim. triangles, $CR^2 : GP^2 :: \triangle CRI : \triangle GPK$,
 and - - - - $CR^2 : GH^2 :: \triangle CRI : \triangle GHM$;
 theref. by division, $CR^2 : GP^2 - GH^2 :: CRI : KPHM$.
 Again, by sim. tri. $CE^2 : CH^2 :: \triangle CTE : \triangle CMH$;
 and by division, $CE^2 : CH^2 - CE^2 :: \triangle CTE : TEHM$.

But, by cor. 5, theor. 21, the $\triangle CTE = \triangle CIR$,
 and by cor. 1, theor. 21, $TEHG = KPHG$, or $TEHM = KPHM$;
 theref. by equ. $CE^2 : CH^2 - CE^2 :: CR^2 : GP^2 - GH^2$ or $PH \cdot HQ$.
 In like manner $CE^2 : CH^2 - CE^2 :: cr^2 : pH \cdot Hq$.
 Theref. by equ. $CR^2 : cr^2 :: PH \cdot HQ : pH \cdot Hq$. Q. E. D.


Corol. 1. In like manner, if any other lines $p'H'q'$, parallel to cr or to pq , meet PHQ ; since the rectangles $PH'Q$, $p'H'q'$ are also in the same ratio of CR^2 to cr^2 ; therefore the rect. $PHQ : pHq :: PH'Q : p'H'q'$.

Also, if another line $p'hq'$ be drawn parallel to pq or CR ; because the rectangles $p'hq'$, $p'hq'$ are still in the same ratio, therefore, in general, the rect. $PHQ : pHq :: p'hq' : p'hq'$. That is, the rectangles of the parts of two parallel lines, are to one another, as the rectangles of the parts of two other parallel lines, any where intersecting the former.

Corol. 2. And when any of the lines only touch the curve, instead of cutting it, the rectangles of such become squares, and the general property still attends them.

That

That is,
 $CR^2 : cr^2 :: TE^2 : te^2,$
 or $CR : cr :: TE : te.$
 and $CR : cr :: tE : te.$




Corol. 3. And hence $TE : te :: tE : te.$

THEOREM XXV.

If a Line be drawn through any Point of the Curves, Parallel to either of the Axes, and terminated at the Asymptotes; the Rectangle of its Segments, measured from that Point, will be equal to the Square of the Semi-axis to which it is parallel.

That is,
 the rect. HEK or HEK = CA²,
 and rect. hek or hek = CA².



For, draw AL parallel to ca, and aL to CA. Then by the parallels, CA² : ca² or AL² :: CD² : DH²; and by theor. 2, CA² : CA² :: CD² - CA² : DE²; theref. by subtr. CA² : ca² :: CA² : DH² - DE² or HEK. But the antecedents CA², CA² are equal, theref. the consequents ca², HEK must also be equal.

In like manner it is again, by the parallels, CA² : ca² or AL² :: CD² : DH²; and by theor. 3, CA² : ca² :: CD² + CA² : DE²; theref. by subtr. CA² : ca² :: CA² : DE² - DH² or HEK.

But the antecedents CA², CA² are the same, theref. the conseq. ca², HEK must be equal. In like manner, by changing the axes, is hek or hek = CA².

Corol. 1. Because the rect. HEK = the rect. hek, therefore EH : eH :: eK : EK.

And consequently HE : is always greater than he.

Corol. 2. The rectangle hek = the rect. HEK.

For, by sim. tri. eh : EH :: ek : EK.

SCHOLIUM.


SCHOLIUM.

It is evident that this proposition is general for any line oblique to the axis also, namely, that the rectangle of the segments of any line, cut by the curve, and terminated by the asymptotes, is equal to the square of the semi diameter to which the line is parallel. Since the demonstration is drawn from properties that are common to all diameters.

THEOREM XXVI.

All the rectangles are equal which are made of the Segments of any Parallel Lines cut by the Curve, and limited by the Asymptotes.

That is,
 the rect. $HEK = hEk$.
 and rect. $hek = hek$.



For, each of the rectangles HEK or hEk is equal to the square of the parallel semi-diameter cs ; and each of the rectangles hek or hek is equal to the square of the parallel semi-diameter ci . And therefore the rectangles of the segments of all parallel lines are equal to one another. Q. E. D.

Corol. 1. The rectangle HEK being constantly the same, whether the point E is taken on the one side or the other of the point of contact I of the tangent parallel to HK , it follows that the parts HE, KE , of any line HK , are equal.

And because the rectangle hek is constant, whether the point e is taken in the one or the other of the opposite hyperbolas, it follows, that the parts he, ke , are also equal.

Corol. 2. And when HK comes into the position of the tangent DIL , the last corollary becomes $IL = ID$, and $IM = IN$, and $LM = DN$.

Hence also the diameter CIR bisects all the parallels to DI which are terminated by the asymptote, namely $RH = RK$.

Corol.

Corol. 3. From the proposition, and the last corollary, it follows that the constant rectangle HEK or EHE is $=IL^2$. And the equal constant rect. HEK or EHE $=MIN$ or $IM^2 - IL^2$.

Corol. 4. And hence $IL =$ the parallel semi-diameter cs .


For, the rect. $EHE = IL^2$,
and the equal rect. $eHe = IM^2 - IL^2$,
theref. $IL^2 = IM^2 - IL^2$, or $IM^2 = 2IL^2$;
but, by cor. 4, theor. 23, $IM^2 = 2cs^2$;
and therefore - - - $IL = cs$.

And so the asymptotes pass through the opposite angles of all the inscribed parallelograms.

THEOREM XXVII.

The rectangle of any two Lines drawn from any Point in the Curve, Parallel to two given Lines, and limited by the Asymptotes, is a Constant Quantity.

That is, if AP, EG, DI be parallels,
as also AQ, EK, DM parallels,
then shall the rect. PAQ = rect. GEK = rect. IDM.



For, produce KE, MD to the other asymptote at H, L.

Then, by the parallels, $HE : GE :: LD : ID$;

but - - - - $EK : EK :: DM : DM$;

theref the rectangle $HEK : GEK :: LDM : IDM$.

But, by the last theor. the rect. $HEK = LDM$;

and therefore the rect. $GEK = IDM = PAQ$.


THEOREM XXVIII.

Every Inscribed Triangle, formed by any Tangent and the two Intercepted Parts of the Asymptotes, is equal to a Constant Quantity ; namely Double the Inscribed Parallelogram.

That

That is, the triangle $cts = 2$ paral. gk .

For, since the tangent ts is bisected by the point of contact e , and ek is parallel to tc , and ge to ck ; therefore ck, ks, ge are all equal, as are also cg, gt, ke . Consequently the triangle $gte =$ the triangle kes , and each equal to half the constant inscribed parallelogram ck . And therefore the whole triangle cts , which is composed of the two smaller triangles and the parallelogram, is equal to double the constant inscribed parallelogram gk .




Q. E. D.

THEOREM XXIX.

If from the Point of Contact of any Tangent, and the two Intersections of the Curve with a Line parallel to the Tangent, three parallel Lines be drawn in any Direction, and terminated by either Asymptote; those three Lines shall be in continued Proportion.

That is, if hkm and the tangent il be parallel, then are the parallels dh, ei, gk in continued proportion.




For, by the parallels, $ei : il :: dh : hm$;
and, by the same $ei : il :: gk : km$;
theref. by compos. $ei^2 : il^2 :: dh \cdot gk : hmk$;

but, by theor. 26, the rect. $hmk = il^2$;
and theref. the rect. $dh \cdot gk = ei^2$,

or - - - $dh : ei :: ei : gk$. Q. E. D.

THEOREM XXX.

Draw the semi-diameters ch, cin, ck ;
Then shall the sector $chi =$ the sector ck ,



For, because HK and all its parallels are bisected by CIN ,
 therefore the triangle $CNH = \text{tri. CNK}$,
 and the segment $INH = \text{seg. INK}$;
 consequently the sector $CIH = \text{sec. CIK}$.

Corol. If the geometricals DH, EI, GK be parallel to the other asymptote, the spaces $DHIE, EIKG$ will be equal; for they are equal to the equal sectors CHI, CIK .

So that by taking any geometricals $CD, CE, CG, \&c.$ and drawing $DH, EI, GK, \&c.$ parallel to the other asymptote, as also the radii CH, CI, CK ;

then the sectors $CHI, CIK, \&c.$
 or the spaces $DHIE, EIKG, \&c.$
 will be all equal among themselves.
 Or the sectors $CHI, CHK, \&c.$
 or the spaces $DHIE, DHKG, \&c.$
 will be in arithmetical progression.

And therefore these sectors, or spaces, will be analogous to the logarithms of the lines or bases $CD, CE, CG, \&c.$; namely CHI or $DHIE$ the log. of the ratio of CD to CE , or of CE to $CG, \&c.$; or of EI to DH , or of GK to $EI, \&c.$; and CHK or $DHKG$ the log. of the ratio of CD to $CG, \&c.$ or of GK to $DH, \&c.$




OF THE PARABOLA.

THEOREM I.

The Abscisses are Proportional to the Squares of their Ordinates.

Let AVM be a section through the axis of the cone, and $AGIH$ a parabolic section by a plane perpendicular to the former, and parallel to the side VM of the cone; also let AFH be the common intersection of the two planes, or the axis of the parabola, and FG, HI ordinates perpendicular to it.



Then it will be, as $AF : AH :: FG^2 : HI^2$.

For, through the ordinates FG, HI draw the circular sections, KGL, MIN , parallel to the base of the cone, having KL, MN

MN for their diameters, to which FG, HI are ordinates, as well as to the axis of the parabola.

Then, by similar triangles, $AF : AH :: FL : HN$;
 but, because of the parallels, $KF = MH$;
 therefore - - - $AF : AH :: KF . FL : MH . HN$.
 But, by the circle, $KF . FL = FG^2$, and $MH . HN = HI^2$;
 Therefore - - - $AF : AH :: FG^2 : HI^2$. Q. E. D.

Corol. Hence the third proportional $\frac{FG^2}{AF}$ or $\frac{HI^2}{AH}$ is a constant quantity, and is equal to the parameter of the axis by defin. 16.


Or $AF : FG :: FG : P$ the parameter.
 Or the rectangle $P . AF = FG^2$.

THEOREM II.

As the Parameter of the Axis :
 Is to the Sum of any Two Ordinates : :
 So is the Difference of those Ordinates :
 To the Difference of their Abscisses ;

That is,

$P : GH + DE :: GH - DE : DG$
 Or, $P : KI :: IH : IE$.



For, by cor. theor. 1, $P . AG = GH^2$,
 and - - - $P . AD = DE^2$;
 theref. by subtraction, $P . DG = GH^2 - DE^2$
 Or, - - - $P . DG = KI . IH$,
 therefore - - - $P : KI :: IH : DG$ or EI . Q. E. D.

Corol. Hence because $P . EI = KI . IH$,
 and, by cor. theor. 1, $P . AG = GH^2$,
 therefore - - - $AG : EI :: GH^2 : KI . IH$.


So that any diameter EI is as the rectangle of the segments KI, IH of the double ordinate KH.

THEOREM III.

The Distance from the Vertex to the Focus is equal to $\frac{1}{4}$ of the Parameter, or to Half the Ordinate at the Focus.

That

That is,
 $AF = \frac{1}{2}FE = \frac{1}{4}P$,
 where F is the focus.




For, the general property is $AF : FE :: FE : P$.
 But by definition 17 - $FE = \frac{1}{2}P$;
 therefore also - $AF = \frac{1}{2}FE = \frac{1}{4}P$. Q. E. D.

THEOREM IV.

A Line drawn from the Focus to any Point in the Curve, is equal to the Sum of the Focal Distance and the Absciss of the Ordinate to that Point.


That is,
 $FE = FA + AD = GD$,
 taking $AG = AF$.



For, since $FD = AD \cup AF$,
 theref. by squaring, $FD^2 = AF^2 - 2AF \cdot AD + AD^2$,
 But, by cor. theor. 1, $DE^2 = P \cdot AD = 4AF \cdot AD$;
 theref. by addition, $FD^2 + DE^2 = AF^2 + 2AF \cdot AD + AD^2$,
 But, by right-ang. tri. $FD^2 + DE^2 = FE^2$;
 therefore - $FE^2 = AF^2 + 2AF \cdot AD + AD^2$,
 and the root or side is $FE = AF + AD$,
 or - - - $FE = GD$, by taking $AG = AF$.

Q. E. D.

Corol. 1. If, through the point G, the line GH be drawn perpendicular to the axis, it is called the directrix of the parabola. The property of which, from this theorem, it appears, is this: That drawing any line HE parallel to the axis, HE is always equal to FE the distance of the focus from the point E.




Corol. 2. Hence also the curve is easily described by points, Namely, in the axis produced take $AG = AF$ the focal distance, and draw a number of lines EE perpendicular to the axis AD; then with the distances GD, GD, GD, &c. as radii and the centre F, draw arcs crossing the parallel ordinates in E, E, E, &c. Then draw the curve through all the points, E, E, E.

THEOREM

THEOREM V.

If a Tangent be drawn to any Point of the Parabola, meeting the Axis produced ; and if an Ordinate to the Axis be drawn from the Point of Contact ; then the Absciss of that Ordinate will be equal to the External Part of the Axis.

That is,
if TC touch the curve
at the point c ;
then is $AT = AM$.



For, from the point T , draw any line cutting the curve in the two points E, H : to which draw the ordinates DE, GH ; also draw the ordinate MC to the point of contact c .

Then, by th. 1, $AD : AG :: DE^2 : GH^2$;
and by sim. tri. $TD^2 : TG^2 :: DE^2 : GH^2$;
theref. by equality, $AD : AG :: TD^2 : TG^2$;
and, by division, $AD : DG :: TD^2 : TG^2 - TD^2$ or $DG \cdot (TD + TG)$;
or - - - $AD : TD :: TD : TD + TG$;
and, by division, $AD : AT :: TD : TG$,
and again by div. $AD : AT :: AT : AG$;
or - - - AT is a mean propor. between AD, AG .

Now if the line TH be supposed to revolve about the point T ; then, as it recedes farther from the axis, the points E and H approach towards each other, the point E descending and the point H ascending, till at last they meet in the point c , when the line becomes a tangent to the curve at c . And then the points D and G meet in the point M , and the ordinates DE, GH in the ordinates CM . Consequently AD, AG , becoming each equal to AM , their mean proportional AT will be equal to the absciss AM . That is, the external part of the axis, cut off by a tangent, is equal to the absciss of the ordinate to the point of contact.


Q. E. D.

THEOREM VI.

If a tangent to the Curve meet the Axis produced ; then the Line drawn from the Focus to the Point of Contact, will be equal to the Distance of the Focus from the Intersection of the Tangent and Axis.

That

That is,
 $FC = FT.$



For, draw the ordinate DC to the point of contact c .
 Then, by theor. 5, $AT = AD$;
 therefore $FT = AF + AD$.
 But, by theor. 4, $FC = AF + AD$;
 theref. by equality, $FC = FT$.

Corol. 1. If CG be drawn perpendicular to the curve, or to the tangent, at c ; then shall $FG = FC = FT$.

For, draw FH perpendicular to TC , which will also bisect TC , because $FT = FC$; and therefore, by the nature of the parallels, FH also bisects TG in F . And consequently $FG = FT = FC$.

So that F is the centre of a circle passing through t, c, g .

Corol. 2. The tangent at the vertex AH is a mean proportional between AF and AD .

For, because FHT is a right angle,
 therefore AH is a mean between AF, AT ,
 or between AF, AD , because $AD = AT$.
 Likewise, FH is a mean between FA, FT ,
 or between FA, FC .

Corol. 3. The tangent TC makes equal angles with FC and the axis FT .

For, because $FT = FC$,
 therefore the $\angle FCT = \angle FTC$.
 Also, the angle $GCF =$ the angle GCK ,
 drawing ICK parallel to the axis AG .


Corol. 4. And because the angle of incidence GCK is = the angle of reflection GCF ; therefore a ray of light falling on the curve in the direction KC , will be reflected to the focus F . That is, all rays parallel to the axis, are reflected to the focus, or burning point.

THEOREM

THEOREM VII.

If there be any Tangent, and a Double Ordinate drawn from the Point of Contact, and also any Line parallel to the Axis, limited by the Tangent and Double Ordinate : then shall the Curve divide that Line in the same Ratio, as the Line divides the double Ordinate.

That is,
 $IE : EK :: CK : KL.$




FOR, by sim. triangles, $CK : KI :: CD : DT$ or $2DA$;
 but by the def. the param. $P : CL :: CD : 2DA$;
 therefore, by equality, $P : CK :: CL : KI.$
 But, by theor. 2, - - - $P : CK :: KL : KE$;
 therefore, by equality $CL : KL :: KI : KE$;
 and by division, - - - $CK : KL :: IE : EK.$ Q. E. D.

THEOREM VIII.

The same being supposed as in theor. 7 ; then shall the External Part of the Line between the Curve and Tangent be proportional to the Square of the intercepted Part of the Tangent, or to the Square of the intercepted Part of the Double Ordinate.

That is, IE is as CI^2 or as CK^2
 and $IE, TA, ON, PL, \&c.$
 are as $CI^2, CT^2, CO^2, CP^2, \&c.$
 or as $CK^2, CD^2, CM^2, CL^2, \&c.$



FOR, by theor. 7, $IE : EK :: CK : KL,$
 or, by equality, $IE : EK :: CK^2 : CK . KL.$
 But, by cor. th. 2, EK is as the rect. $CK . KL,$
 therefore - - - IE is as $CK^2,$ or as $CI^2.$ Q. E. D.


Corol. As this property is common to every position of the tangent, if the lines $IE, TA, ON, \&c.$ be appended on the points $I, T, O, \&c.$ and moveable about them, and of such lengths as that their extremities $E, A, N, \&c.$ be in the curve of a parabola
 in

in some one position of the tangent; then making the tangent revolve about the point c , it appears that the extremities E, A, N , &c. will always form the curve of some parabola, in every position of the tangent.

THEOREM IX.

The Abscisses of any Diameter, are as the Squares of their Ordinates.

That is, CQ, CR, CS , &c.
are as QE^2, RA^2, SN^2 , &c.
Or $CQ : CR :: QE^2 : RA^2$,
&c.



For, draw the tangent CT , and the externals EI, AT, NO , &c. parallel to the axis, or to the diameter CS .

Then, because the ordinates QE, RA, SN , &c. are parallel to the tangent CT , by the definition of them, therefore all the figures IQ, TR, OS , &c. are parallelograms, whose opposite sides are equal;

namely, - - - IE, TA, ON , &c.

are equal to - - - CQ, CR, CS , &c.

Therefore, by theor. 8, CQ, CR, CS , &c.

are as - - - CI^2, CT^2, CO^2 , &c.

or as their equals : : QE^2, RA^2, SN^2 , &c.


Q. E. D.

Corol. Here like as in theor. 2, the difference of the abscisses is as the difference of the squares of their ordinates, or as the rectangles under the sum and difference of the ordinates, the rectangle of the sum and difference of the ordinates being equal to the rectangle under the difference of the abscisses and the parameter of that diameter, or a third proportional to any absciss and its ordinate.

THEOREM X.

If a Line be drawn parallel to any Tangent, and cut the Curve in two points; then if two Ordinates be drawn to the Intersections, and a third to the Point of Contact, these three Ordinates will be in Arithmetical Progression, or the Sum of the Extremes will be equal to Double the Mean.

That is,
 $EG + HI = 2CD.$



For, draw EK parallel to the axis, and produce HI to L.

Then, by sim. triangles, $EK : HK :: TD \text{ or } 2AD : CD;$

but, by theor. 2, $EK : HK :: KL : P \text{ the param.}$

theref. by equality, $2AD : KL :: CD : P.$

But, by the defin. $2AD : 2CD :: CD : P;$

theref. the 2d terms are equal, $KL = 2CD,$

that is, $EG + HI = 2CD.$


Q. E. D.

Corol. When the point E is on the other side of AI; then
 $HI - GE = 2CD.$

THEOREM XI.

Any Diameter bisects all its Double Ordinates, or Lines parallel to the tangent at its Vertex.

That is,
 $ME = MH.$



For, to the axis AI draw the ordinates EG, CD, HI, and MN parallel to them, which is equal to CD.

Then, by theor. 10, $2MN$ or $2CD = EG + HI$,
therefore M is the middle of EH .

And, for the same reason, all its parallels are bisected.


Q. E. D.

SCHOL. Hence, as the abscisses of any diameter and their ordinates have the same relations as those of the axis, namely, that the ordinates are bisected by the diameter, and their squares proportional to the abscisses; so all the other properties of the axis and its ordinates and abscisses before demonstrated, will likewise hold good for any diameter and its ordinates and abscisses. And also those of the parameters, understanding the parameter of any diameter, as a third proportional to any absciss and its ordinate. Some of the most material of which are demonstrated in the following theorems.

THOREM XII.

The Parameter of any Diameter is equal to four Times the Line drawn from the Focus to the Vertex of that Diameter.

That is, $4FC = p$,
the param. of the diam. CM .



FOR, draw the ordinate MA parallel to the tangent CT ; also CD, MN perpendicular to the axis AN , and FH perpendicular to the tangent CT .

Then the abscisses AD, CM or AT , being equal, by theor. 5, the parameters will be as the squares of the ordinates GD, MA or CT , by the definition;

that is,	-	-	$P : p :: CD^2 : CT^2.$
But, by sim. tri.	-	-	$FH : FT :: CD : CT;$
therefore	-	-	$P : p :: FH^2 : FT^2.$
But, by cor. 2, th. 6,	-	-	$FH^2 = FA \cdot FT;$
therefore	-	-	$P : p :: FA \cdot FT : FT^2.$
or, by equality	-	-	$P : p :: FA : FT$ or $FC.$

But,


But, by theor. 3, $p = 4FA$,
 and therefore - $p = 4FT$ or $4FC$. Q. E. D.

Corol. Hence the parameter p of the diameter CM is equal to $4FA + 4AD$, or to $p + 4AD$, that is, the parameter of the axis added to $4AD$.

THEOREM XIII.

If an Ordinate to any Diameter, pass through the Focus, it will be equal to Half its Parameter ; and its Absciss equal to One Fourth of the same Parameter.

That is, $CM = \frac{1}{4}p$,
 and $ME = \frac{1}{2}p$.




FOR, join FC , and draw the tangent CT .

By the parallels, $CM = FT$;
 and, by theor. 6, $FC = FT$;
 also, by theor. 12, $FC = \frac{1}{4}p$;
 therefore - - $CM = \frac{1}{4}p$.

Again, by the defin. CM or $\frac{1}{4}p : ME :: ME : p$,
 and consequently $ME = \frac{1}{2}p = 2CM$. Q. E. D.

Corol. 1. Hence, of any diameter, the double ordinate which passes through the focus, is equal to the parameter, or to quadruple its absciss.

Corol. 2. Hence, and from cor. 1 to theor. 4, and theor. 6 and 12, it appears, that if the directrix GH be drawn, and any lines HE , HE , parallel to the axis ; then every parallel HE will be equal to EF , or $\frac{1}{4}$ of the parameter of the diameter to the point E .




THEOREM

THEOREM XIV.

If there be a Tangent, and any line drawn from the point of Contact and meeting the Curve in some other Point, as also another Line parallel to the Axis, and limited by the First line and the tangent : then shall the Curve divide this Second Line in the same Ratio, as the Second Line divides the first Line.

That is,
 $IE : EK :: CK : KL.$



FOR, draw LP parallel to IK, or to the axis.

Then by theor. 8, $IE : PL :: CI^2 : CP^2,$

or, by sim. tri. - $IE : PL :: CK^2 : CL^2.$

Also, by sim. tri. $IK : PL :: CK : CL,$

or - - - $IK : PL :: CK^2 : CK \cdot CL ;$

therefore by equality $IE : IK :: CK : CL \cdot CL^2 ;$

or, - - - $IE : IK :: CK : CL ;$


and, by division, $IE : EK :: CK : KL ;$

Q. E. D.

Corol. When $CK = KL,$ then $IE = EK = \frac{1}{2}IK.$

THEOREM XV.

If from any point of the curve there be drawn a Tangent, and also Two Right Lines to cut the Curve ; and Diameters be drawn through the points of Intersection E and L, meeting those Two Right Lines in two other Points G, and K : Then will the Line KG joining these last Two Points be parallel to the Tangent.




FOR,

For, by theor. 14, $CK : KL :: EI : EK$;
 and by composition, $CK : CL :: EI : KI$;
 and by the parallels $CK : CL :: GH : LH$;
 But, by sim. tri. - $CK : CL :: KI : LH$;
 theref. by equal. - $KI : LH :: GH : LH$;
 consequently - $KI = GH$,
 and therefore - KG is parallel and equal to IH . Q. E. D.

THEOREM XVI.

If an Ordinate be drawn to the Point of Contact of any Tangent, and another Ordinate produced to cut the Tangent ;
 It will be, as the Difference of the Ordinates :
 Is to the Difference added to the external Part : :
 So is Double the first Ordinate :
 To the Sum of the Ordinates.

That is, $KH : KI :: KL : KG$.



For, by cor. 1 theor. 1, $P : DC :: DC : DA$,
 and - - - - $P : 2DC :: DC : DT$ or $2DA$.
 But, by sim. triangles, $KI : KC :: DC : DT$;
 therefore by equality, $P : 2DC :: KI : KC$,
 or, - - - - $P : KI :: KL : KC$.
 Again, by theor. 2, $P : KH :: KG : KC$;
 therefore by equality, $KH : KI :: KL : KG$. Q. E. D.

Corol. 1. Hence, by composition and division,
 it is, $KH : KI :: GK : GI$,
 and $HI : HK :: HK : KL$,
 also $IH : IK :: IK : IG$;
 that is, IK is a mean proportional between IG and IH .

Corol. 2. And from this last property a tangent can easily be drawn to the curve from any given point i . Namely, draw iHG perpendicular to the axis, and take ik a mean proportional between iH , iG ; then draw kc parallel to the axis, and c will be the point of contact, through which and the given point i the tangent ic is to be drawn.

THEOREM


THEOREM XVII.

If a Tangent cut any Diameter produced, and if an Ordinate to that Diameter be drawn from the Point of Contact; then the Distance in the Diameter produced, between the Vertex and the Intersection of the Tangent, will be equal to the Absciss of that Ordinate.

That is, $IE = EK$.

For, by the last th. $IE : EK :: CK : KL$.

But, by theor. 11, $CK = KL$,
and therefore $IE = EK$.




Corol. 1. The two tangents CI, LI , at the extremities of any double ordinate CL , meet in the same point of the diameter of that double ordinate produced. And the diameter drawn through the intersection of two tangents, bisects the line connecting the points of contact.

Corol. 2. Hence we have another method of drawing a tangent from any given point I without the curve, Namely, from I draw the diameter IK , in which take $EK = EI$, and through K draw CL parallel to the tangent at K ; then C and L are the points to which the tangents must be drawn from I .

THEOREM XVIII.

If a Line be drawn from the Vertex of any Diameter, to cut the Curve in some other Point, and an Ordinate of that Diameter be drawn to that Point, as also another Ordinate any where cutting the Line, both produced if necessary: The Three will be continual Proportionals, namely, the two Ordinates and the Part of the Latter limited by the said Line drawn from the vertex.

That is, DE, GH, GI are continual proportionals, or $DE : GH :: GH : GI$.



For, by theor. 9 - - - $DE^2 : GH^2 :: AD : AG$;

and, by sim. tri. - - - $DE : GI :: AD : AG$;

theref. by equality, - - - $DE : GI :: DE^2 : GH^2$;

that is, of the three DE, GH, GI , 1st : 3d :: 1st² : 2d²;

therefore - - - - - 1st : 2d :: 2d : 3d,

that is, - - - - - $DE : GH :: GH : GI$. Q. E. D.

Corol.

Corol. 1. Or their equals, CK, CH, GI , are proportionals; where EK is parallel to the diameter AD .

Corol. 2. Hence it is $DE : AG :: p : GI$, where p is the parameter, or $AG : GI :: DE : p$.

For, by the defin. $AG : CH :: CH : p$.

Corol. 3. Hence also the three MN, MI, MO , are proportionals, where MO is parallel to the diameter, and AM parallel to the ordinates.

For, by theor. 9, MN, MI, MO ,

or their equals AP, AG, AD ,

are as the squares of PK, CH, DE ,

or of their equals GI, CH, CK ,

which are proportionals by cor. 1

THEOREM XIX.


If a Diameter cut any Parallel Lines terminated by the Curve, the Segments of the Diameter will be as the rectangles of the Segments of those Lines.

That is, $EK : EM :: CK . KL : EM . MO$.

Or, EK is as the rectangle $CK . KL$.

For, draw the diameter AD to which the parallels CL, NO are ordinates and the ordinate EQ parallel to them.

Then CK is the difference, and KL the sum of the ordinates EQ, CR ; also EM the difference, and MO the sum of the ordinates EQ, CR . And the differences of the abscissas, are QR, QS, OR, EK, EM .



Then by cor. theor. 9, $QR : QS :: CK . KL : EM . MO$, that is $EK : EM :: CK . KL : EM . MO$.

Corol. 1. The rect. $CK . KL = \text{rect. } EK$ and the param. of AD . For the rect. $CK . KL = \text{rect. } QR$ and the param. of AD .

Corol. 2. If any line CL be cut by two diameters, KE, CH the rectangles of the parts of the line, are as the segments of the diameters.

For EK is as the rectangle $CK . KL$.

and CH is as the rectangle $CH . HL$;

therefore $EK : CH :: CK . KL : CH . HL$.

Corol.

Corol. 3. If two parallels, cl , no , be cut by two diameters, em , gi ; the rectangles of the parts of the parallels, will be as the segments of the respective diameters.

For - - - $EK : EM :: CK . KL : NM . MO$,
 and - - - $EK : GH :: CK . KL : CH . HL$,
 theref. by equal. $EM : GH :: NM . MO : CH . HL$,

Corol. 4. When the parallels come into the position of the tangent at p their two extremities, or points in the curve unite in the point of contact p ; and the rectangle of the parts becomes the square of the tangent, and the same properties still follow them.

So that, $EV : PV :: PV : p$ the param.

$GW : PW :: PW : p$,


$EV : GW :: PV^2 : PW^2$,

$EV : GH :: PV^2 : CH . HL$.

THEOREM XX.

If two Parallels intersect any other two Parallels; the Rectangles of the Segments will be respectively Proportional.

That is, $CK . KL : DK . KE :: GI . IH : NI . IO$.



For, by cor. 3, theor. 19, $PK : QI :: CK . KL : GI . IH$;

and by the same, $PK : QI :: DK : KE . NI . IO$;

theref. by equal. $CK . KL : DK . KE :: GI . IH : NI . IO$;

Corol. When one of the pairs of intersecting lines comes into the position of their parallel tangents, meeting and limiting each other, the rectangles of their segments become the squares of their respective tangents. So that the constant ratio of the rectangles, is that of the square of their parallel tangents, namely,

$CK . KL : DK . KE :: \text{tang}^2 . \text{parallel to } CL : \text{tang}^2 . \text{parallel to } DE$.


THEOREM XXI.

If there be Three Tangents intersecting each other; their Segments will be in the same Proportion.

That

That is, $GI : IH :: CG : CD :: DH : HE$.

For, through the points G, I, D, H , draw the diameters GK, IL, DM, HN ; as also the lines CI, EI , which are double ordinates to the diameters GK, HN , by cor. 1, theor. 16; therefore the diameters GK, DM, HN , bisect the lines CL, CE, LE ;



hence $KM = CM - CK = \frac{1}{2}CE - \frac{1}{2}CL = \frac{1}{2}LE = LN$ of NE ,

and $MN = ME - NE = \frac{1}{2}CE - \frac{1}{2}LE = \frac{1}{2}CL = CK$ of KL .

But, by parallels, $GI : IH :: KL : LN$,

and $CG : GD :: CK : KM$,

also $DH : HE :: MN : NE$.

But the 3d terms KL, CK, MN are all equal;

as also the 4th terms LN, KM, NE .

Therefore the first and second terms, in all the lines, are proportional, namely $GI : IH :: CG : GD :: DH : HE$. Q. E. D.


THEOREM XXII.

If a Rectangle be described about a Parabola, having the same Base and Altitude; and a diagonal Line be drawn from the Vertex to the Extremity of the Base of the Parabola, forming a right-angled Triangle, of the same Base and Altitude also; then any Line or Ordinate drawn across the three Figures, perpendicular to the Axis, will be cut in Continual Proportion by the Sides of those Figures.

That is,

$$EF : EG :: EG : EH,$$

Or, EF, EG, EH , are in continued proportion.



For, by theor 1, $AB : AE :: BC^2 : EG^2$,

and, by sim. tri. $AB : AE :: BC : EF$,

theref. of equality, $EF : BC :: EG^2 : BC^2$

that is $EF : EH :: EG^2 : EH^2$,

theref. by Geom. th. 78, EF, FG, EH are proportionals,

or $EF : EG :: EG : EH$. Q. E. D.

THEOREM XXIII.

The Area or Space of a Parabola, is equal to Two-Thirds of its Circumscribing Parallelogram.

That is, the space $ABCGA = \frac{2}{3} ABCD$;
or the space $ADCGA = \frac{1}{3} ABCD$.

FOR, conceive the space $ADCGA$ to be composed of, or divided into, indefinitely small parts, by lines parallel to DC or AB , such as IG , which divide AD into like small and equal parts, the number or sum of which is expressed by the line AD . Then,

by the parabola, $BC^2 : EG^2 :: AB : AE$,
that is, - - - $AD^2 : AI^2 :: DC : IG$.

Hence it follows, that any one of these narrow parts, as IG , is $= \frac{DC}{AD^2} \times AI^2$; hence, AD and DC being given or constant quantities, it appears that the said parts IG , &c. are proportional to AI^2 , &c. or proportional to a series of square numbers, whose roots are in arithmetical progression, and the area $ADCGA$ equal to $\frac{DC}{AD^2}$ drawn into the sum of such a series of arithmetics, the number of which is expressed by AD .

Now, by the remark at pag. 227 this vol. the sum of the squares of such a series of arithmetics, is expressed by $\frac{1}{6}n \cdot n+1 \cdot 2n+1$, where n denotes the number of them. In the present case, n represents an infinite number, and then the two factors $n+1$, $2n+1$, become only n and $2n$, omitting the 1 as inconsiderable in respect of the infinite number n : hence the expression above becomes barely $\frac{1}{6}n \cdot n \cdot 2n = \frac{1}{3}n^3$.

To apply this to the case above : n will denote AD or EC ; and the sum of all the AI^2 's becomes $\frac{1}{3} AD^3$ or $\frac{1}{3} BC^3$; consequently the sum of all the $\frac{DC}{AD^2} \times AI^2$'s, is $\frac{DC}{AD^2} \times \frac{1}{3} AD^3 = \frac{1}{3} AD \cdot DC = \frac{1}{3} BD$, which is the area of the exterior part $ADCGA$. That is, the said exterior part $ADCGA$, is $\frac{1}{3}$ of the parallelogram $ABCD$; and consequently the interior part $ABCGA$ is $\frac{2}{3}$ of the same parallelogram.

Q. E. D.

Corol.


Corol. The part $AFCGA$, inclosed between the curve and the right line AFC , is $\frac{1}{6}$ of the same parallelogram, being the difference between $ABCGA$ and the triangle $ABCFA$, that is between $\frac{2}{3}$ and $\frac{1}{2}$ of the parallelogram.

THEOREM XXIV.

The Solid Content of a Paraboloid (or Solid generated by the Rotation of a Parabola about its Axis), is equal to Half its Circumscribing Cylinder.

LET ABC be a paraboloid, generated by the rotation of the parabola AC about its axis AD . Suppose the axis AD be divided into an infinite number of equal parts, through which let circular planes pass, as EFG , all those circles making up the whole solid paraboloid.

Now if $c =$ the number 3.1416 , then $2c \times FG$ is the circumference of the circle EFG whose radius is FG ; therefore $c \times FG^2$ is the area of that circle.



But, by *cor. theor. 1, Parabola*, $p \times AF = FG^2$, where p denotes the parameter of the parabola; consequently $pc \times AF$ will also express the same circular section EG , and therefore $pc \times$ the sum of all the AF 's will be the sum of all those circular sections, or the whole content of the solid paraboloid.

But all the AF 's form an arithmetical progression, beginning at 0 or nothing, and having the greatest term and the sum of all the terms each expressed by the whole axis AD . And since the sum of all the terms of such a progression, is equal to $\frac{1}{2} AD \times AD$ or $\frac{1}{2} AD^2$, half the product of the greatest term and the number of terms; therefore $\frac{1}{2} AD^2$ is equal to the sum of all the AF 's, and consequently $pc \times \frac{1}{2} AD^2$, or $\frac{1}{2} c \times p \times AD^2$, is the sum of all the circular sections, or the content of the paraboloid.

But, by the parabola, $p : DC :: DC : AD$, or $p = \frac{DC^2}{AD}$; consequently $\frac{1}{2} c \times p \times AD^2$ becomes $\frac{1}{2} c \times AD \times DC^2$ for the solid content of the paraboloid. But $c \times AD \times DC^2$ is equal to the cylinder $BCIH$; consequently the paraboloid is the half of its circumscribing cylinder.

Q. E. D.


THEOREM XXV.

The Solidity of the Frustum BEGC of the Paraboloid, is equal to a Cylinder whose height is DF, and its Base Half the Sum of the two Circular Bases EG, BC.

For, by the last theor. $\frac{1}{2}pc \times AD^2 =$ the solid ABC,
 and, by the same, $\frac{1}{2}pc \times AF^2 =$ the solid AEG,
 theref. the diff. $\frac{1}{2}pc \times (AD^2 - AF^2) =$ the frust. BEGC,
 But $AD^2 - AF^2 = DF \times (AD + AF)$,
 theref. $\frac{1}{2}pc \times DF \times (AD + AF) =$ the frust. BEGC.
 But, by the parab. $p \times AD = DC$. and $p \times AF = FG^2$;
 theref. $\frac{1}{2}c \times DF \times (DC + FG^2) =$ the frust. BEGC.

Q. E. D.

ON THE CONIC SECTIONS AS EXPRESSED BY ALGEBRAIC EQUATIONS, CALLED THE EQUATIONS OF THE CURVE.



1. For the Ellipse.

Let d denote AB, the transverse or any diameter ;
 $c = HI$ its conjugate ;
 $x = AK$, any absciss, from the extremity of the diam.
 $y = DK$ the correspondent ordinate.

Then, theor. 2, $AB^2 : HI^2 :: AK \cdot KB : DK^2$,
 that is, $d^2 : c^2 :: x(d-x) : y^2$, hence $d^2 y^2 = c^2 (dx - x^2)$,
 or $dy = c \sqrt{(dx - x^2)}$, the equation of the curve.

And from these equations, any one of the four letters or quantities, d, c, x, y , may easily be found, by the reduction of equations, when the other three are given.

Or, if p denote the parameter, $= c^2 \div d$ by its definition ;
 then, by cor. th. 2, $d : p :: x(d-x) : y^2$, or $dy^2 = p(dx - x^2)$,
 which is another form of the equation of the curve.

Otherwise.

Otherwise.

Or, if $d = AC$ the semi-axis ; $c = CH$ the semi-conjugate ; $p = c^2 \div d$ the semi-parameter ; $x = CK$ the absciss counted from the centre ; and $y = DK$ the ordinate as before. Then is $AK = d - x$, and $KB = d + x$, and $AK \cdot KB = (d - x) \times (d + x) = d^2 - x^2$.

Then, by th. 2, $d^2 : c^2 :: d^2 - x^2 : y^2$, and $d^2 y^2 = c^2 (d^2 - x^2)$, or $dy = c \sqrt{(d^2 - x^2)}$, the equation of the curve.

Or, $d : p :: d^2 - x^2 : y^2$, and $dy^2 = p (d^2 - x^2)$, another form of the equation to the curve : from which any one of the quantities may be found, when the rest are given.

2. For the Hyperbola.

Because the general property of the opposite hyperbolas, with respect to their abscisses and ordinates, is the same as that of the ellipse, therefore the process here is the very same as in the former case for the ellipse ; and the equation to the curve must come out the same also, with sometimes only the change of the sign of a letter or term from $+$ to $-$, or from $-$ to $+$, because here the abscisses lie beyond or without the transverse diameter, whereas they lie between or upon them in the ellipse. Thus, making the same notation for the whole diameter, conjugate, absciss. and ordinate, as at first in the ellipse ; then, the one absciss AK being x , the other BK will be $d + x$, which in the ellipse was $d - x$; so the sign of x must be changed in the general property and equation, by which it becomes $d^2 : c^2 :: x (d + x) : y^2$; hence $d^2 y^2 = c^2 (dx + x^2)$ and $dy = c \sqrt{(dx + x^2)}$, the equation of the curve.

Or, using p the parameter as before, it is, $d : p :: x (d + x) : y^2$, or $dy^2 = p (dx + x^2)$, another form of the equation to the curve.

Otherwise, by using the same letters d , c , p , for the halves of the diameters and parameter, and x for the absciss CK counted from the centre ; then is $AK = x - d$, and $BK = x + d$ and the property $d^2 : c^2 :: (x - d) \times (x + d) : y^2$, gives $d^2 y^2 = c^2 (x^2 - d^2)$, or $dy = c \sqrt{(x^2 - d^2)}$, where the signs of d^2 and x^2 are changed from what they were in the ellipse.

Or again, using the semi-parameter, $d : p :: x^2 - d^2 : y^2$, and $dy^2 = p (x^2 - d^2)$ the equation of the curve.

But for the conjugate hyperbola, as in the figure to theorem 3, the signs of both x^2 and d^2 will be positive ; for the property in that theorem being $CA^2 : ca^2 :: CD^2 + CA^2 : de^2$,


it

it is $d^2 : c^2 :: x^2 + d^2 : y^2 = De^2$, or $d^2 y^2 = c^2 (x^2 + d^2)$, and $dy = c \sqrt{x^2 + d^2}$, the equation to the conjugate hyperbola.

Or, as $d : p :: x^2 + d^2 : y^2$, and $dy^2 = p (x^2 + d^2)$, also the equation to the same curve.

On the Equation to the Hyperbola between the Asymptotes.

Let CE and CB be the two asymptotes to the hyperbola dFD , its vertex being F , and EF, bd, AF, BD ordinates parallel to the asymptotes. Put AF or $EF = a$, $CB = x$, and $BD = y$. Then, by theor. 28, $AF \cdot EF = CB \cdot BD$, or $a^2 = xy$, the equation to the hyperbola, when the abscisses and ordinates are taken parallel to the asymptotes.



3. For the Parabola.

If x denote any absciss beginning at the vertex, and y its ordinate, also p the parameter. Then, by cor. theorem 1, $AK : KD :: KD : p$, or $x : y :: y : p$; hence $px = y^2$ is the equation to the parabola.

4. For the Circle.

Because the circle is only a species of the ellipse, in which the two axes are equal to each other; therefore, making the two diameters d and c equal in the foregoing equations to the ellipse, they become $y^2 = dx - x^2$, when the absciss x begins at the vertex of the diameter: and $y^2 = d^2 - x^2$, when the absciss begins at the centre.

Scholium.

In every one of these equations, we perceive that they rise to the 2d or quadratic degree, or to two dimensions; which is also the number of points in which every one of these curves may be cut by a right line. Hence it is also that these four curves are said to be lines of the 2d order. And these four are all the lines that are of that order, every other curve being of some higher, or having some higher equation, or may be cut in more points by a right line.

ELEMENTS OF ISOPERIMETRY.

Def. 1. When a variable quantity has its mutations regulated by a certain law, or confined within certain limits, it is called a *maximum* when it has reached the greatest magnitude it can possibly attain ; and, on the contrary, when it has arrived at the least possible magnitude, it is called a *minimum*.

Def. 2. *Isoperimeters*, or *Isoperimetrical figures*, are those which have equal perimeters.

Def. 3. The *Locus* of any point, or intersection, &c. is the right line or curve in which these are always situated.

The problem in which it is required to find, among figures of the same or of different kinds, those which within equal perimeters, shall comprehend the greatest surfaces, has long engaged the attention of mathematicians. Since the admirable invention of the method of Fluxions, this problem has been elegantly treated by some of the writers on that branch of analysis ; especially by Maclaurin and Simpson. A much more extensive problem was investigated at the time of " the war of problems," between the two brothers John and James Bernoulli : namely, " To find, among all the isoperimetrical curves between given limits, such a curve, that constructing a second curve, the ordinates of which shall be functions of the ordinates or arcs of the former, the area of the second curve shall be a maximum or minimum." While, however, the attention of mathematicians was drawn to the most abstruse inquiries connected with isoperimetry, the *elements* of the subject were lost sight of. Simpson was the first who called them back to this interesting branch of research, by giving in his neat little book of Geometry a chapter on the maxima and minima of geometrical quantities, and some of the simplest problems concerning isoperimeters. The next who treated this subject in an elementary manner was Simon Lhuillier, of Geneva, who in 1782, published his treatise *De Relatione mutua Capacitatis et Terminorum Figurarum*, &c. His principal object in the composition of that work was to supply the deficiency in this respect which he found in most of the elementary Courses, and to determine, with regard to both, the most usual surfaces and solids, those which possessed the minimum of contour with the same capacity ; and, reciprocally, the maximum of capacity with the same boundary. M. Legendre has also considered the same subject in a manner somewhat different from either Simpson or Lhuillier, in his *Elements de Géométrie*. An elegant geometrical tract, on the
same


same subject, was also given, by Dr. Horsley, in the *Philos. Trans.* vol. 75, for 1775; contained also in the *New Abridgment*, vol. 13, page 653. The chief propositions deduced by these four geometers, together with a few additional propositions, are reduced into one system in the following theorems.

SECTION I. SURFACES.

THEOREM I.

Of all Triangles of the same Base, and whose Vertices fall in a right Line given in Position, the one whose Perimeter is a Minimum is that whose sides are equally inclined to that Line.

Let AB be the common base of a series of triangles ABC' , ABC , &c. whose vertices c' , c , fall in the right line LM , given in position, then is the triangle of least perimeter that whose sides AC , BC , are inclined to the line LM in equal angles.



For, let BM be drawn from B , perpendicularly to LM , and produced till $DM = BM$: join AD , and from the point c where AD cuts LM draw BC : also, from any other point c' , assumed in LM , draw $c'A$, $c'B$, $c'D$. Then the triangles DMC , BMC , having the angle $DCM = \text{angle } ACL$ (th. 7 Geom.) $= \angle MCB$ (by hyp.) $\angle DMC = \angle BMC$, and $DM = BM$, and MC common to both, have also $DC = BC$ (th. 1 Geom.)

So also, we have $c'D = c'B$. Hence $AC + CB = AC + CD = AD$, is less than $AC' + c'D$ (theor. 10 Geom.), or than its equal $AC' + c'B$. And consequently, $AB + BC + AC$ is less than $AB + BC' + AC'$.

Q. E. D.

Cor. 1. Of all triangles of the same base and the same altitude, or of all equal triangles of the same base, the isosceles triangle has the smallest perimeter.

For, the locus of the vertices of all triangles of the same altitude will be a right line LM parallel to the base; and when LM in the above figure becomes parallel to AB , since $\angle MCB = \angle ACL$, $\angle MCB = \angle CBA$ (th. 12 Geom.), $\angle ACL = \angle CAB$; it follows that $\angle CAE = \angle CBA$, and consequently $AC = CB$ (th. 4 Geom.).

Cor. 2. Of all triangles of the same surface, that which has the minimum perimeter is equilateral.

For

For the triangle of the smallest perimeter, with the same surface, must be isosceles, whichever of the sides be considered as base: therefore, the triangle of smallest perimeter has each two or each pair of its sides equal, and consequently it is equilateral.

Cor. 3. Of all rectilinear figures, with a given magnitude and a given number of sides, that which has the smallest perimeter is equilateral.

For so long as any two adjacent sides are not equal, we may draw a diagonal to become a base to those two sides, and then draw an isosceles triangle equal to the triangle so cut off, but of less perimeter: whence the corollary is manifest.

Scholium.

To illustrate the second corollary above, the student may proceed thus: assuming an isosceles triangle whose base is *not* equal to either of the two sides, and then, taking for a new base one of those sides of that triangle, he may construct another isosceles triangle equal to it, but a smaller perimeter. Afterwards, if the base and sides of this second isosceles triangle are not respectively equal, he may construct a third isosceles triangle equal to it, but of a still smaller perimeter: and so on, by performing these successive operations, he will find that all the triangles will approach nearer and nearer to an equilateral triangle.


THEOREM II.

Of all Triangles of the Same Base, and of Equal Perimeters, the Isosceles Triangle has the Greatest Surface.

Let ABC , ABD , be two triangles of the same base AB and with equal perimeters, of which the one ABC is isosceles, the other is not: then the triangle ABC has a surface (or an altitude) greater than the surface (or than the altitude) of the triangle ABD .

Draw $c'D$ through D , parallel to AB , to cut CE (drawn perpendicular to AB) in c' : then it is to be demonstrated that CE is greater than $c'E$.

The triangles $AC'B$, ADB , are equal both in base and altitude; but the triangle $AC'B$ is isosceles, while ADB is scalene: therefore the triangle $AC'B$ has a smaller perimeter than the triangle ADB (th. 1 cor. 1), or than ACB (by hyp.) Consequently AC'



$<$ AC ; and in the right-angled triangles AEC', AEC, having AE common, we have $C'E < CE^*$. Q. E. D.

Cor. Of all isoperimetrical figures, of which the number of sides is given, that which is the greatest has all its sides equal. And in particular, of all isoperimetrical triangles, that whose surface is a maximum, is equilateral.

For, so long as any two adjacent sides are not equal, the surface may be augmented without increasing the perimeter.

Remark Nearly as in this theorem may it be proved that, of all triangles of equal heights, and of which the sum of the two sides is equal, that which is isosceles has the greatest base. And, of all triangles standing on the same base and having equal vertical angles, the isosceles one is the greatest.

THEOREM III.


Of all Right Lines that can be drawn through a Given Point between Two Right Lines Given in Position, that which is Bisected by the Given Point forms with the other two Lines the Least Triangle.

Of all right lines GD, AB, GD, that can be drawn through a given point P to cut the right lines CA, CD, given in position, that AB, which is bisected by the given point P, forms with GA, CD, the least triangle, ABC.

For, let EE be drawn through A parallel to CD, meeting DG (produced if necessary) in E : then the triangles PBD, PAE, are manifestly equiangular ; and, since the corresponding sides PB, PA are equal, the triangles are equal also. Hence PBD will be less or greater than PAG, according as CG is greater or less than CA. In the former case, let PACD, which is common, be added to both ; then will BAC be less than DCC (ax. 4 Geom.). In the latter case, if PGCB be added, DCG will be the greater than BAC ; and consequently in this case also BAC is less than DCG.

Q. E. D.

Cor. If PM and PN be drawn parallel to CB and CA respectively, the two triangles PAM, PBN, will be equal, and



* When two mathematical quantities are separated by the character $<$, it denotes that the preceding quantity is *less than* the succeeding one: when, on the contrary, the separating character is $>$, it denotes that the preceding quantity is *greater than* the succeeding one.

these

these two taken together (since $AM = PN = MC$) will be equal to the parallelogram $PMCN$: and consequently the parallelogram $PMCN$ is equal to half ABC , but less than half DGC . From which it follows (consistently with both the algebraical and geometrical solution of prob. 8, Application of Algebra to Geometry), that a parallelogram is always less than half a triangle in which it is inscribed, except when the base of the one is half the base of the other, or the height of the former half the height of the latter; in which case the parallelogram is just half the triangle: this being the maximum parallelogram inscribed in the triangle.

Scholium.

From the preceding corollary it might easily be shown, that the least triangle which can possibly be described about, and the greatest parallelogram which can be inscribed in, any curve concave to its axis, will be when the subtangent is equal to half the base of the triangle, or to the whole base of the parallelogram: and that the two figures will be in the ratio of 2 to 1. But this is foreign to the present enquiry.

THEOREM IV.

Of all Triangles in which two Sides are Given in Magnitude, the Greatest is that in which the two Given Sides are Perpendicular to each other.

For, assuming for base one of the given sides, the surface is proportional to the perpendicular let fall upon that side from the opposite extremity of the other given side: therefore, the surface is the greatest when that perpendicular is the greatest; that is to say, when the other side is not inclined to that perpendicular, but *coincides* with it: hence the surface is a maximum when the two given sides are perpendicular to each other.

Otherwise. Since the surface of a triangle, in which two sides are given, is proportional to the sine of the angle included between those two sides; it follows, that the triangle is the greatest when that sine is the greatest: but the greatest sine is the sine total, or the sine of a quadrant; therefore the two sides given make a quadrantal angle, or are perpendicular to each other. Q. E. D.

THEOREM V.



Of all Rectilinear Figures in which all the Sides except one are known, the Greatest is that which may be Incribed in a Semicircle whose Diameter is that Unknown Side.

For, if you suppose the contrary to be the case. then whenever the figure made with the sides given, and the side unknown is not inscribable in a semicircle of which this latter is the diameter, viz. whenever any one of the angles, formed by lines drawn from the extremities of the unknown side to one of the summits of the figure, is not a right angle; we may make a figure greater than it, in which that angle shall be right, and which shall only differ from it in that respect: therefore, whenever all the angles formed by right lines drawn from the several vertices of the figure to the extremities of the unknown line, are not right angles, or do not fall in the circumference of a semicircle, the figure is not in its maximum state. Q. E. D.

THEOREM VI.

Of all Figures made with sides Given in Number and Magnitude, that which may be Incribed in a Circle is the Greatest.

Let $ABCDEFGG$ be the polygon inscribed, and $abcdefg$ a polygon with equal sides, but not inscribable in a circle; so that $AB=ab$, $BC=bc$, &c.; it is affirmed that the polygon $ABCDEFGG$ is greater than the polygon $abcdefg$.



Draw the diameter EP ; join AP , PB ; upon $ab = AB$ make the triangle abp , equal in all respects to ABP ; and join ep . Then, of the two figures $edcbp$, $pagfe$, one at least is not (by hyp.) inscribable in the semicircle of which ep is the diameter. Consequently, one at least of these two figures is smaller than the corresponding part of the figure $APBCDEFG$ (th. 5). Therefore the figure $APBCDEFG$ is greater than the figure $apbcdefg$: and if from these there be taken away the respective triangles APB , apb , which are equal by construction, there will remain (ax. 5 Geom.) the polygon $ABCDEFGG$ greater than the polygon $abcdefg$. Q. E. D.

THEOREM

THEOREM VII.

The Magnitude of the Greatest Polygon which can be contained under any number of Unequal Sides, does not at all depend on the Order in which those Lines are connected with each other.

For, since the polygon is a maximum under given sides, it is inscribable in a circle (th. 6). And this inscribed polygon is constituted of as many isosceles triangles as it has sides, those sides forming the bases of the respective triangles, the other sides of all the triangles being radii of the circle, and their common summit the centre of the circle. Consequently, the magnitude of the polygon, that is, of the assemblage of these triangles, does not at all depend on their disposition, or arrangement round the common centre. Q. E. D.

THEOREM VIII.

If a Polygon Inscribed in a Circle have all its Sides Equal, all its Angles are likewise Equal, or it is a regular Polygon.

For, if lines be drawn from the several angles of the polygon, to the centre of the circumscribing circle, they will divide the polygon into as many isosceles triangles as it has sides; and each of these isosceles triangles will be equal to either of the others in all respects, and of course they will have the angles at their bases all equal: consequently, the angles of the polygon, which are each made up of two angles at the bases of two contiguous isosceles triangles, will be equal to one another. Q. E. D.

THEOREM IX.

Of all Figures having the Same Number of Sides and Equal Perimeters, the Greatest is Regular.

For, the greatest figure under the given conditions has all sides equal (th. 2 cor.). But since the sum of the sides and the number of them are given, each of them is given: therefore (th. 6), the figure is inscribable in a circle: and consequently (th. 8) all its angles are equal; that is, it is regular.

Q. E. D.

Cor. Hence we see that regular polygons possess the property of a maximum of surface, when compared with any other figures of the same name and with equal perimeters.

THEOREM

THEOREM X.


A Regular Polygon has a Smaller Perimeter than an Irregular one Equal to it in Surface, and having the Same Number of Sides.

This is the converse of the preceding theorem, and may be demonstrated thus: Let R and I be two figures equal in surface and having the same number of sides, of which R is regular, I irregular: let also R' be a regular figure similar to R , and having a perimeter equal to that of I . Then (th. 9) $R' : I$; but $I = R$; therefore $R' > R$. But R' and R are similar; consequently, perimeter of $R' >$ perimeter of R ; while per. $R' =$ per. I (by hyp.). Hence, per. $I >$ per. R . Q. E. D.

THEOREM XI.

The Surfaces of Polygons. Circumscribed about the Same or Equal Circles, are respectively as their Perimeters*.

Let the polygon $ABCD$ be circumscribed about the circle $EFGH$; and let this polygon be divided into triangles, by lines drawn from its several angles to the centre O of the circle. Then, since each of the tangents, AB , BC , &c. is perpendicular to its corresponding radius OE , OF , &c. drawn to the point of contact (th. 46 Geom.); and since the area of a triangle is equal to the rectangle of the perpendicular and half the base (Mens. of Surfaces, pr. 2); it follows, that the area of each of the triangles AEO , BCO , &c. is equal to the rectangle of the radius of the circle and half the corresponding side AE , BC , &c.: and consequently, the area of the polygon $ABCD$, circumscribing the circle, will be equal to the rectangle of the radius of the circle and half the perimeter of the polygon. But, the surface of the circle is equal to the rectangle of the radius and half the circumference (th. 94 Geom.). Therefore, the surface of the circle, is to that of the polygon, as half the cir-



* This theorem, together with the analogous ones respecting bodies circumscribing cylinders and spheres, were given by Emerson in his Geometry, and their use in the theory of Isoperimeters was just suggested; but the full application of them to that theory is due to Simon Lhuillier.

circumference of the former, to half the perimeter of the latter; or, as the circumference of the former, to the perimeter of the latter. Now, let P and P' be any two polygons circumscribing a circle c : then, by the foregoing, we have

$$\text{surf. } c : \text{surf. } P :: \text{circum. } c : \text{perim. } P,$$

$$\text{surf. } c : \text{surf. } P' :: \text{circum. } c : \text{perim. } P'.$$

But, since the antecedents of the ratios in both these proportions, are equal, the consequents are proportional: that is, $\text{surf. } P : \text{surf. } P' :: \text{perim. } P : \text{perim. } P'$. Q. E. D.

Corol. 1. And one of the triangular portions ABO , of a polygon circumscribing a circle, is to the corresponding circular sector, as the side AB of the polygon, to the arc of the circle included between AO and BO .

Cor. 2. Every circular arc is greater than its chord, and less than the sum of the two tangents drawn from its extremities and produced till they meet.

The first part of this corollary is evident, because a right line is the shortest distance between two given points. The second part follows at once from this proposition: for $EA + AH$ being to the arch EIH , as the quadrangle $AECI$ to the circular sector $IIEO$; and the quadrangle being greater than the sector, because it contains it; it follows that $EA + AH$ is greater than the arch EIH *.

Cor. 3. Hence also, any single tangent EA , is greater than its corresponding arc EI .

THEOREM XII.

If a Circle and a Polygon, Circumscribable about another Circle, are Isoperimeters, the Surface of the Circle is a Geometrical Mean Proportional between that Polygon and a Similar Polygon (regular or irregular) Circumscribed about that Circle.

Let c be a circle, P a polygon isoperimetrical to that circle, and circumscribable about some other circle, and P' a polygon similar to P and circumscribable about the circle c : it is affirmed that $P : c :: c : P'$.

* This second corollary is introduced, not because of its immediate connection with the subject under discussion, but because, notwithstanding its simplicity, some authors have employed whole pages in attempting its demonstration, and failed at last.

For, $P : P' :: \text{perim}^2 . P :: \text{perim}^2 . P' :: \text{circum}^2 . C : \text{perim}^2 . P'$
 by th. 89, Geom. and the hypothesis.

But (th. 11) $P' : C :: \text{per} . P' : \text{cir} . C :: \text{per}^2 . P' : \text{per} . P' \times \text{cir} . C$
 Therefore $P : C :: - - - - - \text{cir}^2 . C : \text{per} . P \times \text{cir} . C$
 $:: \text{cir} . C : \text{per} . P' :: C : P' . \quad \text{Q. E. D.}$

THEOREM XIII.

If a Circle and a Polygon, Circumscribable about another Circle, are Equal in Surface, the Perimeter of that Figure is a Geometrical Mean Proportional between the Circumference of the first Circle and the Perimeter of a Similar Polygon Circumscribed about it.

Let $c = P$, and let P' be circumscribed about c and similar to c : then it is affirmed that $\text{cir} . c : \text{per} . P :: \text{per} . P : \text{per} . P'$
 For, $\text{cir} . c : \text{per} . P' :: C : P' :: P : P' :: \text{per}^2 . P : \text{per}^2 . P'$
 Also, $\text{per} . P' : \text{per} . P - - - - - :: \text{per}^2 . P' : \text{per} . P \times \text{per} . P'$
 Therefore, $\text{cir} . c : \text{per} . P - - - - - :: \text{per}^2 . P : \text{per} . P \times \text{per} . P'$
 $:: \text{per} . P : \text{per} . P' . \quad \text{Q. E. D.}$

THEOREM XIV.

The Circle is Greater than any Rectilinear Figure of the Same Perimeter; and it has a Perimeter Smaller than any Rectilinear Figure of the Same Surface.

For, in the proportion, $P : C :: C : P'$, (th. 12), since $c < P'$,
 therefore $P < C$.

And, in the propor. $\text{cir} . c : \text{per} . P :: \text{per} . P : \text{per} . P'$ (th. 13),
 or, $\text{cir} . c : \text{per} . P' :: \text{cir}^2 . C : \text{per}^2 . P$,
 $\text{cir} . c < \text{per} . P'$;

therefore, $\text{cir} . c < \text{per}^2 . P$, or $\text{cir} . c < \text{per} . P . \quad \text{Q. E. D.}$

Cor. 1. It follows at once, from this and the two preceding theorems, that rectilinear figures which are isoperimeters, and each circumscribable about a circle, are respectively in the inverse ratio of the perimeters, or of the surfaces, of figures similar to them, and both circumscribed about one and the same circle. And that the perimeters of equal rectilineal figures, each circumscribable about a circle, are respectively in the subduplicate ratio of the perimeters or of the surfaces, of figures, similar to them, and both circumscribed about one and the same circle.


Cor. 2. Therefore, the comparison of the perimeters of equal regular figures, having different numbers of sides, and that

that of the surfaces of regular isoperimetrical figures, is reduced to the comparison of the perimeters, or of the surfaces of regular figures respectively similar to them, and circumscribable about one and the same circle.

Lemma 1.

If an acute angle of a right-angled triangle be divided into any number of equal parts, the side of the triangle opposite to that acute angle is divided into unequal parts, which are greater as they are more remote from the right angle.

Let the acute angle c , of the right-angled triangle ACF , be divided into equal parts, by the lines CB , CD , CE , drawn from that angle to the opposite side; then shall the parts AB , BD , &c. intercepted by the lines drawn from c ; be successively longer as they are more remote from the right angle A .



For the angles ACD , BCE , &c. being bisected by CB , CD , &c. therefore by theor. 83 Geom. $AC : CD :: AB : BD$, and $BC : CE :: BD : DE$, and $DC : CF :: DE : EF$. And by th. 21 Geom. $CD > CA$, $CE > CB$, $CF > CD$, and so on: whence it follows, that $DB > AB$, $DE > DB$, and so on. Q. E. D.

Cor. Hence it is obvious that, if the part the most remote from the right angle A , be repeated a number of times equal to that in'o which the acute angle is divided, there will result a quantity greater than the side opposite to the divided angle.


THEOREM XV.

If two Regular Figures, Circumscribed about the Same Circle, differ in their Number of Sides by Unity, that which has the Greatest number of Sides shall have the Smallest Perimeter.

Let CA be the radius of a circle, and AB , AD , the half sides of two regular polygons circumscribed about that circle, of which the number of sides differ by unity, being respectively $n + 1$ and n . The angles ACB , ACD ,

therefore are respectively the $\frac{1}{n+1}$ and the $\frac{1}{n}$ th

part of two right angles; consequently these angles are as n and $n + 1$: and hence, the angle may be conceived divided into $n + 1$ equal parts, of which BCD is one.



Consequently, (cor. to the lemma) $(n + 1) BD > AD$. Taking, then, unequal quantities from equal quantities we shall have

$$(n + 1) AD - (n + 1) BD < (n + 1) AD - AD,$$

$$\text{OR } (n + 1) AB < n \cdot AD.$$

That is, the semiperimeter of the polygon whose half side is AB is smaller than the semiperimeter of the polygon whose half side is AD : whence the proposition is manifest.

Cor. Hence, augmenting successively by unity the number of sides, it follows generally, that the perimeters of polygons circumscribed about any proposed circle, become smaller as the number of their sides become greater.

THEOREM XVI.

The Surfaces of Regular Isoperimetrical Figures are Greater as the Number of their Sides is Greater: and the Perimeters of Equal Regular Figures are Smaller as the Number of their Sides is Greater.

For, 1st. Regular isoperimetrical figures are (cor. 1. th. 14) in the inverse ratio of figures similar to them circumscribed about the same circle. And (th. 15) these latter are smaller when the number of sides is greater: therefore, on the contrary, the former become greater as they have more sides.

2dly. The perimeters of equal regular figures are (cor. 1 th. 14) in the subduplicate ratio of the perimeters of similar figures circumscribed about the same circle: and (th. 15) these latter are smaller as they have more sides: therefore the perimeters of the former also are smaller when the number of their sides is greater. Q. E. D.



SECTION II. SOLIDS.

THEOREM XVII.

Of all Prisms of the Same Altitude, whose Base is Given in Magnitude and Species, or Figure, or Shape, the Right Prism has the Smallest Surface.

For, the area of each face of the prism is proportional to its height; therefore the area of each face is the smallest when its height is the smallest, that is to say, when it is equal to the altitude of the prism itself: and in that case the prism is evidently a right prism. Q. E. D.

THEOREM

THEOREM XVIII.

Of all Prisms whose Base is Given in Magnitude and Species, and whose Lateral Surface is the same, the Right Prism has the Greatest Altitude, or the Greatest Capacity.

This is the converse of the preceding theorem, and may readily be proved after the manner of theorem 2.

THEOREM XIX.

Of all Right Prisms of the Same Altitude, whose Bases are Given in Magnitude and of a Given number of Sides, that whose Base is a Regular Figure has the Smallest Surface.

For, the surface of a right prism of given altitude, and base given in magnitude, is evidently proportional to the perimeter of its base. But (th. 10) the base being given in magnitude, and having a given number of sides, its perimeter is smallest when it is regular: whence, the truth of the proposition is manifest.

THEOREM XX.

Of two Right Prisms of the Same Altitude, and with Irregular Bases Equal in Surface, that whose Base has the Greatest Number of sides has the smallest Surface: and, in particular, the Right Cylinder has a Smaller Surface than any Prism of the Same Altitude and the Same Capacity.

The demonstration is analogous to that of the preceding theorem, being at once deducible from theorems 16 and 14.

THEOREM XXI.

Of all Right Prisms whose Altitudes and whose Whole Surfaces are Equal, and whose Bases have a Given Number of Sides, that whose Base is a Regular Figure is the Greatest.

Let P, P' , be two right prisms of the same name, equal in altitude, and equal whole surface, the first of these having a regular, the second an irregular base; then is the base of the prism P , less than the base of the prism P' .

For, let P'' be a prism of equal altitude, and whose base is equal to that of the prism P' and similar to that of the prism P . Then the lateral surface of the prism P'' is smaller than the lateral surface of the prism P' (th. 19): hence, the total surface

face

face of P'' is smaller than the total surface of P' , and therefore (by hyp.) smaller than the whole surface of P . But the prisms P' and P have equal altitudes and similar bases; therefore the dimensions of the base of P'' are smaller than the dimensions of the base of P . Consequently the base of P'' , or that of P' , is less than the base of P ; or the base of P greater than that of P' . Q. E. D.

THEOREM XXII.

Of Two Right Prisms, having Equal Altitudes, Equal Total Surfaces, and Regular Bases, that whose Base has the Greatest number of Sides, has the Greatest Capacity. And, in particular, a right Cylinder is Greater than any Right Prism of Equal Altitude and Equal Total Surface.

The demonstration of this is similar to that of the preceding theorem, and flows, from th. 20.

THEOREM XXIII.

The Greatest Parallelopiped which can be contained under the Three parts of a Given Line, any way taken, will be that constituted of Equal length, breadth, and depth.

For, let AB be the given line, and, if possible, let two parts AE, ED , be unequal. Bisect AD in C , then will the rectangle under $AE (= AC + CE)$ and $ED (= AC - CE)$, be less than AC^2 , or than $AC \cdot CD$, by the square of CE (th. 33 Geom.). Consequently, the solid $AE \cdot ED \cdot DB$, will be less than the solid $AC \cdot CD \cdot DB$; which is repugnant to the hypothesis.




Cor. Hence, of all the rectangular parallelopipeds, having the sum of their three dimensions the same, the cube is the greatest.

THEOREM XXIV.

The Greatest Parallelopiped that can possibly be contained under the Square of one Part of a Given Line, and the other Part, any way taken, will be when the former Part is the Double of the latter.

Let AB be a given line, and $AC = 2CB$, then is $AC^2 \cdot CB$ the greatest possible.



For,

For, let ac' and $c'b$ be any other parts into which the given line ab may be divided; and let ac, ac' , be bisected in d, d' , respectively. Then shall $ac^2 \cdot cb = 4ad \cdot dc \cdot cb$ (COR. to theor. 31 Geom.) $> 4ad' \cdot d'c \cdot cb$, or greater than its equal $c'a^2 \cdot c'b$, by the preceding theorem.

THEOREM XXV.

Of all Right Parallelopipeds Given in Magnitude, that which has the Smallest Surface has all its Faces Squares, or is a Cube. And reciprocally, of all Parallelopipeds of Equal Surface, the Greatest is a Cube.

For, by theorems 19 and 21, the right parallelopiped having the smallest surface with the same capacity, or the greatest capacity with the same surface, has a square for its base. But, any face whatever may be taken for base: therefore, in the parallelopiped whose surface is the smallest with the same capacity, or whose capacity is the greatest with the same surface, any two opposite faces whatever are squares: consequently, this parallelopiped is a cube.

THEOREM XXVI.

The Capacities of Prisms Circumscribing the Same Right Cylinder, are Respectively as their Surfaces, whether Total or Lateral.

For, the capacities are respectively as the bases of the prisms; that is to say (th. 11), as the perimeters of their bases; and these are manifestly as the lateral surfaces: whence the proposition is evident.

Cor. The surface of a right prism circumscribing a cylinder, is to the surface of that cylinder, as the capacity of the former, to the capacity of the latter.

Def. The Archimedean cylinder is that which circumscribes a sphere, or whose altitude is equal to the diameter of its base.

THEOREM XXVII.

The Archimedean Cylinder has a Smaller Surface than any other Right Cylinder of Equal Capacity; and it is Greater than any other Right Cylinder of Equal Surface.

Let c and c' denote two right cylinders, of which the first is Archimedean, the other not: then,

1st,

1st; If . . . $c=c'$, surf. $c <$ surf. c' .

2dly, if surf. $c =$ surf. c' , $c >$ c' .

For having circumscribed about the cylinders, c, c' , the right prisms P, P' , with square bases, the former will be a cube, the second not: and the following series of equal ratios will obtain, viz. $c : P ::$ surf. $c : surf. P ::$ base $c : base P ::$ base $c' : base P' :: c' : P' ::$ surf. $c' : surf. P'$.

Then, 1st: when $c = c'$. Since $c : P :: c' : P'$, it follows that $P = P'$; and therefore (th. 25) surf. $P <$ surf. P' . But, surf. $c : surf. P ::$ surf. $c' : surf. P'$; consequently surf. $c <$ surf. c' . Q. E. 1D.

2dly: when surf. $c =$ surf. c' . Then, since surf. $c : surf. P ::$ surf. $c' : surf. P'$, it follows that surf. $P =$ surf. P' ; and therefore (th. 25) $P >$ P' . But $c : P :: c' : P'$; consequently $c >$ c' . Q. E. 2D.

THEOREM XXVIII.

Of all Right Prisms whose Bases are Circumscribable about Circles, and Given in Species, that whose Altitude is Double the Radius of the Circle Inscribed in the Base, has the Smallest Surface with the Same Capacity, and the Greatest Capacity with the Same Surface.

This may be demonstrated exactly as the preceding theorem, by supposing cylinders inscribed in the prisms.

Scholium.

If the base cannot be circumscribed about a circle, the right prism which has the minimum surface or the maximum capacity, is that whose lateral surface is quadruple of the surface of one end, or that whose lateral surface is two-thirds of the total surface. This is manifestly the case with the Archimedean cylinder; and the extension of the property depends solely on the mutual connexion subsisting between the properties of the cylinder, and those of circumscribing prisms.

THEOREM XXIX.

The Surfaces of Right Cones Circumscribed about a Sphere, are as their solidities.


For, it may be demonstrated, in a manner analogous to the demonstrations of theorems 11 and 26, that these cones are

are equal to right cones whose altitude is equal to the radius of the inscribed sphere, and whose bases are equal to the total surfaces of the cones : therefore the surfaces and solidities are proportional.

THEOREM XXX.

The Surface or the Solidity of a Right Cone Circumscribed about a Sphere, is Directly as the Square of the Cone's Altitude, and Inversely as the Excess of that Altitude over the Diameter of the Sphere.

Let VAT be a right-angled triangle which, by its rotation upon VA as an axis, generates a right cone ; and BDA the semicircle which by a like rotation upon VA forms the inscribed sphere : then, the surface or the solidity of the cone varies as $\frac{VA^2}{VB}$.



For, draw the radius CD to the point of contact of the semicircle and VT . Then, because the triangles VAT , VDC , are similar, it is $AT : VT :: CD : VC$.

And, by compos. $AT : AT + VT : CD : CD + CV = VA$;

Therefore $AT^2 : (AT + VT) AT :: CD : VA$, by multiplying the terms of the first ratio by AT .

But, because VB , VD , VA are continued proportionals, it is $VB : VA :: VD^2 : VA^2 :: CD^2 : AT^2$ by sim. triangles.

But $CD : VA :: AT^2 : (AT + VT) AT$ by the last ; and these mult. give $CD \cdot VB : VA^2 :: CD^2 : (AT + VT) AT$,

$$\text{OR } VB : CD :: VA^2 : (AT + VT) AT = CD \cdot \frac{VA^2}{VB}.$$

But the surface of the cone, which is denoted by $\pi \cdot AT^2 + \pi \cdot AT \cdot VT^*$, is manifestly proportional to the first member of this equation, is also proportional to the second member,


or, since CD is constant, it is proportional to $\frac{AV^2}{BV}$, or to a third proportional to BV and AV . And, since the capacities of these circumscribing cones are as their surfaces (th. 29), the truth of the whole proposition is evident.

Lemma 2.

The difference of two right lines being given, the third proportional to the less and the greater of them is a minimum when the greater of those lines is double the other.

* π being = 3.141593.

Let AV and BV be two right lines, whose difference AB is given, and let AP be a third proportional to BV and AV ; then is AP a minimum when $AV = 2BV$.



For, since $AP : AV :: AV : BV$;

By division $AP : AP - AV :: AV : AV - BV$;

That is, $AP : VP :: AV : AB$.

Hence $VP \cdot AV = AP \cdot AB$.

But $VP \cdot AV$ is either $=$ or $< \frac{1}{4}AP^2$ (cor. to th. 31 Geom. and th. 23 of this chapter.)

Therefore $AP \cdot AB < \frac{1}{4}AP^2$: whence $4AB < AP$, or $AP > 4AB$, Consequently the minimum value of AP is the quadruple of AB ; and in that case $PV = VA = 2AB$. Q. E. D.*

THEOREM XXXI.

Of all Right Cones Circumscribed about the Same Sphere, the Smallest is that whose Altitude is Double the Diameter of the Sphere.

For, by th. 30, the solidity varies as $\frac{VA^2}{VB}$ (see the fig. to that theorem): and, by lemma 2, since $VA - VB$ is given, the third proportional $\frac{VA^2}{VB}$ is a minimum when $VA = 2AB$. Q. E. D.

Cor. 1. Hence, the distance from the centre of the sphere to the vertex of the least circumscribing cone, is triple the radius of the sphere.

Cor. 2. Hence also, the side of such cone is triple the radius of its base.

* Though the evidence of a single demonstration, conducted on sound mathematical principles, is really irresistible, and therefore needs no corroboration; yet it is frequently conducive as well to mental improvement, as to mental delight, to obtain like results from different processes. In this view it will be advantageous to the student, to confirm the truth of several of the propositions in this chapter by means of the fluxional analysis. Let the truth enunciated in the above lemma be taken for an example: and let AB be denoted by a , AV by x , BV by $x - a$. Then we shall have

$x - a : x :: x : \frac{x^2}{x - a}$: the third proportional; which is to be a minimum. Hence the fluxion of this fraction will be equal to zero (Flux. art. 51). That is, (Flux. arts. 19 and 30), $\frac{x^2 x - 2axx}{(x - a)^3} = 0$. Consequently, $x^2 - 2ax = 0$, and $x = 2a$,

∴ $AV = 2AB$, as above.

THEOREM XXXII.

The Whole Surface of a Right Cone being given, the Inscribed Sphere is the Greatest when the Slant Side of the Cone is Triple the Radius of its Base.

For, let c and c' be two right cones of equal whole surface, the radii of their respective inscribed spheres being denoted by R and R' ; let the side of the cone c be triple the radius of its base, the same ratio not obtaining in c' ; and let c'' be a cone similar to c , and circumscribed about the same sphere with c' . Then, (by th. 31) surf. $c'' < \text{surf. } c'$; therefore surf. $c'' < \text{surf. } c$. But c'' and c are similar, therefore all the dimensions of c'' are less than the corresponding dimensions of c : and consequently the radius R' of the sphere inscribed in c'' or in c' , is less than the radius R of the sphere inscribed in c , or $R > R'$. Q. E. D.

Cor. The capacity of a right cone being given, the inscribed sphere is the greatest when the side of the cone is triple the radius of its base.

For the capacities of such cones vary as their surfaces (th. 29).

THEOREM XXXIII.

Of all Right Cones of Equal Whole Surface, the Greatest is that whose side is Triple the Radius of its Base: and reciprocally, of all Right Cones of Equal Capacity, that whose Side is Triple the Radius of its Base has the Least Surface.

For, by th. 29, the capacity of a right cone is in the compound ratio of its whole surface and the radius of its inscribed sphere. Therefore, the whole surface being given, the capacity is proportional to the radius of the inscribed sphere: and consequently is a maximum when the radius of the inscribed sphere is such: that is, (th. 32) when the side of the cone is triple the radius of the base*.

Again,

* Here again a similar result may easily be deduced from the method of fluxions. Let the radius of the base be denoted by x , the slant side of the cone by z , its whole surface by a^2 , and 3.141593 by π . Then the circumference of the cone's base will be $2\pi x$, its area πx^2 and the convex surface πxz . The whole surface is, therefore, $=\pi x^2 + \pi xz$: and this being $=a^2$, we have $z = \frac{a^2}{\pi x} - x$. But the altitude of the cone is equal to the square root of the difference of the squares

Again, reciprocally, the capacity being given, the surface is in the inverse ratio of the sphere inscribed: therefore, it is the smallest when that radius is the greatest; that is (th. 32) when the side of the cone is triple the radius of its base.

Q. E. D.

THEOREM XXXIV.

The Surfaces, whether Total or Lateral, of Pyramids Circumscribed about the Same Right Cone, are respectively as their Solidities. And, in particular, the Surface of a Pyramid Circumscribed about a Cone, is to the Surface of that Cone, as the Solidity of the Pyramid is to the Solidity of the Cone; and these Ratios are Equal to those of the Surfaces or the Perimeters of the Bases.

For, the capacities of the several solids are respectively as their bases; and their surfaces are as the perimeters of those bases: so that the proposition may manifestly be demonstrated by a chain of reasoning exactly like that adopted in theorem 11.

of the side and of the radius of the base; that is, it is $= \sqrt{\left(\frac{a^4}{\pi^2 x^2} - \frac{2a^2}{\pi}\right)}$. And this multiplied into $\frac{1}{3}$ of the area of the base, viz. by $\frac{1}{3}\pi x^2$, gives $\frac{1}{3}\pi x^2 \sqrt{\left(\frac{a^4}{\pi^2 x^2} - \frac{2a^2}{\pi}\right)}$, for the capacity of the cone. Now, this being a maximum, its square must be so likewise (Flux. art. 53), that is, $\frac{a^4 x^2 - 2\pi a^2 x^4}{9}$, or, rejecting the denominator, as constant, $a^4 x^2 - 2\pi a^2 x^4$ must be a maximum. This, in fluxions, is $2a^4 x - 8\pi a^2 x^3 = 0$; whence we have $a^2 - 4\pi x^2 = 0$, and consequently $x = \sqrt{\frac{a^2}{4\pi}}$; and $a^2 = 4\pi x^2$. Substituting this value of a^2 for it, in the value of z above given, there results $z = \frac{a^2}{\pi x} - x = \frac{4\pi x^2}{\pi x} - x = 4x - x = 3x$. Therefore, the side of the cone is triple the radius of its base. Or, the square of the altitude is to the square of the radius of the base, as 8 to 1, or, to the square of the diameter of the base, as 2 to 1.

THEOREM

THEOREM XXXV.

The Base of a Right Pyramid being Given in Species, the Capacity of that Pyramid is a Maximum with the same Surface, and on the contrary, the Surface is a Minimum with the Same Capacity, when the Height of One Face is Triple the Radius of the Circle Inscribed in the Base.

LET P and P' be two right pyramids with similar bases, the height of one lateral face of P being triple the radius of the circle inscribed in the base, but this proportion not obtaining with regard to P' : then

1st. If surf. $P = \text{surf. } P'$, $P > P'$.

2dly. If . . . $P = . . . P'$, surf. $P < \text{surf. } P'$.

For, let c and c' be right cones inscribed within the pyramids P and P' : then in the cone c , the slant side is triple the radius of its base, while this is not the case with respect to the cone c' . Therefore, if $c = c'$, surf. $c < \text{surf. } c'$ and if surf. $c = \text{surf. } c'$, $c > c'$ (th. 33).

But, 1st. surf. $P : \text{surf. } c :: \text{surf. } P' : \text{surf. } c'$;
whence, if surf. $P = \text{surf. } P'$, surf. $c = \text{surf. } c'$;
therefore $c > c'$. But $P : c :: P' : c'$. Therefore $P > P'$.

2dly, $P : c :: P' : c'$. Theref. if $P = P'$, $c = c'$: consequently surf. $c < \text{surf. } c'$. But, surf. $P : \text{surf. } c :: \text{surf. } P' : \text{surf. } c'$. Whence, surf. $P < \text{surf. } P'$.

Cor. The regular tetraedon possesses the property of the minimum surface with the same capacity, and of the maximum capacity with the same surface, relatively to all right pyramids with equilateral triangular bases, and, *a fortiori*, relatively to every other triangular pyramid.

THEOREM XXXVI.

A Sphere is to any Circumscribing Solid, Bounded by Plane Surfaces, as the Surface of the Sphere to that of the Circumscribing Solid.

For, since all the planes touch the sphere, the radius drawn to each point of contact will be perpendicular to each respective plane. So that, if planes be drawn through the centre of the sphere and through all the edges of the body, the body will be divided into pyramids whose bases are the respective planes, and their common altitude the radius of the sphere. Hence, the sum of all these pyramids, or the whole circumscribing solid, is equal to a pyramid or a cone whose base

base is equal to the whole surface of that solid, and altitude equal to the radius of the sphere. But the capacity of the sphere is equal to that of a cone whose base is equal to the surface of the sphere, and altitude equal to its radius. Consequently, the capacity of the sphere, is to that of the circumscribing solid, as the surface of the former to the surface of the latter: both having in this mode of considering them, a common altitude.

Q. E. D.

Cor. 1. All circumscribing cylinders, cones, &c. are to the sphere they circumscribe, as their respective surfaces.

For the same proportion will subsist between their indefinitely small corresponding segments, and therefore between their wholes.

Cor. 2. All bodies circumscribing the same sphere, are respectively as their surfaces.

THEOREM XXXVII.

The Sphere is Greater than any Polyedron of Equal Surface.

For, first it may be demonstrated by a process similar to that adopted in theorem 9, that a *regular* polyedron has a greater capacity than any other polyedron of equal surface. Let p , therefore, be a regular polyedron of equal surface to a sphere s . Then p must either circumscribe s , or fall partly within it and partly out of it, or fall entirely within it. The first of these suppositions is contrary to the hypothesis of the proposition, because in that case the surface of p could not be *equal* to that of s . Either the 2d or 3d supposition therefore must obtain; and then each plane of the surface of p must fall either partly or wholly within the sphere s : whichever of these be the case, the perpendiculars demitted from the centre of s upon the planes, will be each less than the radius of that sphere: and consequently the polyedron p must be less than the sphere s , because it has an equal base, but a less altitude.

Q. E. D.

Cor. If a prism, a cylinder, a pyramid, or a cone, be equal to a sphere either in capacity, or in surface; in the first case, the surface of the sphere is less than the surface of any of those solids; in the second, the capacity of the sphere is greater than that of either of those solids.

The theorems in this chapter will suggest a variety of practical examples to exercise the student in computation. A few such are given in the following page.

EXERCISES.

EXERCISES.

Ex. 1. Find the areas of an equilateral triangle, a square, a hexagon, a dodecagon, and a circle, the perimeter of each being 36.

Ex. 2. Find the difference between the area of a triangle whose sides are 3, 4, and 5, and of an equilateral triangle of equal perimeter.

Ex. 3. What is the area of the greatest triangle which can be constituted with two given sides 8 and 11 : and what will be the length of its third side ?

Ex. 4. The circumference of a circle is 12, and the perimeter of an irregular polygon which circumscribes it is 15 : what are their respective areas ?

Ex. 5. Required the surface and the solidity of the greatest parallelepiped, whose length, breadth, and depth, together make 18 ?

Ex. 6. The surface of a square prism is 546 : what is its solidity when a maximum ?

Ex. 7. The content of a cylinder is 169.645968 : what is its surface when a minimum ?

Ex. 8. The whole surface of a right cone is 201.061952 : what is its solidity when a maximum ?

Ex. 9. The surface of a triangular pyramid is 43.30127 : what is its capacity when a maximum ?

Ex. 10. The radius of a sphere is 10. Required the solidities of this sphere, of its circumscribed equilateral cone, and of its circumscribed cylinder.

Ex. 11. The surface of a sphere is 28.274337, and of an irregular polyedron circumscribed about it 35 : what are their respective solidities ?

Ex. 12. The solidity of a sphere, equilateral cone, and Archimedean cylinder, are each 500 : what are the surfaces and respective dimensions of each ?

Ex. 13. If the surface of a sphere be represented by the number 4, the circumscribed cylinder's convex surface and whole surface will be 4 and 6, and the circumscribed equilateral cone's convex and whole surface, 6 and 9 respectively. Show how these numbers are deduced.

Ex. 14. The solidity of a sphere, circumscribed cylinder, and circumscribed equilateral cone, are as the numbers 4, 6, and 9. Required the proof.


PROBLEMS RELATIVE TO THE DIVISION OF FIELDS OR OTHER SURFACES.

PROBLEM I.

To Divide a Triangle into two parts having a Given Ratio,
 $m : n$

1st. By a line drawn from one angle of the triangle.


Make $AD : AB :: m : m + n$; draw CD . So shall $ADC, BDC,$ be the parts required.



Here, evidently, $AD = \frac{m}{m+n} AB, DB = \frac{n}{m+n} AB.$

2dly. By a line parallel to one of the sides of the triangle.

Let ABC be the given triangle, to be divided into two parts, in the ratio of m to n , by a line parallel to the base AB . Make CE to EB as m to n : erect ED perpendicularly to CB , till it meet the semicircle described on CB , as a diameter, in D . Make $CF = CD$: and draw through $F, GF \parallel AB$. So shall GF divide the triangle ABC in the given ratio:



For, $CE : CB :: \frac{CD^2}{CE} :: CD^2 (=CF^2) : CB^2.$ But $CE : EB :: m : n,$


or $CE : CB :: m : m + n,$ by the construction; therefore $CF^2 : CB^2 :: m : m + n.$ And since $\triangle CGF : \triangle CAB :: CF^2 : CB^2$; it follows that $CGF : CAB :: m : m + n,$ as required.

Computation. Since $CB^2 : CF^2 :: m + n : m,$ therefore, $(m + n) CF^2 = m \cdot CB^2$; whence $CF \sqrt{(m + n)} = CB \sqrt{m},$ or $CF = CB \sqrt{\frac{m}{m+n}}.$ In like manner, $CG = CA \sqrt{\frac{m}{m+n}}.$

3dly. By a line parallel to a given line.

Let HI be the line parallel to which a line is to be drawn, so as to divide the triangle ABC in the ratio of m to n .

By case 2d draw GF parallel to AB , so as to divide ABC in the given ratio. Through F draw FE parallel to HI . On CE as a diameter describe a semicircle; draw GD perp. to AC , to cut the semicircle in D . Make $CP = CD$: through P , parallel to EF , draw PQ , the line required.



The

The demonstration of this follows at once from case 2 ; because it is only to divide FCE , by a line parallel to FE , into two triangles having the ratio of FCE , to FCG , that is, of CE , to CG .


Computation. CG and CF being computed, as in case 1, the distances CH , CI being given, and CP being to CQ as CH to CI : the triangles CGF , GPQ , also having a common verticle angle, are to each other, as $CG \cdot CF$ to $CQ \cdot CP$. These products therefore are equal ; and since the factors of the former are known, the latter product is known. We have hence given the ratio of the two lines $CP (=x)$ to $CQ (=y)$ as CH to CI ; say, as p to q ; and their product $= CF \cdot CG$, say, $= ab$: to find x and y . Here we find $x = \sqrt{\frac{abp}{q}}$, $y = \sqrt{\frac{abq}{p}}$. That is,

$$CP = \sqrt{\frac{CF \cdot CG \cdot CH}{CI}} ; CQ = \sqrt{\frac{CF \cdot CG \cdot CI}{CH}}$$

N. B. If the line of division were to be perpendicular to one of the sides, as to CA , the construction would be similar :

CP would be a geometrical mean between CA and $\frac{m}{m+n} cb$, b being the foot of the perpendicular from B upon AC .

4thly. By a line drawn through a given point P .



By any of the former cases draw lm (fig. 1) to divide the triangle ABC , in the given ratio of m to n : bisect cl in r , and through r and m let pass the sides of the rhomboid $crsm$. Make $ca = pe$, which is given, because the point P is given in position : make cd a fourth proportional to ca , cr , cm ; that is, make $ca : cr :: cm : cd$; and let a and d , be two angles of the rhomboid $cabd$, figs. 1 and 2. pe , in figure 2, being drawn parallel to ac , describe on ed as a diameter the semicircle efd , on which set off $ef = ce = ap$: then set off dm or dm' on CA equal to df , and through P and M , P and M' draw the lines

draw the lines $LM, L'M'$, either of which will divide the triangle in the given ratio.—The construction is given in 2 figs. merely to avoid complexness in the diagrams.

The limitations are obvious from the construction: for, the point L must fall between B and c , and the point M between A and c ; AP must also be less than Pb , otherwise ef cannot be applied to the semicircle on ed .

Demon. Because $cr = \frac{1}{2}cl$, the rhomboid $crsm =$ triangle clm , and because $ca : cr :: cm : cd$, we have $ca \cdot cd = cm \cdot cr$, therefore rhomboid $cabd =$ rhomboid $crsm =$ triangle clm . By reason of the parallels CB, bd , and CA, ab , the triangles aLP, dGM, bGP , are similar, and are to each other as the squares of their homologous sides AP, dM, bP : now $ed^2 = ef^2 + df^2$, by construction; and $ed = pb, ef = AP, df = dM$; therefore $pb^2 = AP^2 + dM^2$, or, the triangle PbG taken away from the rhomboid, is equal to the sum of the triangles aPL, dMG , added to the part $capcd$: consequently $CLM = cabd$, as required. By a like process, it may be shown that $aL'P, dG'M', P'BG'$, are similar, and $aL'P + dG'M' = P'BG'$; whence $Pb'dM' = aL'P$, and $CL'M' = cabd$, as required.



Computation. cl, cm , being known, as well as, ca, AP , or $ce, eP, cr = \frac{1}{2}cl$, is known: and hence cd may be found by the proportion $ca : cr :: cm : cd$. Then $cd - ce = ed$, and $\sqrt{ed^2 - ef^2} = \sqrt{ed^2 - AP^2} = df = dM = dM'$. Thus CM is determined. Then we have $\frac{cl \cdot cm}{CM} = CL$.

N. B. When the point is in one of the sides, as at M ; then make $CL \cdot CM \cdot (m + n) = CA \cdot CB \cdot m$, or $CL : CA :: m \cdot CB : (m + n) CM$, and the thing is done.

5thly. By the shortest line possible.

Draw any line PQ dividing the triangle in the given ratio, and so that the summit of the triangle CPQ shall be c the most acute of the three angles of the triangle. Make $CM = CN$, a geometrical mean proportional between CP and CQ ; so shall MN be the shortest line possible dividing the triangle in the given ratio.—The computation is evident.

Demons. Suppose MN to be the shortest line cutting off the given triangle CMN , and $CG \perp MN$. $MN = MG + GN = CG \cdot \cot M + CG \cdot \cot N = CG (\cot M + \cot N)$. But, $\cot M + \cot N = \frac{\cos M}{\sin M} + \frac{\cos N}{\sin N} = \frac{\sin(M+N)}{\sin M \cdot \sin N}$. And equa.



xviii, Analyt. pl. trigonom.) $\sin. M \cdot \sin. N = \frac{1}{2} \cos. (M-N) - \frac{1}{2} \cos. (M+N) = \frac{1}{2} \cos. (M-N) + \frac{1}{2} \cos. c$. Theref. $MN = CG \cdot \frac{\sin. (M+N)}{\frac{1}{2} \cos. (M-N) + \frac{1}{2} \cos. c}$; which expression is a minimum when

its denominator is a maximum; that is, when $\cos (M-N)$ is the greatest possible, which is manifestly when $M-N=0$, or $M=N$, or when the triangle CMN is isosceles. That the isosceles triangle must have the most acute angle for its summit, is evident from the consideration, that since $2\Delta_{CMN} = CG \cdot MN$, MN varies inversely as CG ; and consequently MN is shortest when CG is longest, that is, when the angle c is the most acute.


N. B. A very simple and elegant demonstration to this case is given in Simpson's Geometry: vide the book on Max. and Min. See also another demonstration at case 2d prob. 6th, below.

PROBLEM II.

To Divide a Triangle into Three Parts, having the Ratio of the quantities m, n, p .

1st. By lines drawn from one angle of the triangle to the opposite side.


Divide the side AB , opposite the angle c from whence the lines are to proceed, in the given ratio at D, E ; join CD, CE ; and ACD, DCE, ECB , are the three triangles required. The demonstration is manifest; as is also the computation.



If it be wished that the lines of division be the shortest the nature of the case will admit of, let them be drawn from the most obtuse angle, to the opposite or longest side.



2dly. By lines parallel to one of the sides of the triangle.

Make $CD : DH : HB :: m : n : p$. Erect DE, HI , perpendicularly to CB , till they meet the semicircle described on the diameter CB , in E and I . Make $CF = CE$, and $CK = CI$. Draw GF through F , and LK through K , parallel to AB ; so shall the lines GF and LK , divide the triangle ABC as required,



The demonstration and computation will be similar to those in the second case of prob. 1.

3dly. By lines drawn from a given point on one of the sides.





Let P (fig. 1) be the given point, *a* and *b* the points which divide the side AB in the given ratio of *m*, *n*, *p*: the point P falling between *a* and *b*. Join PC, parallel to which draw *ac*, *bd*, to meet the sides AC, BC, in the points *c* and *d*: join PC, *rd*, so shall the lines *cp*, *pd*, divide the triangle in the given ratio.

In fig. 2, where P falls nearer one of the extremities of AB than both *a* and *b*, the construction is essentially the same; the sole difference in the result is, that the points *c* and *d*, both fall on one side AC of the triangle.

Demon. The lines *ca*, *cb*, divide the triangle into the given ratio, by case 1st. But by reason of the parallel lines *ac*, *pc*, *bd*, $\triangle acc = \triangle acp$, and $\triangle bdc = \triangle bdp$. Therefore, in fig. 1, $aac + acp = aac + acc$ that is, $acp = aac$: and $bbd + bdp = bbd + bdc$, that is, $bdp = bbc$. Consequently, the remainder $cpd = cab$.—In fig. 2, $acp = aac$, and $adp = acb$; therefore $cpd = acp$; and $acb - adp = acb - acb$, that is, $cbpd = cbb$.

Computation. The perpendiculars *cg*, *cd* being demitted, $\triangle acp : \triangle acb :: m : m + n + p :: ap . cg : ab . cd$. Therefore $(m+n+p) ap . cg = m . ab . cd$, and $cg = \frac{m . ab . cd}{(m+n+p) ap}$. The line *cg* being thus known, we soon find *ac*; for $cd : ac :: cg : ac = \frac{ac . cg}{cd} = \frac{m . ab . ac}{(m+n+p) ap}$. Indeed this expression may be deduced more simply; for, since $acb : acp :: ac . ab : ac . ap :: m+n+p : m$, we have $(m+n+p) ac . ap = m . ab . ac$, and $ac = \frac{m . ab . ac}{(m+n+p) ap}$. By a like process is obtained, in fig. 1, $bd = \frac{p . ab . bc}{(m+n+p) pb}$; and, in fig. 2, $ad = \frac{(m+n) ab . ac}{(m+n+p) ap}$.

4thly. By lines drawn from a given point P within the triangle.



Const. Through p and c draw the line cpp , and let the triangle be divided into the given ratio by lines pc , pd , drawn from p to intersect ac , bc , or either of them; according to the method described in case 3 of this problem Through p draw pc , pd , and respectively parallel to them, from p draw the lines pm , pn : join pm , pn ; so shall these lines with fp , divide the triangle in the given ratio.

Demon. The triangles cpm , cpp , are manifestly equal, as are also dpn , dpp ; therefore $cpm = cpc$, and $cpn = cpd$; whence also, in fig. 1, $cnpm = cdpc$, and, in fig. 2, $cbpfn = cbpd$.

Comput. Since $cp \cdot cn = cp \cdot cd$, we have $cn = \frac{cp \cdot cd}{cp}$.

In like manner $cm = \frac{cp \cdot cc}{cp}$.



Remark. It will generally be best to contrive that the *smallest* share of the triangle shall be laid off nearest the vertex c of the triangle, in order to ensure the possibility of the construction. Even this precaution however may sometimes fail, of ensuring the construction by the method above given: when this happens, proceed thus:

By case 1, draw the lines cd , ce , from the vertex c to the opposite side ab , to divide the triangle in the given ratio. Upon ab set off any where mn , so that $mn : ab :: pp$ (the perp. from p on ab) : cp , the altitude of the triangle. If mp and pn are together to be the least possible, then set off $\frac{1}{2} mn$ on each side the point p : so will the triangle mpn be isosceles, and its perimeter (with the given base and area) a minimum.



5thly. By lines, one of which is drawn from a given angle to a given point, which is also the point of concurrence of the other two lines.

Const.



Const. By case 1st draw the lines ca, cb , dividing the triangle in the given ratio, and so that the smaller portions shall lie nearest the angles A and B (unless the conditions of the division require it to be otherwise). From P and a demit upon ac , the perpendiculars fp, ac ; and from P and b , on bc , the perpendiculars pq, bd . Make $cm : ca :: ac : fp$, and $cn : cb :: bd : pq$. Draw pm, pn , which, with cp , will divide the triangle as required.


When the perpendicular from b or from a , upon bc or ac , is longer than the corresponding perpendicular from P , the point n or m will fall further from c than B or A does. Suppose it to be n ; then make $n'e : eb :: ne : ep$, and draw pn' for the line of division.

The demonstration of all this is too obvious to need tracing here.

Comput. The perp. $ca = aa \cdot \sin A$; and $cm = \frac{ca \cdot ac}{fp}$.
 $bd = bb \cdot \sin B$; and $cn = \frac{cb \cdot bd}{pq}$.

6thly. By lines, one of which falls from the given point of concurrence of all three, upon a given side, in a given angle.

Suppose the given angle to be a right angle, and ef the given perpendicular: which will simplify the operation, though the principles of construction will be the same.



Const. Let ca, cb , divide the triangle in the given ratio. Make $fn : cb :: bd : pf$, and $fm : ca :: ac : pf$; and draw pn, pm , thus forming two triangles pfN, pfM , equal to cbB, caA , respectively. If n fall between f and B , and m between A and f , this construction manifestly effects the division. But if one of the points, suppose m , falls beyond the corresponding point A , the line pm intersecting ac in e : then make $m'e : ea :: em : ep$, and draw pm' : so shall pf, pm', pn , divide the triangle as required.


Comput.

Comput. Here ca and bd , are found as in case 5th; and hence $f_N = \frac{CB \cdot bd}{Pf}$; and $f_M = \frac{CA \cdot ac}{Pf}$. Then $PM = \sqrt{(Mf^2 + f_P^2)}$ and $\frac{Pf}{PM} = \sin. M$. Also $180^\circ - (M + A) = MeA$. Then $\sin. MeA : \sin. M : \sin A \propto MA (=Mf - Af) : Ae : Me$. Again $pe = PM - Me$; and lastly $M'e = \frac{Ae \cdot eM}{eP}$

Here also the demonstration is manifest,

7thly. By lines drawn from the angles to meet in a determinate point.

Construc. On one of the sides, as AC , set off AD , so that $AD : AC :: m : m + n + p$. And on the other, as AB , set off BE , so that $BE : BC :: n : m + n + p$. Through D draw DG parallel to AB ; and through E , EH parallel to BC ; to their point of intersection I draw lines AI, BI, CI , which will divide the triangle ABC into the portions required.



Demon. Any triangle whose base is AB , and whose vertex falls in DG parallel to it, will manifestly be to AEC , as AD to AC , or as m to $m + n + p$: so also, any triangle whose base is BC , and whose vertex falls in EH parallel to it, will be to AEC , as BE to BA , that is, as n to $m + n + p$.

Thus we have $AIB : ACB :: m : m + n + p$,
and . . . $BIC : ACB :: n : m + n + p$,
therefore . . . $AIB : BIC :: m : n$.

And the first two proportions give, by composition, $AIB + BIC : ACB :: m + n : m + n + p$; and by division, $ACB - (AIB + BIC) : ACB :: m + n + p - (m + n) : m + n + p$, or $AIC : ACB :: p : m + n + p$, consequently $AIB : BIC : AIC \propto m : n : p$.

Comput. $BE = GI = \frac{n \cdot AB}{m + n + p}$; $EG = \frac{m \cdot BC}{m + n + p}$; angle $BGI = 2$ right angles $- B$. Hence, in the triangle BGI , there are known two sides and the included angle, to find the third side BI .

Remark. When $m = n = p$, the construction becomes simpler. Thus: from the vertex draw CD to bisect AB ; and from B draw BE in like manner to the middle of AC : the point of intersection I of the lines CD, BE , will the point sought.



For, on BE and BE produced, demit, from the angle c and

A, the perpendiculars CI, AK : then the triangles CEI, AEK , are equal in all respects, because $AE=CE, KAE=ICE$, and the angles at E are equal. Hence $AK=CI$. But these are the perpendicular altitudes of the triangles BPC, BPA , which have the common base BP . Consequently those two triangles are equal in area. In a similar manner it may be proved, that $APC=APB$ or CPB : therefore these three triangles are equal to each other, and the lines PA, PB, PC , trisect the $\triangle ABC$.


PROBLEM III.

To Divide a Triangle into Four Parts, having the Proportion of the Quantities m, n, p, q .

This, like the former problems, might be divided into several cases, the consideration of all which would draw us to a very great length, and which is in a great measure unnecessary, because the method will in general be suggested immediately on contemplating the method of proceeding in the analogous case of the preceding problem. We shall therefore only take one case, namely, that in which the lines of division must all be drawn from a given point of one of the sides.

Let P be the given point in the side AB .

Let the points l, m, n , divide the base AB in the given proportion; so will the lines cl, cm, cn , divide the surface of the triangle in the same proportion. Join CP , and parallel to it draw, from l, m, n , the lines Ll, Mm, Nn , to cut the other two sides of the triangle in L, M, N . Draw PL, PM, PN , which will divide the triangle as required.



The demonstration is too obvious to need tracing throughout: for the triangles LlP, LlC , having the same base Ll , and lying between the same two parallels Ll, CP , are equal; to each of these adding the triangle AlL , there results $AlP=AlC$. And in like manner the truth of the whole construction may be shown.


The computation may be conducted after the manner of that in case 3d. prob 2.

PROBLEM IV.

To Divide a Quadrilateral into Two Parts having a Given Ratio, $m : n$.

1st. By a line drawn from any point in the perimeter of the figure,

Construc. From P draw lines PA, PB , to the opposite angles A, B . Through D draw DF parallel to PA , to meet BA produced in F : and through c draw CE parallel to PB to meet AB produced in E .



Divide FE in M, in the given ratio of m to n : join P, M ; so shall the line PM divide the quadrilateral as required.

Demon. That the triangle FPE is equal to the quadrangle ABCD, may be shown by the same process as is used to demonstrate the construction of prob. 36, Geometry ; of which, in fact, this is only a modification. And the line PM evidently divides FPE in the given ratio. But $FPM = ADPM$, and $EPM = BCPM$: therefore PM divides the quadrangle also in the given ratio.

Remark 1. If the line PM cut either of the sides AD, BC, then its position must be changed by a process similar to that described in the 5th and 6th cases of the last problem.


Remark 2. The quadrilateral may be divided into three, four, or more parts, by a similar method, being subject however to the restriction mentioned in the preceding remark.

Remark 3. The same method may obviously be used when the given point P is in one of the angles of the figure.

Comput. Suppose I to be the point of intersection of the sides DC and AB, produced ; and let the part of the quadrilateral laid off towards I, be to the other, as n to m . Then we have $IM = \frac{n(ID \cdot IA - IB \cdot IC)}{(m+n)IP}$. As to the distances DI, AI, (since the angles at A and D, and consequently that at I, are known), they are easily found from the proportionality of the sides of triangles to the sines of their opposite angles.

2dly. By a line drawn parallel to a given line.

Construc. Produce DC, AB, till they meet, as at I. Join DB parallel to which draw CF. Divide AF in the given ratio in H. Through D draw DG parallel to the given line. Make IP a mean proportional between IH, IG ; through P draw PM parallel to GD : so shall PM divide the quadrilateral ABCD as required.



Demon. It is evident, from the transformation of figures, so often resorted to in these problems, that the triangle ADF = quadrilateral AECD (th. 36 Geom.) : and that DH divides the triangle ADF in the given ratio, is evident from prob. 1, case 1. We have only then to demonstrate that the triangle IHD is equal to the triangle IPM, for in that case HDF will manifestly be equal to BCMP. Now, by construction, $IH : IP :: IP : IG ::$ (by the parallels) $IM : ID$; whence, by making the products of the means and extremes equal, we have $ID \cdot IH = IP \cdot IM$; but when


tions (as from 8 sides to 6, from 6 to 4, &c.) if the sides be more than four. Divide the triangle NPS into two parts having the given ratio, by the line PH . In like manner, reduce $ADEFGA$ into an equivalent triangle having H for its vertex, and FE produced for its base; and divide this triangle into the given ratio by a line from H , as HK . The compound line PHK will manifestly divide the whole polygon into two parts having the given ratio. To reduce this to a right line, join PK , and through H draw HM parallel to it; join PM ; so will the right line PM divide the polygon as required, provided M fall between F and E . If it do not, the reduction may be completed by the process described in cases 5th and 6th prob. 2d.

All this is too evident to need demonstration.

Remark. There is a *direct* method of solving this problem, without subdividing the figure: but as it requires the computation of the area, it is not given here.

2dly. By the shortest line possible.

Construc. From any point P' , in one of those two sides of the polygon which, when produced, meet in the most acute angle I , draw a line $P'M'$, to the other of those sides (EF), dividing the polygon in the given ratio. Find the points P and M , so that IP or IM shall be a mean proportional between IP' , IM' ; then will PM be the line of division required.




The demonstration of this is the same as has been already given, at case 5 prob. 1. Those, however, who wish for a proof, independent of the arithmetic of sines, will not be displeased to have the additional demonstration below.


The *shortest* line which, with two other lines given in position includes a given area, will make equal angles with those two lines, or with the segments of them it cuts off from an isosceles triangle.

Let the two triangles ABC , AFF , having the common angle A , be equal in surface, and let the former triangle be isosceles, or have $AB=AC$; then is BC shorter than EF .

First, the oblique base EF cannot pass through D , the middle point of BC , as in the annexed figure. For, drawing CG parallel to AB , to meet EF produced in G . Then the two triangles DBE , DCG are identical, or mutually equal in all respects. Consequently the triangle DCF is less than DCE , and therefore ABC less than AEF .



EF must therefore cut BC in some point H between B and D , and cutting the perp. AD in some point I above D , as in the 2d fig. Upon EF (produced if necessary) demit the perp. AK . Then in the right-angled $\triangle AIK$, the perp. AK is less than the hypotenuse AI , and therefore much more less than the other perp. AD . But, of equal triangles, that which has the greatest perpendicular, has the least base. Therefore the base BC is less than the base EF . Q. E. D.



This series of problems might have been extended much further; but the preceding will furnish a sufficient variety, to suggest to the student the best method to be adopted in almost any other case that may occur. The following practical examples are subjoined by way of exercise.

Ex. 1. A triangular field, whose sides are 20, 18, and 16 chains, is to have a piece of 4 acres in content fenced off from it, by a right line drawn from the most obtuse angle to the opposite side. Required the length of the dividing line, and its distance from either extremity of the line on which it falls?

Ex. 2. The three sides of a triangle are 5, 12, and 13. If two-thirds of this triangle be cut off by a line drawn parallel to the longest side, it is required to find the length of the dividing line, and the distance of its two extremities from the extremities of the longest side.

Ex. 3. It is required to find the length and position of the shortest possible line, which shall divide, into two equal parts, a triangle whose sides are 25, 24, and 7 respectively.

Ex. 4. The sides of a triangle are 6, 8, and 10: it is required to cut off nine-sixteenths of it, by a line that shall pass through the centre of its inscribed circle.

Ex. 5. Two sides of a triangle, which include an angle of 70° , and 14 and 17 respectively. It is required to divide it into three equal parts, by lines drawn parallel to its longest side.

Ex. 6. The base of a triangle is 112.65, the vertical angle $57^\circ 57'$, and the difference of the sides about that angle is 8. It is to be divided into three equal parts, by lines drawn from the angles to meet in a point within the triangle. The lengths of those lines are required

Ex. 7. The legs of a right-angled triangle are 28 and 45. Required the lengths of lines drawn from the middle of the hypotenuse, to divide it into four equal parts.

Ex. 8. The length and breadth of a rectangle are 15 and 9. It is proposed to cut off one-fifth of it, by a line which shall be drawn from a point on the longest side at the distance of 4 from a corner.

Ex. 9. A regular hexagon, each of whose sides is 12, is to be divided into four equal parts, by two equal lines; both passing through the centre of the figure. What is the length of those lines when a minimum?

Ex. 10. The three sides of a triangle are 5, 6, and 7. How may it be divided into four equal parts, by two lines which shall cut each other perpendicularly;

* * The student will find that some of these examples will admit of two answers.



On the Construction of Geometrical Problems.

Problems in Plane Geometry are solved either by means of the modern or algebraical analysis, or of the ancient or geometrical analysis. Of the former, some specimens are given in the Application of Algebra to Geometry, page 369, &c. of this volume. Of the latter, we here present a few examples, premising a brief account of this kind of analysis.

Geometrical analysis is the way by which we proceed from the thing demanded, granted for the moment, till we have connected it by a series of consequences with something anteriorly known, or placed it among the number of principles known to be true.


Analysis

Analysis may be distinguished into two kinds. In the one, which is named by Pappus, contemplative, it is proposed to ascertain the truth or the falsehood of a proposition advanced; the other is referred to the solution of problems, or to the investigation of unknown truths. In the first we assume as true, or as previously existing, the subject of the proposition advanced, and proceed by the consequences of the hypothesis to something known; and if the result be thus found true, the proposition advanced is likewise true. The direct demonstration is afterwards formed, by taking up again, in an inverted order, the several parts of the analysis. If the consequence at which we arrive in the last place is found false, we thence conclude that the proposition analysed is also false. When a *problem* is under consideration, we first suppose it resolved, and then pursue the consequences thence derived till we come to something known. If the ultimate result thus obtained be comprised in what the geometers call data, the question proposed may be resolved: the demonstration (or rather the construction), is also constituted by taking the parts of the analysis in an inverted order. The impossibility of the last result of the analysis, will prove evidently, in this case as well as in the former, that of the thing required.

In illustration of these remarks take the following examples.

Ex. 1. It is required to draw, in a given segment of a circle, from the extremes of the base A and B , two lines AC , BC , meeting at a point c in the circumference, such that they shall have to each other a given ratio, viz. that of M to N .

Analysis. Suppose that the thing is affected, that is to say, that $AC : CB :: M : N$, and let the base AB of the segment be cut in the same ratio in the point E . Then EC , being drawn, will bisect the angle ACB (by th. 83 Geom.); consequently, if the circle be completed, and CE be produced to meet it in F , the remaining circumference will also be bisected in F , or have $FA = FB$, because those arcs are the double measures of equal angles: therefore the point F , as well as E , being given, the point c is also given.




Construction. Let the given base of the segment AB be cut in the point E in the assigned ratio of M to N , and complete the circle; bisect the remaining circumference in F ; join FE , and produce it till it meet the circumference in c : then drawing CA , CB , the thing is done.

Demonstration. Since the arc $FA =$ the arc FB , the angle $ACF =$ angle BCF , by theor. 49 Geom.; therefore $AC : CB ::$

AE : EB, by th. 83. But AE : EB :: M : N, by construction ; therefore AC : CB :: M : N. Q. E. D.

Ex. 2. From a given circle to cut off an arc, such that the sum of m times the sine, and n times the versed sine, may be equal to a given line.

Anal. Suppose it done, and that AEE'B is the given circle, BE'E the required arc, ED its sine, BD its versed sine ; in DA (produced if necessary) take BF and n th part of the given sum ; join PE, and produce it to meet BF \perp to AB or \parallel to ED, in the point F. Then, since $m \cdot ED + n \cdot BD = n \cdot BF = n \cdot PD + n \cdot BD$; consequently $m \cdot ED = n \cdot PD$; hence PD :




ED :: $m : n$. But PD : ED :: (by sim. tri.) PB : BF ; therefore PB : BF :: $m : n$. Now PB is given, and, being at right angles to BF, is also given in position ; therefore the point F is given and consequently PF given in position ; and therefore the point E, its intersection with the circumference of the circle AEE'B, or the arc BE is given. Hence the following

Const. From B, the extremity of any diameter AB of the given circle, draw BM at right angles to AB ; in AB (produced if necessary) take BP an n th part of the given sum ; and on BM take BF so that BF : BP :: $n : m$. Join PF, meeting the circumference of the circle in E and E', and BE or BE' is the arc required.

Demon. From the points E and E' draw ED and E'D' at right angles to AB. Then, since BF : BP :: $n : m$, and (by sim. tri.) BF : BP :: DE : DP ; therefore DE : DP :: $n : m$. Hence $m \cdot DE = n \cdot DP$; add to each $n \cdot BD$, then will $m \cdot DE + n \cdot BD = n \cdot BD + n \cdot DP = n \cdot PB$, or the given sum.

Ex. 3. In a given triangle ABH, to inscribe another triangle abc , similar to a given one, having one of its sides parallel to a line mbn given by position, and the angular points a, b, c , situate in the sides AB, BH, AH, of the triangle ABH respectively.

Analysis. Suppose the thing done, and that abc is inscribed as required. Through any point c in BH draw CD parallel to mbn or to ab , and cutting AB in D ; draw CE parallel to bc , and DE to ac , intersecting each other in E. The triangles DEC, acb , are similar, and



DC : ab :: CE : bc ; also BDC, bab , are similar, and DC : ab ::

BC :


$BC : Bb$. Therefore $BC : CE :: Bb : bc$; and they are about equal angles, consequently B, E, c , are in a right line.

Constr. From any point c in BH , draw CD parallel to nm ; on CD constitute a triangle CDE similar to the given one; and through its angles E draw BE , which produce till it cuts AH in c : through c draw ca parallel to ED and cb parallel to EC ; join ab , then abc is the triangle required, having its side ab parallel to mn , and being similar to the given triangle.

Demon. For, because of the parallel lines ac, DE , and cb, EC , the quadrilaterals $BDEC$ and $Bacb$, are similar; and therefore the proportional lines DC, ab , cutting off equal angles BDC, Bab ; BCD, Bba ; must make the angles EDC, ECD , respectively equal to the angles cab, cba ; while ab is parallel to DC , which is parallel to mn , by construction.

Ex. 4. Given, in a plane triangle, the vertical angle, the perpendicular, and the rectangle of the segments of the base made by that perpendicular; to construct the triangle.

Anal. Suppose ABC the triangle required, BD the given perpendicular to the base AC , produce it to meet the periphery of the circumscribing circle $ABCH$, whose centre is O , in H ; then, by th. 61 Geom. the rectangle $BD \cdot DH = AD \cdot DC$, the given rectangle: hence, since BD is given, DH and BH are given; therefore $BI = HI$ is given: as also $ID = OE$: and the angle EOC is = ABC the given one, because EOC is measured by the arc KC , and ABC by half the arc AKC or by KC . Consequently EC and $AC = 2EC$ are given. Whence this



Construction. Find DH such, that $DB \cdot DH =$ the given rectangle, or find $DH = \frac{AD \cdot DC}{BD}$; then on any right line GF take $FE =$ the given perpendicular, and $EG = DH$; bisect FG in O , and make $EOC =$ the given vertical angle; then will OC cut EC , drawn perpendicular to OE , in c . With centre O and radius OC , describe a circle, cutting CE produced in A : through F parallel to AC draw FB , to cut the circle in B ; join AB, CB , and ABC is the triangle required.

Remark. In a similar manner we may proceed, when it is required to divide a given angle into two parts, the rectangle

angle of whose tangents may be of a given magnitude. See prob. 40, Simpson's Select Exercises.

Note. For other exercises, the student may construct all the problems, except the 24th in the Application of Algebra to Geometry, at page 369, &c. of this volume. And that he may be the better able to trace the relative advantages of the ancient and the modern analysis, it will be advisable that he solve those problems both geometrically and algebraically.

PRACTICAL EXERCISES IN MENSURATION.

QUEST. 1. WHAT difference is there between a floor 28 feet long by 20 broad, and two others, each of half the dimensions : and what do all three come to at 45s. per square, or 100 square feet ?

Ans. diff. 280 sq. feet. Amount 18 guineas.

QUEST. 2. An elm plank is 14 feet 3 inches long, and I would have just a square yard slit off it ; at what distance from the edge must the line be struck ?

Ans. $7\frac{1}{9}$ inches.

QUEST. 3. A ceiling contains 114 yards 6 feet of plastering, and the room 28 feet broad ; what is the length of it ?

Ans. $36\frac{2}{7}$ feet.

QUEST. 4. A common joist is 7 inches deep, and $2\frac{1}{2}$ thick ; but wanting a scantling just as big again, that shall be 3 inches thick ; what will the other dimensions be ?

Ans. $11\frac{2}{3}$ inches.

QUEST. 5. A wooden cistern cost me 3s. 2d. painting within, at 6d. per yard ; the length of it was 102 inches, and the depth 21 inches ; what was the width ?

Ans. $27\frac{1}{4}$ inches.

QUEST. 6. If my court-yard be 47 feet 9 inches square, and I have laid a foot-path with Purbeck stone, of 4 feet wide, along one side of it, what will paving the rest with flints come to, at 6d. per square yard ?

Ans. 5l. 16s. $0\frac{1}{2}$ d.

QUEST. 7. A ladder $26\frac{2}{3}$ feet long, may be so planted, that it shall reach a window 22 feet from the ground on one side
of

of the street ; and by only turning it over, without moving the foot out of its place, it will do the same by a window 14 feet high on the other side ; what is the breadth of the street ?

Ans. 37 feet $9\frac{1}{2}$ inches.

QUEST. 8. The paving of a triangular court, at 18d. per foot, came to 100*l.* ; the longest of the three sides was 88 feet ; required the sum of the other two equal sides ?

Ans. 106·85 feet.

QUEST. 9. There are two columns in the ruins of Persepolis left standing upright : the one is 64 feet above the plain, and the other 50 : in a straight line between these stands an ancient small statue, the head of which is 97 feet from the summit of the higher, and 86 feet from the top of the lower column, the base of which measures just 76 feet to the centre of the figure's base Required the distance between the tops of the two columns ?

Ans. 157 feet nearly.

QUEST. 10. The perambulator, or surveying wheel, is so contrived, as to turn just twice in the length of 1 pole, or $16\frac{1}{2}$ feet ; required the diameter ?

Ans. 2·626 feet.

QUEST. 11. In turning a one-horse chaise within a ring of a certain diameter, it was observed that the outer wheel made two turns, while the inner made but one : the wheels were both 4 feet high ; and supposing them fixed at the distance of 5 feet asunder on the axletree, what was the circumference of the track described by the outer wheel ?

Ans. 62·83 feet.

QUEST. 12. What is the side of that equilateral triangle, whose area cost as much paving at 8d. a foot, as the pallsading the three sides did at a guinea a yard ?

Ans. 72·746 feet.

QUEST. 13. In the trapezium ABCD, are given, $AB = 13$, $BC = 31\frac{1}{2}$, $CD = 24$, and $DA = 18$, also B a right angle ; required the area ?

Ans. 410·122.

QUEST. 14. A roof which is 24 feet 8 inches by 14 feet 6 inches, is to be covered with lead at 8lb. per square foot : what will it come to at 18s. per cwt. ;

Ans. 22*l.* 19s. 10¼*d.*

QUEST. 15. Having a rectangular marble slab, 58 inches by 27, I would have a square foot cut off parallel to the shorter edge ; I would then have the like quantity divided from the remainder parallel to the longer side ; and this alternately repeated, till there shall not be the quantity of a foot left :

left : what will be the dimensions of the remaining piece ?

Ans. 20·7 inches by 6·086.

QUEST. 16. Given two sides of an obtuse-angled triangle, which are 20 and 40 poles ; required the third side, that the triangle may contain just an acre of land ?

Ans. 58·876 or 23·099.

QUEST. 17. The end wall of a house is 24 feet 6 inches in breadth, and 40 feet to the eaves ; $\frac{1}{3}$ of which is 2 bricks thick, $\frac{1}{3}$ more is $1\frac{1}{2}$ brick thick, and the rest 1 brick thick. Now the triangular gable rises 38 courses of bricks, 4 of which usually make a foot in depth, and this is but $4\frac{1}{2}$ inches, or half a brick thick : what will this piece of work come to at 5*l.* 10*s.* per statute rod ?

Ans. 20*l.* 11*s.* 7½*d.*

QUEST. 18. How many bricks will it take to build a wall, 10 feet high, and 500 feet long, of a brick and half thick : reckoning the brick 10 inches long, and 4 courses to the foot in height ?

Ans. 72000.

QUEST. 19. How many bricks will build a square pyramid of 100 feet on each side at the base, and also 100 feet perpendicular height : the dimensions of a brick being supposed 10 inches long, 5 inches broad, and 3 inches thick ?

Ans. 3840000.

QUEST. 20. If, from a right-angled triangle, whose base is 12, and perpendicular 16 feet, a line be drawn parallel to the perpendicular, cutting off a triangle whose area is 24 square feet ; required the sides of this triangle ?

Ans. 6, 8, and 10.

QUEST. 21. The ellipse in Grosvenor-square measures 340 links across the longest way, and 612 the shortest, within the rails : now the walls being 14 inches thick, what ground do they enclose, and what do they stand upon ?

Ans. } enclose 4 ac. 0 r. 6 p.
 } stand on 1760½ sq. feet.

QUEST. 22. If a round pillar, 7 inches over, have 4 feet of stone in it : of what diameter is the column, of equal length, that contains 10 times as much ?

Ans. 22·136 inches.

QUEST. 23. A circular fish-pond is to be made in a garden, that shall take up just half an acre ; what must be the length of the chord that strikes the circle ?

Ans. 27¾ yards.

QUEST. 24. When a roof is of a true pitch, or making a right angle at the ridge, the rafters are nearly $\frac{3}{4}$ of the breadth of the building: now supposing the eaves-boards to project 10 inches on a side, what will the new ripping a house cost, that measures 32 feet 9 inches long, by 22 feet 9 inches broad on the flat, at 15s. per square? Ans. 8*l.* 15*s.* 9 $\frac{1}{2}$ *d.*

QUEST. 25. A cable, which is 3 feet long, and 9 inches in compass, weighs 22*lb.*; what will a fathom of that cable weigh, which measures a foot about? Ans. 78 $\frac{2}{3}$ *lb.*

QUEST. 26. My plumber has put 28*lb.* per square foot into a cistern, 74 inches and twice the thickness of the lead long, 26 inches broad, and 40 deep: he has also put three stays across it within, of the same strength, and 16 inches deep, and reckons 22*s.* per cwt. for work and materials. I, being a mason, have paved him a workshop, 22 feet 10 inches broad, with Purbeck stone, at 7*d.* per foot; and on the balance, I find there is 3*s.* 6*d.* due to him; what was the length of the workshop supposing sheet lead of $\frac{1}{10}$ of an inch thick to weigh 5·899*lb.* the square foot?

Ans. 32 feet, 0 $\frac{3}{4}$ inch.

QUEST. 27. The distance of the centres of two circles, whose diameters are each 50. being given, equal to 30; what is the area of the space enclosed by their circumferences?

Ans. 559·119.

QUEST. 28. If 20 feet of iron railing weigh half a ton, when the bars are an inch and quarter square; what will 50 feet come to at 3 $\frac{1}{2}$ *d.* per *lb.* the bars being $\frac{7}{8}$ of an inch square?

Ans. 20*l.* 0*s.* 2*d.*

QUEST. 29. The area of an equilateral triangle, whose base falls on the diameter, and its vertex in the middle of the arc of a semicircle, is equal to 100: what is the diameter of the semicircle?

Ans. 26·32148.

QUEST. 30. It is required to find the thickness of the lead in a pipe, of an inch and quarter bore, which weighs 14*lb.* per yard in length; the cubic foot of lead weighing 11325 ounces?

Ans. ·20737 inches.

QUEST. 31. Supposing the expense of paving a semicircular plot, at 2*s.* 4*d.* per foot, come to 10*l.*; what is the diameter of it?

Ans. 14·7737 feet.

QUEST.

QUEST. 32. What is the length of a chord which cuts off $\frac{1}{3}$ of the area from a circle whose diameter is 289?

Ans. 278·6716.

QUEST. 33. My plumber has set me up a cistern, and his shop-book being burnt, he has no means of bringing in the charge, and I do not choose to take it down to have it weighed; but by measure he finds it contains $64\frac{3}{10}$ square feet, and that it is precisely $\frac{1}{8}$ of an inch in thickness. Lead was then wrought at 2*l.* per fother of $19\frac{1}{2}$ cwt. It is required from these items to make out the bill, allowing $6\frac{5}{8}$ oz. for the weight of a cubic inch of lead?

Ans. 4*l.* 11*s.* 2*d.*

QUEST. 34. What will the diameter of a globe be, when the solidity and superficial content are expressed by the same number?

Ans. 6.

QUEST. 35. A sack, that would hold 3 bushels of corn, is $22\frac{1}{2}$ inches broad when empty; what will another sack contain, which being of the same length, has twice its breadth, or circumference?

Ans. 12 bushels.

QUEST. 36. A carpenter is to put an oaken curb to a round well, at 8*d.* per foot square: the breadth of the curb is to be $7\frac{1}{2}$ inches, and the diameter within $3\frac{1}{2}$ feet; what will be the expense?

Ans. 5*s.* 2*d.*

QUEST. 37. A gentleman has a garden 100 feet long and 80 feet broad; and a gravel walk is to be made of an equal width half round it; what must the breadth of the walk be to take up just half the ground?

Ans. 25·968 feet.

QUEST. 38. The top of a may-pole, being broken off by a blast of wind, struck the ground at 10 feet distance from the foot of the pole; what was the height of the whole may-pole, supposing the length of the broken piece to be 26 feet?

Ans. 50 feet.

QUEST. 39. Seven men bought a grinding stone, of 60 inches diameter, each paying $\frac{1}{7}$ part of the expense; what part of the diameter must each grind down for his share?

Ans. the 1st 4·4508, 2d 4·8400, 3d 5·3535, 4th 6·0765,
5th 7·2079, 6th 9·3935, 7th 22·6778 inches.

QUEST. 40. A maltster has a kiln, that is 16 feet 6 inches square: but he wants to pull it down, and build a new one, that

QUEST. 50. A cubic foot of brass is to be drawn into wire, of $\frac{1}{40}$ of an inch in diameter; what will the length of the wire be, allowing no loss in the metal?

Ans. 97784·797 yards, or 55 miles 984·797 yards.

QUEST. 51. Of what diameter must the bore of a cannon be, which is cast for a ball of 24lb. weight, so that the diameter of the bore may be $\frac{1}{10}$ of an inch more than that of the ball?

Ans. 5·647 inches.

QUEST. 52. Supposing the diameter of an iron 9lb. ball to be 4 inches, as it is very nearly; it is required to find the diameters of the several balls weighing 1, 2, 3, 4, 6, 12, 18, 24, 32, 36, and 42lb, and the caliber of their guns allowing $\frac{1}{30}$ of the caliber, or $\frac{1}{30}$ of the ball's diameter, for windage.

Answer.

Wt. of ball.	Diameter ball.	Caliber of gun.
1	1·9230	1·9622
2	2·4228	2·4723
3	2·7734	2·8301
4	3·0526	3·1149
6	3·4943	3·5656
9	4·0000	4·0816
12	4·4026	4·4924
18	5·0397	5·1425
24	5·5469	5·6601
32	6·1051	6·2297
36	6·3496	6·4792
42	6·6844	6·8208

QUEST. 53. Supposing the windage of all mortars to be $\frac{1}{30}$ of the caliber, and the diameter of the hollow part of the shell to be $\frac{7}{10}$ of the caliber of the mortar: it is required to determine the diameter and weight of the shell, and the quantity or weight of powder requisite to fill it, for each of the several sorts of mortars, namely, the 13, 10, 8, 5·8, and 4·6 inch mortar.

Answer.

Answer,

Calib. of mort.	Diameter of shell.	Wt. of shell empty.	Wt. of powder.	Wt. of shell filled.
4.6	4.523	8.320	0.583	8.903
5.8	5.703	16.677	1.168	17.845
8	7.867	43.764	3.065	46.829
10	9.833	85.476	5.986	91.462
13	12.783	187.791	13.151	200.942

QUEST. 54. If a heavy sphere, whose diameter is 4 inches, be let fall into a conical glass, full of water, whose diameter is 3, and altitude 6 inches; it is required to determine how much water will run over?

Ans. 26.272 cubic inches, or nearly $\frac{3}{4}$ of a pint.

QUEST. 55. The dimensions of the sphere and cone being the same as in the last question, and the cone only $\frac{1}{2}$ full of water; required what part of the axis of the sphere is immersed in the water?

Ans. .546 parts of an inch.

QUEST. 56. The cone being still the same, and $\frac{1}{2}$ full of water; required the diameter of a sphere which shall be just all covered by the water?

Ans. 2.445996 inches.

QUEST. 57. If a person, with an air balloon, ascend vertically from London. to such a height that he can just see Oxford appear in the horizon; it is required to determine his height above the earth, supposing its circumference to be 25000 miles, and the distance between London and Oxford 49.5933 miles?

Ans. $\frac{311}{10000}$ of a mile, or 547 yards 1 foot.


QUEST. 58. In a garrison there are three remarkable objects A, B, C, the distances of which from one to another are known to be, AB 213, AC 424, and BC 262 yards; I am desirous of knowing my position and distance at a place or station s, from which I observed the angle ASB $13^{\circ} 30'$, and the angle CSB $29^{\circ} 50'$ both by geometry and trigonometry.

Answer,

AS 605.7122;

BS 429.6814;

CS 524.2365.




QUEST. 59. Required the same as in the last question, when the point B is on the other side of AC, supposing AB 9,

AC

AC 12, and BC 6 furlongs ; also the angle ASB $33^{\circ} 45'$, and the angle BSC $22^{\circ} 30'$.

Answer,

AS 10.64, BS 15.64, CS 14.01.



QUEST. 60. It is required to determine the magnitude of a cube of gold, of the standard fineness, which shall be equal to a sum of 480 million of pounds sterling, supposing a guinea to weigh 5 dwts $9\frac{1}{2}$ grains. Ans. 18.691 feet.

QUEST. 61. The ditch of a fortification is 1000 feet long, 9 feet deep, 20 feet broad at bottom, and 22 at top ; how much water will fill the ditch ? Ans. 1158127 gallons nearly.

QUEST. 62. If the diameter of the earth be 7920 miles, and that of the moon 2160 miles : required the ratio of their surfaces, and also of their solidities : supposing them both to be globular, as they are very nearly ?

Ans. the surfaces are as $13\frac{1}{2}$ to 1 nearly ;
and the solidities as $49\frac{1}{2}$ to 1 nearly.





Qn

37

499

1822

v.1



